

PACKAGE LEAFLET: INFORMATION FOR THE USER

Abavalex 10 mg film-coated tablets
Abavalex 25 mg film-coated tablets
Abavalex 50 mg film-coated tablets
Abavalex 75 mg film-coated tablets
Abavalex 100 mg film-coated tablets

brivaracetam

Read all of this leaflet carefully before you or your child start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Abavalex is and what it is used for
2. What you need to know before you take Abavalex
3. How to take Abavalex
4. Possible side effects
5. How to store Abavalex
6. Contents of the pack and other information

1. What Abavalex is and what it is used for

What Abavalex is

Abavalex contains the active substance brivaracetam. It belongs to a group of medicines called 'anti-epileptics'. These medicines are used to treat epilepsy.

What Abavalex is used for

- Abavalex is used in adults, adolescents and children from 2 years of age.
- It is used to treat a type of epilepsy that has partial seizures with or without a secondary generalisation.
- Partial seizures are fits that start by only affecting one side of the brain. These partial seizures can spread and extend to larger areas on both sides of the brain – this is called a 'secondary generalisation'.
- You have been given this medicine to lower the number of fits (seizures) you have.
- Abavalex is used together with other medicines for epilepsy.

2. What you need to know before you take Abavalex

Do not take Abavalex if:

- you are allergic to brivaracetam, other similar chemical compounds as levetiracetam or piracetam or any of the other ingredients of this medicine (listed in section 6). If you are not sure, talk to your doctor or pharmacist before taking Abavalex.

- you have ever developed as severe skin rash or skin peeling, blistering and/or mouth sores after taking Abavalex.
Serious skin reactions including Stevens-Johnson syndrome have been reported in association with Abavalex treatment. Stop using Abavalex and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

Warnings and Precautions

Talk to your doctor or pharmacist before taking Abavalex if:

- you have thoughts of harming or killing yourself. A small number of people being treated with anti-epileptic medicines such as Abavalex have had thoughts of harming or killing themselves. If you have any of these thoughts at any time, contact your doctor immediately.
- you have liver problems – your doctor may need to adjust your dose.

Children

Abavalex is not recommended for use in children under 2 years of age.

Other medicines and Abavalex

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, tell your doctor if you are taking any of the following medicines – this is because your doctor may need to adjust your Abavalex dose:

- Rifampicin – a medicine used to treat bacterial infections.
- St John's wort (also known as Hypericum perforatum) – a herbal medicine used to treat depression and anxiety as well as other conditions.

Abavalex with alcohol

Combining this medicine with alcohol is not recommended. If you drink alcohol while taking Abavalex the negative effects of alcohol may be increased.

Pregnancy and breast-feeding

Fertile women should discuss the use of contraceptives with the doctor.

If you are pregnant or breast-feeding, think you may be pregnant or planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

It is not recommended to take Abavalex if you are pregnant, as the effects of Abavalex on pregnancy and the unborn baby are not known.

It is not recommended to breast-feed your baby while taking Abavalex, as Abavalex passes into breast-milk.

Do not stop treatment without talking to your doctor first. Stopping treatment could increase your seizures and harm your baby.

Driving and using machines

You may feel sleepy, dizzy or tired while taking Abavalex. These effects are more likely at the start of the treatment or after a dose increase. Do not drive, cycle or use any tools or machines until you know how the medicine affects you.

Abavalex contains lactose and sodium

Abavalex film-coated tablets contain:

- lactose (a type of sugar) – If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.
- sodium – This medicine contains less than 1 mmol sodium (23mg) per tablet, that is to say essentially ‘sodium free’.

3. How to take Abavalex

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. Other form(s) of brivaracetam may be more suitable for certain patients e.g. children (if tablets cannot be swallowed in whole, or if the dose cannot be met with the use of whole tablets for example); ask your doctor or pharmacist.

You will take Abavalex together with other medicines for epilepsy.

How much to take

Your doctor will work out the right daily dose for you. Take the daily dose in two equal divided doses, approximately 12 hours apart.

Adolescents and children weighing 50 kg or more, and adults

The recommended dose is from 25 mg to 100 mg taken twice a day. Your doctor may then decide to adjust your dose to find the best dose for you.

Adolescents and children weighing from 20 kg to less than 50 kg

The recommended dose is from 0.5 mg to 2 mg for each kg of bodyweight, taken twice a day. Your doctor may then decide to adjust your dose to find the best dose for you.

Children weighing from 10 kg to less than 20 kg

The recommended dose is from 0.5 mg to 2.5 mg for each kg of bodyweight, taken twice a day. Your child’s doctor may then decide to adjust your child’s dose to find the best dose for your child.

People with liver problems

If you have problems with your liver:

- As an adolescent or child weighing 50 kg or more, or as an adult, the maximum dose you will take is 75 mg twice a day.
- As an adolescent or child weighing from 20 kg to less than 50 kg, the maximum dose you will take is 1.5 mg for each kg of bodyweight twice a day.
- As a child weighing from 10 kg to less than 20 kg, the maximum dose your child will take is 2 mg for each kg of bodyweight twice a day.

How to take Abavalex tablets

Swallow the tablets with an appropriate amount of liquid. It is recommended to swallow the tablets whole since the active substance has a bitter taste. The medicine may be taken with or without food.

How long to take Abavalex for

Abavalex is a long-term treatment – keep taking Abavalex until your doctor tells you to stop.

If you take more Abavalex than you should

If you have taken more Abavalex than you should, talk to your doctor. You may feel dizzy and sleepy. You may also have any of the following symptoms: feeling sick, a feeling of 'spinning', problems of keeping your balance, anxiety, feeling very tired, irritability, being aggressive, not being able to sleep, depression, thoughts or attempts of harming or killing yourself.

If you forget to take Abavalex

If you miss a dose, take it as soon as you remember. Then take your next dose at the time you would normally take it.

Do not take a double dose to make up for a forgotten dose. If you are not sure what to do, ask your doctor or pharmacist.

If you stop taking Abavalex

Do not stop taking this medicine unless your doctor tells you to. This is because stopping treatment could increase the number of fits you have.

If your doctor asks you to stop taking this medicine, they will lower your dose gradually. This helps to stop your fits coming back or getting worse.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Very common: may affect more than 1 in 10 people

- feeling sleepy or dizzy

Common: may affect up to 1 in 10 people

- flu
- feeling very tired (fatigue)
- convulsion, a feeling of 'spinning' (vertigo)
- feeling and being sick, constipation
- depression, anxiety, not being able to sleep (insomnia), irritability
- infections of the nose and throat (such as the 'common cold'), cough
- decreased appetite

Uncommon: may affect up to 1 in 100 people

- allergic reactions
- abnormal thinking and/or loss of touch with reality (psychotic disorder), being aggressive, nervous excitement (agitation)
- thoughts or attempts of harming or killing yourself: tell your doctor straight away
- a decrease in white blood cells (called 'neutropenia') - shown in blood tests

Not known: frequency cannot be estimated from the available data

- a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome)

Additional side effects in children

Common: may affect up to 1 in 10 people

- restlessness and hyperactivity (psychomotor hyperactivity)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V*](#). By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Abavalex

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton box and blister after EXP:. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Abavalex contains

The active substance is brivaracetam.

Each film-coated tablet contains 10 mg, 25 mg, 50 mg, 75 mg, or 100 mg brivaracetam.

The other ingredients are:

Core

Lactose monohydrate, Croscarmellose sodium (E 468), Lactose (see section 2 „Abavalex contains lactose and sodium”), Magnesium stearate (E 470b)

Film-coating

- 10 mg film-coated tablets: poly(vinyl alcohol) (E 1203), titanium dioxide (E 171), macrogol 3350 (E 1521), talc (E 553b).
- 25 mg film-coated tablets: poly(vinyl alcohol) (E 1203), titanium dioxide (E 171), macrogol 3350 (E 1521), talc (E 553b), Iron oxide black (E 172), iron oxide yellow (E 172).
- 50 mg film-coated tablets: poly(vinyl alcohol) (E 1203), titanium dioxide (E 171), macrogol 3350 (E 1521), talc (E 553b), iron oxide yellow (E 172), iron oxide red (E 172).
- 75 mg film-coated tablets: poly(vinyl alcohol) (E 1203), titanium dioxide (E 171), macrogol 3350 (E 1521), talc (E 553b), iron oxide red (E 172), Iron oxide black (E 172), iron oxide yellow (E 172).
- 100 mg film-coated tablets: poly(vinyl alcohol) (E 1203), titanium dioxide (E 171), macrogol 3350 (E 1521), talc (E 553b), iron oxide yellow (E 172), Iron oxide black (E 172).

What Abavalex looks like and contents of the pack

Abavalex 10 mg are white to off-white, round and biconvex film-coated tablets of 5.1 mm in diameter.

Abavalex 25 mg are grey, oval and biconvex film-coated tablets of 9.0 mm in length.

Abavalex 50 mg are yellow, oval and biconvex film-coated tablets of 11.8 mm in length.
Abavalex 75 mg are pale pink, oval and biconvex film-coated tablets of 13.1 mm in length.
Abavalex 100 mg are beige, oval and biconvex film-coated tablets of 14.6 mm in length.

Abavalex film-coated tablets are supplied in opaque blister packs.
Blister packs contain 14, 20, 56, 60, 70 and 168 film-coated tablets

Not all pack sizes may be marketed.

Marketing authorisation holder

[To be completed nationally.]

Manufacturer

[To be completed nationally.]

This medicinal product is authorised in the Member States of the EEA under the following names:

[To be completed nationally]

This leaflet was last revised in 2025-08-27