

Package leaflet: Information for the user

Amlodipin Accord 5 mg tablets Amlodipin Accord 10 mg tablets Amlodipine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Amlodipin Accord is and what it is used for
2. What you need to know before you take Amlodipin Accord
3. How to take Amlodipin Accord
4. Possible side effects
5. How to store Amlodipin Accord
6. Contents of the pack and other information

1. What Amlodipin Accord is and what it is used for

Amlodipin Accord contains the active substance amlodipine which belongs to a group of medicines called calcium antagonists.

Amlodipin Accord is used to treat high blood pressure (hypertension) or a certain type of chest pain called angina, a rare form of which is Prinzmetal's or variant angina.

In patients with high blood pressure this medicine works by relaxing blood vessels, so that blood passes through them more easily. In patients with angina, Amlodipin Accord works by improving blood supply to the heart muscle which then receives more oxygen and as a result chest pain is prevented. This medicine does not provide immediate relief of chest pain from angina.

2. What you need to know before you take Amlodipin Accord

Do not take Amlodipin Accord

- If you are allergic (hypersensitive) to amlodipine, or any of the other ingredients of this medicine listed in section 6, or to any other calcium antagonists. This may be itching, reddening of the skin or difficulty in breathing.
- If you have severe low blood pressure (hypotension)
- If you have narrowing of the aortic heart valve (aortic stenosis) or cardiogenic shock (a condition where your heart is unable to supply enough blood to the body).
- If you suffer from heart failure after a heart attack

Warnings and precautions

Talk to your doctor or pharmacist before taking Amlodipin Accord. You should inform your doctor if you have or have had any of the following conditions:

- Recent heart attack
- Heart failure
- Severe increase in blood pressure (Hypertensive crisis)
- Liver disease
- You are elderly and your dose needs to be increased

Children and adolescents

Amlodipin Accord has not been studied in children under the age of 6 years. Amlodipin Accord should only be used for hypertension in children and adolescents from 6 years to 17 years of age (see section 3).

For more information, talk to your doctor.

Other medicines and Amlodipin Accord

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Amlodipin Accord may affect or be affected by other medicines, such as:

- ketoconazole, itraconazole (anti-fungal medicines)
- ritonavir, indinavir, nelfinavir (so called protease inhibitors used to treat HIV)
- rifampicin, erythromycin, clarithromycin (antibiotics)
- hypericum perforatum (St. John's Wort)
- verapamil, diltiazem (heart medicines)
- dantrolene (infusion for severe body temperature abnormalities)
- tacrolimus, sirolimus, temsirolimus, and everolimus (medicines used to alter the way your immune system works)
- simvastatin (cholesterol lowering medicine)
- cyclosporine (an immunosuppressant)

Amlodipin Accord may lower your blood pressure even more if you are already taking other medicines to treat your high blood pressure.

Amlodipin Accord with food and drink

Grapefruit juice and grapefruit should not be consumed by people who are taking Amlodipin Accord. This is because grapefruit and grapefruit juice can lead to an increase in the blood levels of the active ingredient amlodipine, which can cause an unpredictable increase in the blood pressure lowering effect of Amlodipin Accord.

Pregnancy and breast-feeding

Pregnancy

The safety of amlodipine in human pregnancy has not been established. If you think you might be pregnant, or are planning to get pregnant, you must tell your doctor before you take Amlodipin Accord.

Breast-feeding

Amlodipine has been shown to pass into breast milk in small amounts. If you are breast-feeding or about to start breast-feeding you must tell your doctor before taking Amlodipin Accord.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Amlodipin Accord may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Amlodipin Accord contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Amlodipin Accord

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended initial dose is 5 mg once daily. The dose can be increased to 10 mg once daily.

This medicine can be used before or after food and drinks. You should take this medicine at the same

time each day with a drink of water. Do not take Amlodipin Accord with grapefruit juice.

Use in children and adolescents

For children and adolescents (6-17 years old), the recommended usual starting dose is 2.5 mg a day. The maximum recommended dose is 5 mg a day. Amlodipine 2.5 mg is not currently available and the 2.5 mg dose cannot be obtained with Amlodipin Accord 5 mg tablets as these tablets are not manufactured to break into two equal halves.

It is important to keep taking the tablets. Do not wait until your tablets are finished before seeing your doctor.

If you take more Amlodipin Accord than you should

Taking too many tablets may cause your blood pressure to become low or even dangerously low. You may feel dizzy, lightheaded, faint or weak. If blood pressure drop is severe enough shock can occur.

Excess fluid may accumulate in your lungs (pulmonary oedema) causing shortness of breath that may develop up to 24-48 hours after intake.

Your skin could feel cool and clammy and you could lose consciousness. Seek immediate medical attention if you take too many Amlodipin Accord.

If you forget to take Amlodipin Accord

Do not worry. If you forget to take a tablet, leave out that dose completely. Take your next dose at the right time. Do not take a double dose to make up for a forgotten dose.

If you stop taking Amlodipin Accord

Your doctor will advise you how long to take this medicine. Your condition may return if you stop using this medicine before you are advised.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Visit your doctor **immediately** if you experience any of the following side effects after taking this medicine.

- Sudden wheeziness, chest pain, shortness of breath or difficulty in breathing
- Swelling of eyelids, face or lips
- Swelling of the tongue and throat which causes great difficulty breathing
- Severe skin reactions including intense skin rash, hives, reddening of the skin over your whole body, severe itching, blistering, peeling and swelling of the skin, inflammation of mucous membranes (Stevens Johnson Syndrome, toxic epidermal necrolysis) or other allergic reactions
- Heart attack, abnormal heart beat
- Inflamed pancreas which may cause severe abdominal and back pain accompanied with feeling very unwell

The following **very common side-effect** has been reported. If this causes you problems or if it **lasts for more than one week**, you should **contact your doctor**.

Very common: may affect more than 1 in 10 people

- oedema (Fluid retention)

The following **common side effects** have been reported. If any of these cause you problems or if they **last for more than one week**, you should **contact your doctor**.

Common: may affect up to 1 in 10 people

- Headache, dizziness, sleepiness (especially at the beginning of treatment)
- Palpitations (awareness of your heart beat), flushing
- Abdominal pain, feeling sick (nausea)
- Altered bowel habits, diarrhoea, constipation, indigestion,
- Tiredness, weakness
- Visual disturbances, double vision
- Muscle cramps
- Ankle swelling

Other side-effects that have been reported include the following list. If any of these get serious, or if you notice any side-effects not listed in this leaflet, please tell your doctor or pharmacist.

Uncommon: may affect up to 1 in 100 people

- Mood changes, anxiety, depression, sleeplessness
- Trembling, taste abnormalities, fainting
- Numbness or tingling sensation in your limbs; loss of pain sensation
- Ringing in the ears
- Low blood pressure
- Sneezing/running nose caused by inflammation of the lining of the nose (rhinitis)
- Cough
- Dry mouth, vomiting (being sick)
- Hair loss, increased sweating, itchy skin, red patches on skin, skin discolouration
- Disorder in passing urine, increased need to urinate at night, increased number of times of passing urine
- Inability to obtain an erection; discomfort or enlargement of the breasts in men
- Pain, feeling unwell
- Joint or muscle pain, back pain
- Weight increase or decrease

Rare: may affect up to 1 in 1,000 people

- Confusion

Very rare: may affect up to 1 in 10,000 people

- Decreased numbers of white blood cells, decrease in blood platelets which may result in unusual bruising or easy bleeding
- Excess sugar in blood (hyperglycaemia)
- A disorder of the nerves which can cause muscular weakness, tingling or numbness
- Swelling of the gums
- Abdominal bloating (gastritis)
- Abnormal liver function, inflammation of the liver (hepatitis), yellowing of the skin (jaundice), liver enzyme increase which may have an effect on some medical tests
- Increased muscle tension
- Inflammation of blood vessels, often with skin rash
- Sensitivity to light
- Disorders combining rigidity, tremor, and/or movement disorders

Not known: frequency cannot be estimated from the available data

- Trembling, rigid posture, mask-like face, slow movements and a shuffling, unbalanced walk

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine. [To be completed nationally]

5. How to store Amlodipin Accord

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after “EXP”. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Amlodipin Accord contains

Amlodipin Accord 5 mg tablets:

The active substance is amlodipine. Each tablet contains 5 mg amlodipine (as besilate).

The other ingredients are: microcrystalline cellulose, sodium starch glycolate, magnesium stearate, disodium hydrogen citrate, crospovidone, croscarmellose sodium.

Amlodipin Accord 10 mg tablets:

The active substance is amlodipine. Each tablet contains 10 mg amlodipine (as besilate).

The other ingredients are: microcrystalline cellulose, sodium starch glycolate, magnesium stearate, disodium hydrogen citrate, crospovidone, croscarmellose sodium.

What Amlodipin Accord looks like and contents of the pack

Amlodipin Accord 5 mg tablets:

White to off white, approximately 6.6 mm round, biconvex tablets.

Amlodipin Accord 10 mg tablets:

White to off white, approximately 8.5 mm round, biconvex tablets

Pack sizes

Blister pack:

5mg: 10, 14, 20, 28, 30, 50, 60, 90, 98 and 100 tablets

10mg: 10, 14, 20, 28, 30, 50, 60, 90, 98 and 100 tablets

HDPE bottle:

250,500 and 1000 tablets (for hospital or dose dispensing use only)

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Accord Healthcare Limited
Sage House, 319 Pinner Road,
North Harrow, Middlesex, HA1 4HF,
United Kingdom

Manufacturer

Accord Healthcare B.V.,
Winthontlaan 200,
3526 KV Utrecht,

The Netherlands

Accord Healthcare Polska Sp.z o.o.,
ul. Lutomierska 50,95-200 Pabianice, Poland

This medicinal product is authorised in the Member States of the EEA under the following names:

Name of the Member State	Name of the medicinal product
Austria	Amlodipin Accord 5/10 mg Tabletten
Cyprus	Amlodipine Accord 5 mg/10 mg Tablets
Czech Republic	Amlodipin Accord 5/10 mg Tablety
Denmark	Amlodipin Accord 5/10 mg Tabletter
Estonia	Amlodipine Accord 5/10 mg
Finland	Amlodipin Accord 5 mg/10 mg tabletti
Ireland	Amlodipine 5 mg/10 mg Tablets
Italy	AMLODIPINA ACCORD 5 /10 MG COMPRESSE
Lithuania	Amlodipine Accord 5/10 mg tabletės
Latvia	Amlodipine Accord 5/10 mg tablets
Netherlands (The)	Amlodipine (als besilaat) Accord 5/10 mg tabletten
Norway	Amlodipin Accord
Poland	Almiden
Portugal	Amlodipina Accord
Romania	Almiden 5 mg/10 mg comprimate
Spain	Amlodipine Accord 5 mg/10 mg comprimidos
Sweden	Amlodipin Accord 5/10 mg tabletter

This leaflet was last revised in
2022-09-01