

Summary of risk management plan for Apixaban Reddy

This is a summary of the risk management plan (RMP) for Apixaban Reddy. The RMP details important risks of Apixaban Reddy, how more information will be obtained about Apixaban Reddy's risks and uncertainties (missing information).

Apixaban Reddy's summary of product characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how Apixaban Reddy should be used.

Important new concerns or changes to the current ones will be included in updates of Apixaban Reddy's RMP.

I. The medicine and what it is used for

Apixaban Reddy is authorised for the following indications (see SmPC for the full indication):

Adults:

- for prevention of venous thromboembolic events (VTE) in adult patients who have undergone elective hip or knee replacement surgery (Apixaban Reddy 2.5).
- for prevention of stroke and systemic embolism in adult patients with non-valvular atrial fibrillation (NVAf), with one or more risk factors, such as prior stroke or transient ischaemic attack (TIA); age $\geq$ 75 years; hypertension; diabetes mellitus; symptomatic heart failure (NYHA Class $\geq$ II) (Apixaban Reddy 2.5 and 5 mg).
- Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE), and prevention of recurrent DVT and PE in adults (see section 4.4 for haemodynamically unstable PE patients) (Apixaban Reddy 2.5 and 5 mg).

Paediatric population

- Treatment of venous thromboembolism (VTE) and prevention of recurrent VTE in paediatric patients from 28 days to less than 18 years of age.

It contains apixaban as the active substance and it is given orally.

II. Risks associated with the medicine and activities to minimise or further characterise the risks

Important risks of Apixaban Reddy, together with measures to minimise such risks and the proposed studies for learning more about Apixaban Reddy's risks, are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals;
- Important advice on the medicine's packaging;
- The authorised pack size — the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly;
- The medicine's legal status — the way a medicine is supplied to the patient (e.g. with or without prescription) can help to minimise its risks.

Together, these measures constitute *routine risk minimisation* measures.

In addition to these measures, information about adverse reactions is collected continuously and regularly analysed, including PSUR assessment, so that immediate action can be taken as necessary. These measures constitute *routine pharmacovigilance activities*.

If important information that may affect the safe use of Apixaban Reddy is not yet available, it is listed under 'missing information' below.

II.A. List of important risks and missing information

Important risks of Apixaban Reddy are risks that need special risk management activities to further investigate or minimise the risk, so that the medicinal product can be safely taken. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of Apixaban Reddy. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g. on the long-term use of the medicine).

Summary of safety concerns	
Important identified risks	• Bleeding
Important potential risks	• Liver injury • Potential risk of bleeding or thrombosis due to overdose or underdose
Missing information	• Use in patients with severe renal impairment

II.B. Summary of important risks

Important identified risk: Bleeding	
Evidence for linking the risk to the medicine	The risk of bleeding associated with apixaban has been comprehensively evaluated in the nonclinical and clinical apixaban programmes. The most clinically significant treatment-related ARs associated with apixaban are bleeding ARs. The majority of bleeding-related events were non-serious and mild to moderate in severity. A bleeding event can be serious if it occurs in a critical anatomical site such as in the brain. Intracranial bleeding can be fatal. Low rates of intracranial bleedings and fatal bleeding were reported. The overall bleeding risk of apixaban was found to be similar to ASA and superior to warfarin in the non-valvular AF programme, and superior to enoxaparin/warfarin in VTE treatment patients.
Risk factors and risk groups	Concurrent use of other anticoagulants or antiplatelet therapies. Patient characteristics: comorbid conditions (e.g., congenital or acquired bleeding disorders; active ulcerative gastrointestinal disease; bacterial endocarditis; thrombocytopenia; platelet disorders; history of haemorrhagic stroke; severe uncontrolled

Important identified risk: Bleeding	
	hypertension; and recent brain, spinal, or ophthalmological surgery). Past medical history (e.g., previous stroke, prior GI bleeding) Co-administration of strong inhibitors of both CYP3A4 and P-glycoprotein (P-gp) (e.g., azole antifungals, protease inhibitors) may increase apixaban blood concentration and risk of bleeding. Therefore, co-administration of apixaban with strong inhibitors of both CYP3A4 and P-gp is not recommended. Orthopaedic VTE Prevention indication Patient characteristics: age > 75 years old When neuraxial anaesthesia (spinal/epidural anaesthesia) or spinal/epidural puncture is employed, patients treated with antithrombotic agents for prevention of thromboembolic complications are at risk of developing an epidural or spinal haematoma which can result in long-term or permanent paralysis. The risk of these events may be increased by the post-operative use of indwelling epidural catheters or the concomitant use of medicinal products affecting haemostasis. The risk may also be increased by traumatic or repeated epidural or spinal puncture. VTE Treatment indication Co-administration of strong inducers of both CYP3A4 and P-gp may lead to a reduction in apixaban exposure and is not recommended for the treatment of DVT and PE. In a clinical study in atrial fibrillation patients, diminished efficacy and a higher risk of bleeding were observed with co-administration of apixaban with strong inducers of both CYP3A4 and P-gp compared with using apixaban alone.
Risk minimisation measures	<u>Routine risk minimisation measures:</u> SmPC section 4.2, 4.3, 4.4, 4.5, 4.8, and 4.9 <u>Other routine risk minimisation measures beyond the Product Information:</u> Prescription-only medicine <u>Additional risk minimisation measures:</u> Patient Card

Important potential risk: Liver injury	
Evidence for linking the risk to the medicine	Across the apixaban clinical program, there have been infrequent reports of liver-related AEs, SAEs, and laboratory abnormalities. In the VTE prevention orthopaedic population,

Important potential risk: Liver injury	
	the majority of events were post-operative transient elevations of ALT, AST, total bilirubin, and/or ALP that either resolved while study drug continued or during follow-up period. In the AF indication, the low frequency of LFT elevations and liver-related safety events is clinically important, and supports the favourable safety profile of apixaban for this indication. In VTE Treatment and Prevention of Recurrent VTE indication, most patients who experienced hepatic enzyme elevation were asymptomatic, however, some patients experienced symptoms depending on the severity of the condition.
Risk factors and risk groups	Prior hepatitis, cirrhosis, fatty liver, alcohol consumption, poor nutrition, co-existing chronic disease, co-administration of hepatically metabolized drugs (e.g., statins), medication overdose, hypoperfusion, transfusion, halogen-anaesthetics, analgesics, hepatotoxic antibiotics, autoimmune disease (autoimmune hepatitis), viruses (primarily HAV, HBV, HCV), hereditary conditions (e.g., Wilson's disease).
Risk minimisation measures	<u>Routine risk minimisation measures:</u> SmPC section 4.2, 4.3, 4.4, and 4.8 <u>Other routine risk minimisation measures beyond the Product Information:</u> Prescription-only medicine <u>Additional risk minimisation measures:</u> None
Important potential risk: Potential risk of bleeding or thrombosis due to overdose or under dose	
Evidence for linking the risk to the medicine	Although post-marketing data has shown that medication errors occur infrequently, overdose as the most prevalent medication error has potentially serious consequences because of the increased risk of bleeding. The majority of events reported under the medication errors HGLT for apixaban in pivotal studies were SAEs. The vast majority of cases reporting overdose, accidental overdose, intentional overdose or accidental exposure were asymptomatic. There was a single fatal outcome as a consequence of intentional suicidal overdose with phenazepam and alcohol.
Risk factors and risk groups	Risk factors include complex/ unclear patient information, packaging, and product label, and use of the product in emergency situations

Important potential risk: Liver injury	
Risk minimisation measures	<u>Routine risk minimisation measures:</u> SmPC section 4.2 and 4.9 <u>Other routine risk minimisation measures beyond the Product Information:</u> Prescription-only medicine <u>Additional risk minimisation measures:</u> None
Missing information: Use in patients with severe renal impairment	
Risk minimisation measures	<u>Routine risk minimisation measures:</u> SmPC section 4.2, 4.4, and 5.2 <u>Other routine risk minimisation measures beyond the Product Information:</u> Prescription-only medicine <u>Additional risk minimisation measures:</u> None

II.C. Post-authorisation development plan

II.C.1 Studies which are conditions of the marketing authorisation

There are no studies which are conditions of the marketing authorisation or specific obligation of Apixaban. Reddy.

II.C.2 Other studies in post-authorisation development plan

There are no studies required for Apixaban Reddy.