

PACKAGE LEAFLET

Package leaflet: Information for the patient

Apremilast Anabiosis 10 mg film-coated tablets
Apremilast Anabiosis 20 mg film-coated tablets
Apremilast Anabiosis 30 mg film-coated tablets

apremilast

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Apremilast Anabiosis is and what it is used for
2. What you need to know before you take Apremilast Anabiosis
3. How to take Apremilast Anabiosis
4. Possible side effects
5. How to store Apremilast Anabiosis
6. Contents of the pack and other information

1. What Apremilast Anabiosis is and what it is used for

Apremilast Anabiosis contains the active substance ‘apremilast’. This belongs to a group of medicines called phosphodiesterase 4 inhibitors, which help to reduce inflammation.

Apremilast Anabiosis is used to treat adults with the following conditions:

- **Active psoriatic arthritis** – if you cannot use another type of medicines called ‘Disease-Modifying Antirheumatic Drugs’ (DMARDs) or when you have tried one of these medicines and it did not work.
- **Moderate to severe chronic plaque psoriasis** – if you cannot use one of the following treatments or when you have tried one of these treatments and it did not work:
 - phototherapy – a treatment where certain areas of the skin are exposed to ultraviolet light
 - systemic therapy – a treatment that affects the entire body rather than just one local area, such as ‘ciclosporin’, ‘methotrexate’ or ‘psoralen’.
- **Behçet’s disease (BD)** – to treat the mouth ulcers which is a common problem for people with this illness.

What psoriatic arthritis is

Psoriatic arthritis is an inflammatory disease of the joints, usually accompanied by psoriasis, an inflammatory disease of the skin.

What plaque psoriasis is

Psoriasis is an inflammatory disease of the skin, which can cause red, scaly, thick, itchy, painful patches on your skin and can also affect your scalp and nails.

What Behçet’s disease is

Behçet’s disease is a rare type of inflammatory disease which affects many parts of the body. The most common problem is mouth ulcers.

How Apremilast Anabiosis works

Psoriatic arthritis, psoriasis and Behçet's disease are usually lifelong conditions and there is currently no cure. Apremilast Anabiosis works by reducing the activity of an enzyme in the body called 'phosphodiesterase 4', which is involved in the process of inflammation. By reducing the activity of this enzyme, Apremilast Anabiosis can help to control the inflammation associated with psoriatic arthritis, psoriasis and Behçet's disease, and thereby reduce the signs and symptoms of these conditions.

In psoriatic arthritis, treatment with Apremilast Anabiosis results in an improvement in swollen and painful joints, and can improve your general physical function.

In psoriasis, treatment with Apremilast Anabiosis results in a reduction in psoriatic skin plaque and other signs and symptoms of the disease.

In Behçet's disease, treatment with Apremilast Anabiosis reduces the number of mouth ulcers and can stop them completely. It can also reduce the associated pain.

Apremilast Anabiosis has also been shown to improve the quality of life in patients with psoriasis, psoriatic arthritis or Behçet's disease. This means that the impact of your condition on daily activities, relationships and other factors should be less than it was before.

2. What you need to know before you take Apremilast Anabiosis

Do not take Apremilast Anabiosis:

- if you are allergic to apremilast or any of the other ingredients of this medicine (listed in section 6).
- if you are pregnant or think you may be pregnant

Warnings and precautions

Talk to your doctor or pharmacist before taking Apremilast Anabiosis.

Depression and suicidal thoughts

Tell your doctor before starting Apremilast Anabiosis if you have depression which is getting worse with thoughts of suicide.

You or your caregiver should also tell your doctor straight away of any changes in behaviour or mood, feelings of depression and of any suicidal thoughts you may have after taking Apremilast Anabiosis.

Severe kidney problems

If you have severe kidney problems, your dose will be different – see section 3.

If you are underweight

Talk to your doctor while taking Apremilast Anabiosis if you lose weight without meaning to.

Gut problems

If you experience severe diarrhoea, nausea, or vomiting, you should talk to your doctor.

Children and adolescents

Apremilast Anabiosis has not been studied in children and adolescents, therefore it is not recommended for use in children and adolescents aged 17 years and under.

Other medicines and Apremilast Anabiosis

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines obtained without a prescription and herbal medicines. This is because Apremilast Anabiosis can affect the way some other medicines work. Also some other medicines can affect the way Apremilast Anabiosis works.

In particular, tell your doctor or pharmacist before taking Apremilast Anabiosis if you are taking any of the following medicines:

- rifampicin – an antibiotic for tuberculosis
- phenytoin, phenobarbital and carbamazepine – medicines used in the treatment of seizures of epilepsy
- St. John's Wort – a herbal medicine for mild anxiety and depression.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

There is little information about the effects of Apremilast Anabiosis in pregnancy. You should not become pregnant while taking this medicine and should use an effective method of contraception during treatment with Apremilast Anabiosis.

It is not known if this medicine passes into human milk. You should not use Apremilast Anabiosis while breast-feeding.

Driving and using machines

Apremilast Anabiosis has no effect on the ability to drive and use machines.

Apremilast Anabiosis contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Apremilast Anabiosis contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Apremilast Anabiosis

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

How much to take

- When you first start taking Apremilast Anabiosis, you will receive a 'treatment initiation pack' which contains all the doses as listed in the table below.
- The 'treatment initiation pack' is clearly labelled to make sure you take the correct tablet at the correct time.
- Your treatment will start at a lower dose and will gradually be increased over the first 6 days of treatment.
- The 'treatment initiation pack' will also contain enough tablets for another 8 days at the recommended dose (days 7 to 14).

- The recommended dose of Apremilast Anabiosis is 30 mg twice a day after the titration phase is complete – one 30 mg dose in the morning and one 30 mg dose in the evening, approximately 12 hours apart, with or without food.
- This is a total daily dose of 60 mg. By the end of day 6 you will have reached this recommended dose.
- Once the recommended dose has been reached, you will only get the 30 mg tablet strength in your prescribed packs. You will only ever need to go through this stage of gradually increasing your dose once even if you re-start treatment.

Day	Morning Dose	Evening Dose	Total Daily Dose
Day 1	10 mg (light pink)	Do not take dose	10 mg
Day 2	10 mg (light pink)	10 mg (light pink)	20 mg
Day 3	10 mg (light pink)	20 mg (beige)	30 mg
Day 4	20 mg (beige)	20 mg (beige)	40 mg
Day 5	20 mg (beige)	30 mg (pink)	50 mg
Day 6 onwards	30 mg (pink)	30 mg (pink)	60 mg

People with severe kidney problems

If you have severe kidney problems then the recommended dose of Apremilast Anabiosis is 30 mg **once a day (morning dose)**. Your doctor will talk to you about how to increase your dose when you first start taking Apremilast Anabiosis.

How and when to take Apremilast Anabiosis

- Apremilast Anabiosis is for oral use.
- Swallow the tablets whole, preferably with water.
- You can take the tablets either with or without food.
- Take Apremilast Anabiosis at about the same time each day, one tablet in the morning and one tablet in the evening.
- To remove the tablets from the blister, bend the blister and tear along the perforation to release a single-unit. Then gently peel off the labelled paper foil from the single-unit, beginning at the arrow. Do not push the tablet through the paper foil.

If your condition has not improved after six months of treatment, you should talk to your doctor.

If you take more Apremilast Anabiosis than you should

If you take more Apremilast Anabiosis than you should, talk to a doctor or go to the hospital straight away. Take the medicine pack and this leaflet with you.

If you forget to take Apremilast Anabiosis

- If you miss a dose of Apremilast Anabiosis, take it as soon as you remember. If it is close to the time for your next dose, just skip the missed dose. Take the next dose at your regular time
- Do not take a double dose to make up for a forgotten dose.

If you stop taking Apremilast Anabiosis

- You should continue taking Apremilast Anabiosis until your doctor tells you to stop.
- Do not stop taking Apremilast Anabiosis without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Serious side effects – depression and suicidal thoughts

Tell your doctor straight away about any changes in behaviour or mood, feelings of depression, thoughts of suicide or suicidal behaviour (this is uncommon).

Very common side effects (may affect more than 1 in 10 people)

- Diarrhoea
- Nausea
- Headache
- Upper respiratory tract infections such as cold, runny nose, sinus infection

Common side effects (may affect up to 1 in 10 people)

- Cough
- Back pain
- Vomiting
- Feeling tired
- Stomach pain
- Loss of appetite
- Frequent bowel movements
- Difficulty sleeping (insomnia)
- Indigestion or heartburn
- Inflammation and swelling of the tubes in your lungs (bronchitis)
- Common cold (nasopharyngitis)
- Depression
- Migraine
- Tension headache

Uncommon side effects (may affect up to 1 in 100 people)

- Rash
- Hives (urticaria)
- Weight loss
- Allergic reaction
- Bleeding in the bowel or in the stomach
- Suicidal ideation or behaviour

Not known side effects (frequency cannot be estimated from the available data)

- Severe allergic reaction (may include swelling of the face, lips, mouth, tongue, or throat that may lead to difficulty breathing or swallowing)

If you are 65 years of age or older, you might have a higher risk of complications of severe diarrhoea, nausea and vomiting. If your gut problems become severe, you should talk to your doctor.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system:

Sverige

Läkemedelsverket

Box 26

751 03 Uppsala

Webbplats: www.lakemedelsverket.se

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Apremilast Anabiosis

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and carton after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not use this medicine if you notice any damage or signs of tampering to the medicine packaging.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Apremilast Anabiosis contains

- The active substance is apremilast.
- Apremilast Anabiosis 10 mg film-coated tablets: each film-coated tablet contains 10 mg of apremilast.
- Apremilast Anabiosis 20 mg film-coated tablets: each film-coated tablet contains 20 mg of apremilast.
- Apremilast Anabiosis 30 mg film-coated tablets: each film-coated tablet contains 30 mg of apremilast.

The other ingredients in the tablet core are: lactose monohydrate, microcrystalline cellulose, croscarmellose sodium and magnesium stearate.

The film-coating contains: hypromellose (E464), titanium dioxide (E171), macrogol, iron oxide red (E172).

Apremilast Anabiosis 20 mg film-coated tablets also contain iron oxide yellow (E172).

What Apremilast Anabiosis looks like and contents of the pack

Apremilast Anabiosis 10 mg film-coated tablets are light pink, oval shaped film-coated tablets, debossed “TV” on one side and “Y2” on the other side. The tablet dimensions are approximately 9 mm x 5 mm.

Apremilast Anabiosis 20 mg film-coated tablets are beige, oval shaped, film-coated tablets, debossed “TV” on one side and “Y3” on the other side. The tablet dimensions are approximately 11 mm x 6 mm.

Apremilast Anabiosis 30 mg film-coated tablets are pink, oval shaped, film-coated tablets, debossed “TV” on one side and “Y4” on the other side. The tablet dimensions are approximately 12 mm x 7 mm.

Apremilast Anabiosis 10 mg, 20 mg, 30 mg film-coated tablets (initiation pack)

The treatment initiation pack contains 27 film-coated tablets (4 x 10 mg tablets, 4 x 20 mg tablets, 19 x 30 mg tablets) in PVC/Aluminium foil blisters.

Pack sizes: 27 tablets (in common blister)

27x1 tablets (in single-unit blister)

Apremilast Anabiosis 30 mg film-coated tablets

PVC/Aluminium blister

Pack sizes: 30x1 (in single-unit blister), 56, 56x1 (in single-unit blister), 60, 168x1 (in single-unit blister).

Not all pack sizes may be marketed.

Marketing Authorisation Holder

ANABIOSIS S.A.

Side Road to Kimis Ave. No 3-7

Neo Iraklio, Attica

Greece, 141 22

Tel: +30 210 2711020, +30 210 2711120

Fax: +30 210 2712001

Email: info@anabiosis.gr

Manufacturer

PLIVA Hrvatska d.o.o. (PLIVA Croatia Ltd.),

Prilaz baruna Filipovica 25,

10000 Zagreb,

Croatia

Teva Operations Poland Sp. z o.o.,

ul. Mogilska 80.

31-546, Krakow,

Poland

This medicine is authorised in the Member States of the European Economic Area under the following names:

Sweden: Apremilast Anabiosis

Greece: Apremilast/Anabiosis

This leaflet was last revised in 21 October 2025.

Other sources of information

<Latest approved information on this medicine is available by scanning the QR code included in the <package leaflet><outer carton> with a smartphone/device. The same information is also available on the following URL: [URL to be included] <and the <NCA> website>>