

**PACKAGE LEAFLET**

**Package leaflet: Information for the patient**  
**Asacol 1 g Suppositories**  
mesalazine

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Asacol 1 g Suppositories are and what they are used for
2. What you need to know before you use Asacol 1 g Suppositories
3. How to use Asacol 1 g Suppositories
4. Possible side effects
5. How to store Asacol 1 g Suppositories
6. Contents of the pack and other information

**1. What Asacol 1 g Suppositories are and what they are used for**

Asacol 1 g Suppositories contain the active substance mesalazine, an anti-inflammatory agent used to treat inflammatory bowel disease.

Asacol 1 g Suppositories are used for the treatment of acute mild to moderate ulcerative colitis limited to the rectum (ulcerative proctitis).

**2. What you need to know before you use Asacol 1 g Suppositories**

**Do not use Asacol 1 g Suppositories**

- if you are allergic to salicylic acid, to salicylates such as acetylsalicylic acid (aspirin) or the other ingredients of this medicine (listed in section 6).
- If you have a serious liver or kidney disease.

**Warnings and precautions**

**Talk to your doctor before using Asacol 1 g Suppositories**

- If you have a history of problems with your lungs, particularly if you suffer from **bronchial asthma**.
- If you have a history of allergy to sulphasalazine, a substance related to mesalazine.
- If you suffer with problems of your **liver**.
- If you suffer with problems of your **kidney**.
- If you have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after using mesalazine.

If you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears contact your doctor immediately.

**Further precautions**

During treatment your doctor may want to keep you under close medical supervision, and you may need to have regular blood and urine tests.

Kidney stones may develop with use of mesalazine. Symptoms may include pain in sides of abdomen and blood in urine. Take care to drink sufficient amount of liquid during treatment with mesalazine.

Serious skin reactions including Drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS), and toxic epidermal necrolysis (TEN) have been reported in association with mesalazine treatment. Stop using mesalazine and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

Mesalazine may produce red-brown urine discoloration after contact with sodium hypochlorite bleach in the toilet water. It concerns a chemical reaction between mesalazine and bleach and is harmless.

### **Other medicines and Asacol 1 g Suppositories**

Tell your doctor, if you take or use any of the medicines mentioned below, as the effects of these medicines may change (interactions):

- **Azathioprine, 6-mercaptopurine or thioguanine** (medicines used to treat immune disorders).
- **Certain agents that inhibit blood clotting** (medicines for thrombosis or to thin your blood, e.g., warfarin)

Tell your doctor or pharmacist, if you are taking, have recently taken or might take any other medicines. It may still be all right for you to use Asacol 1 g Suppositories and your doctor will be able to decide, what is suitable for you.

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

You should only use Asacol 1 g Suppositories during pregnancy, if your doctor tells you to.

You should also only use Asacol 1 g Suppositories during breast-feeding, if your doctor tells you to, as this medicine may pass into breast milk.

### **Driving and using machines**

Asacol 1 g Suppositories are not expected to affect your ability to drive or operate machines.

## **3. How to use Asacol 1 g Suppositories**

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

### **Method of administration**

This medicine may only be used rectally, so it has to be inserted through the anus. Do NOT take by mouth.

### **Dosage**

#### Adults and older people

To treat acute phase of ulcerative proctitis the recommended dose is one Asacol 1 g suppository once a day, at bedtime.

The dose should be adjusted by your doctor as per your condition.

#### Use in children

There is little experience and only limited documentation for an effect in children.

### **Duration of treatment**

Your doctor will decide how long you need to continue the treatment with this medicine. This will depend on your condition.

To obtain the maximum benefit from this medicine, you should use Asacol 1 g Suppositories regularly and consistently, as directed.

If you think your Asacol 1 g Suppositories are too strong or too weak, talk to your doctor.

**If you use more Asacol 1 g Suppositories than you should**

Contact a doctor if you are in doubt, so he or she can decide what to do.

If you use too much Asacol 1 g Suppositories on one occasion, just take your next dose as prescribed.

Do not use a smaller amount.

**If you forget to use Asacol 1 g Suppositories**

If you forget to use this medicine, use it as soon as you remember. However, if it is almost time for your next dose, skip the dose you missed. Then use the next dose at its usual time. Do not use a double dose to make up for a forgotten dose.

**If you stop using Asacol 1 g Suppositories**

Do not stop using this product until you have talked to your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

:

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**All medicines can cause allergic reactions although serious allergic reactions are very rare. If you get any of the following symptoms after using this medicine, you should contact your doctor immediately:**

- **Allergic skin rash**
- Reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes, widespread rash, fever and enlarged lymph nodes. These serious skin rashes can be preceded by fever and flu-like symptoms.
- **Fever**
- **Breathing difficulties**

**Tell your doctor immediately** if you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears. These could be symptoms of increased pressure within your skull (idiopathic intracranial hypertension) (frequency not known [cannot be estimated from the available data]).

*If you experience a marked reduction of your general health, especially if accompanied by fever, and/or a sore throat and mouth, stop using these suppositories and contact your doctor immediately.* These symptoms can, very rarely, be due to a fall in the number of white cells in your blood (a condition called agranulocytosis), which may make you more prone to developing a serious infection. A blood test can confirm whether your symptoms are due to an effect of this medicine on your blood.

The following side effects have also been reported by patients using mesalazine:

**Rare side effects** (may affect up to 1 in 1,000 patients):

- Abdominal pain, diarrhoea, wind (flatulence), nausea and vomiting, constipation
- Headache, dizziness
- Chest pain, breathlessness or swollen limbs because of an effect on your heart
- Increased sensitivity of your skin to sun and ultraviolet light (photosensitivity)

**Very rare side effects** (may affect up to 1 in 10,000 patients):

- Changes in kidney function, sometimes with swollen limbs or flank pain
- Severe abdominal pain because of acute inflammation of the pancreas

- Fever, sore throat or malaise, because of blood count changes
- Shortness of breath, cough, wheezing, lung shadow on x-ray due to allergic and/or inflammatory lung conditions
- Severe diarrhoea and abdominal pain because of an allergic reaction to this medicine within the bowel
- Skin rash or inflammation
- Muscle and joint pain
- Jaundice or abdominal pain because of liver or bile flow disorders
- Hair loss and the development of baldness
- Numbness and tingling in the hands and feet (peripheral neuropathy)
- Reversible decrease in semen production

**Not known** (frequency cannot be estimated from the available data)

- Kidney stones and associated kidney pain (see also section 2)

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Asacol 1 g Suppositories**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister strip and on the carton. The expiry date refers to the last day of that month.

Do not store above 30°C.

Store in the original container in order to protect contents from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## **6. Contents of the pack and other information**

### **What Asacol 1 g Suppositories contains**

- The active substance is mesalazine and one suppository of Mesalazine ESPL 1 g Suppositories contains 1 g mesalazine.
- The other ingredient is hard fat.

### **What Asacol 1 g Suppositories look like and contents of the pack**

Asacol 1 g Suppositories are light beige coloured, torpedo-shaped suppositories.

Asacol 1 g Suppositories are available in packs of 10, 20, 30, 60 and 90 suppositories.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

The marketing authorisation holder is

[To be completed nationally]

The manufacturer is

Tillotts Pharma GmbH

Warmbacher Strasse 80

79618 Rheinfelden Germany

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Czech Republic, Denmark, Finland, Germany, Island, Norway, Spain, Sweden: Asacol

Estonia, Latvia, Lithuania: Yaldigo

Ireland: Asacolon

France: Fivasa

Greece, The Netherlands: Kiudro

**This leaflet was last revised in 2025-05-23**

**Other sources of information**

Detailed information on this medicine is available on the website of {name of MS Agency (link)}