




QR code to [www.oeh.fi/xxxx](http://www.oeh.fi/xxxx) [to be completed nationally].

Scan this code or visit [www.oeh.fi/xxxx](http://www.oeh.fi/xxxx) [to be completed nationally] to see instructions on how to use Easyhaler

### Package leaflet: Information for the patient

#### **Bufori Easyhaler, 80 micrograms/4.5 micrograms/inhalation, Inhalation Powder** Budesonide/formoterol fumarate dihydrate

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Bufori Easyhaler 80/4.5 is and what it is used for
2. What you need to know before you use Bufori Easyhaler 80/4.5
3. How to use Bufori Easyhaler 80/4.5
4. Possible side effects
5. How to store Bufori Easyhaler 80/4.5
6. Contents of the pack and other information

#### **1. What Bufori Easyhaler 80/4.5 is and what it is used for**

Bufori Easyhaler is an inhaler that is used to treat asthma in adults, adolescents and children aged 6 years and older. It contains two different medicines: budesonide and formoterol fumarate dihydrate.

- Budesonide belongs to a group of medicines called 'corticosteroids'. It works by reducing and preventing swelling and inflammation in your lungs.
- Formoterol fumarate dihydrate belongs to a group of medicines called 'long-acting beta<sub>2</sub>adrenoceptor agonists' or 'bronchodilators'. It works by relaxing the muscles in your airways. This helps you to breathe more easily.

This medicine is not suitable for people with severe asthma. It can be prescribed for asthma in two different ways.

#### **a) Some people are prescribed two asthma inhalers: Bufori Easyhaler 80/4.5 and a separate 'reliever inhaler'**

- They use Bufori Easyhaler every day. This helps to prevent asthma symptoms from happening.

- They use their ‘reliever inhaler’ when they get asthma symptoms, to make it easier to breathe again.

**b) Some people are prescribed Bufori Easyhaler 80/4.5 as their only asthma inhaler**

- They use Bufori Easyhaler every day. This helps to prevent asthma symptoms from happening.
- They also use Bufori Easyhaler when they need extra doses for relief of asthma symptoms, to make it easier to breathe again. They do not need a separate inhaler for this.

**2. What you need to know before you use Bufori Easyhaler 80/4.5**

**Do not use Bufori Easyhaler 80/4.5:**

- if you are allergic to budesonide, formoterol or to the other ingredient of this medicine (listed in section 6) which is lactose (which contains small amounts of milk protein).

**Warnings and precautions**

Talk to your doctor or pharmacist before using Bufori Easyhaler if you:

- are diabetic
- have a lung infection
- have high blood pressure or you have ever had a heart problem (including an uneven heart beat, a very fast pulse, narrowing of the arteries or heart failure)
- have problems with your thyroid or adrenal glands
- have low levels of potassium in your blood
- have severe liver problems.

Contact your doctor if you experience blurred vision or other visual disturbances.

Rinse your mouth after inhaling your dose in order to avoid fungal infection in the mouth.

**Other medicines and Bufori Easyhaler**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

Some medicines may increase the effects of Bufori Easyhaler and your doctor may wish to monitor you carefully if you are taking these medicines (including some medicines for HIV: ritonavir, cobicistat).

In particular, tell your doctor or pharmacist if you are taking any of the following medicines:

- Beta-blocker medicines (such as atenolol or propranolol for high blood pressure), including eyedrops (such as timolol for glaucoma).
- Medicines for a fast or uneven heart beat (such as quinidine).
- Medicines like digoxin, often used to treat heart failure.
- Diuretics, also known as ‘water tablets’ (such as furosemide). These are used to treat high blood pressure.
- Steroid medicines that you take by mouth (such as prednisolone).
- Xanthine medicines (such as theophylline or aminophylline). These are often used to treat asthma.
- Other bronchodilators (such as salbutamol).
- Tricyclic anti-depressants (such as amitriptyline) and the anti-depressant nefazodone.
- Phenothiazine medicines (such as chlorpromazine and prochlorperazine).
- Medicines to treat infections (such as ketoconazole, itraconazole, voriconazole, posaconazole, clarithromycin and telithromycin).
- Medicines for Parkinson’s disease (such as leva-dopa).
- Medicines for thyroid problems (such as levo-thyroxine).

If any of the above applies to you, or if you are not sure, talk to your doctor or pharmacist before using Bufori Easyhaler.

Also tell your doctor or pharmacist if you are going to have a general anaesthetic for an operation or for dental work.

### **Pregnancy and breast-feeding**

- If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine
- Do not use Bufori Easyhaler unless your doctor tells you to.
- If you get pregnant while using Bufori Easyhaler, do not stop using Bufori Easyhaler but talk to your doctor immediately.

### **Driving and using machines**

Bufori Easyhaler has no or negligible effect on your ability to drive or to use tools or machines.

**Bufori Easyhaler contains lactose** which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, talk to your doctor before using this medicine. The amount of lactose in this medicine does not normally cause problems in people who are lactose intolerant. The excipient lactose contains small amounts of milk proteins, which may cause allergic reaction.

## **3. How to use Bufori Easyhaler 80/4.5**

- Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.
- It is important to use Bufori Easyhaler every day, even if you have no asthma symptoms at the time.
- Your doctor will want to regularly check your asthma symptoms.

If you have been taking steroid tablets for your asthma, your doctor may reduce the number of tablets that you take, once you start to use Bufori Easyhaler. If you have been taking oral steroid tablets for a long time, your doctor may want you to have blood tests from time to time. When reducing oral steroid tablets, you may feel generally unwell even though your chest symptoms may be improving. You might experience symptoms such as a stuffy or runny nose, weakness or joint or muscle pain and rash (eczema). If any of these symptoms bother you, or if symptoms such as headache, tiredness, nausea (feeling sick) or vomiting (being sick) occur, please contact your doctor immediately. You may need to take other medication if you develop allergic or arthritic symptoms. Speak to your doctor if you are concerned as to whether you should continue to use Bufori Easyhaler.

Your doctor may consider adding steroid tablets to your usual treatment during periods of stress (for example, when you have a chest infection or before an operation).

### **Important information about your asthma symptoms**

If you feel you are getting breathless or wheezy while using Bufori Easyhaler, you should continue to use Bufori Easyhaler but go to see your doctor as soon as possible, as you may need additional treatment.

Contact your doctor immediately if:

- your breathing is getting worse or you often wake up at night with asthma.
  - your chest starts to feel tight in the morning or your chest tightness lasts longer than usual.
- These signs could mean that your asthma is not being properly controlled and you may need different or additional treatment immediately.

## **ASTHMA**

Bufori Easyhaler 80/4.5 can be prescribed for asthma in two different ways. The amount of Bufori Easyhaler to use and when to use it depends on how it has been prescribed for you.

- a) If you have been prescribed Bufori Easyhaler and a separate reliever inhaler, read the section called 'A) Using Bufori Easyhaler and a separate reliever inhaler'.
- b) If you have been prescribed Bufori Easyhaler as your only inhaler, read the section called 'B) Using Bufori Easyhaler as your only asthma inhaler'.

### **A) Using Bufori Easyhaler and a separate reliever inhaler**

**Use your Bufori Easyhaler every day.** This helps to prevent asthma symptoms from happening.

#### **Adults (18 years and above)**

- The usual dose is 1 or 2 inhalations, twice a day.
- Your doctor may increase this to 4 inhalations, twice a day.

If your symptoms are well controlled, your doctor may ask you to take your medicine once a day.

#### **Adolescents (12 to 17 years)**

- The usual dose is 1 or 2 inhalations, twice a day.
- If your symptoms are well controlled, your doctor may ask you to take your medicine once a day.

#### **Children (6 to 11 years)**

- The usual dose is 2 inhalations, twice a day.

Bufori Easyhaler 80/4.5 is not recommended to be used in children who are younger than 6 years.

Your doctor (or asthma nurse) will help you to manage your asthma. They will adjust the dose of this medicine to the lowest dose that controls your asthma. However, do not adjust the dose without talking to your doctor (or asthma nurse) first.

**Use your separate 'reliever inhaler' to treat asthma symptoms when they happen.** Always keep your 'reliever inhaler' with you to use when you need it. Do not use Bufori Easyhaler to treat asthma symptoms - use your reliever inhaler.

### **B) Using Bufori Easyhaler as your only asthma inhaler**

Only use Bufori Easyhaler in this way if your doctor has told you to and if you are aged 12 or above.

**Use your Bufori Easyhaler 80/4.5 every day.** This helps to prevent asthma symptoms from happening. You can take:

- 1 inhalation in the morning **and** 1 inhalation in the evening
- or**
- 2 inhalations in the morning
- or**
- 2 inhalations in the evening.

**Also use Bufori Easyhaler 80/4.5 as a 'reliever inhaler' to treat asthma symptoms when they happen.**

- If you get asthma symptoms, take 1 inhalation and wait a few minutes.
- If you do not feel better, take another inhalation.
- Do not take more than 6 inhalations at a single time.

Always keep your Bufori Easyhaler with you, so you can use it when you need it.

A total daily dose of more than 8 inhalations is not normally needed. However, your doctor may allow you to take up to 12 inhalations a day for a limited period.

If you regularly need to use 8 or more inhalations a day, make an appointment to see your doctor or nurse. They may need to change your treatment.

Do not use more than 12 inhalations in total in 24 hours.

If you are doing exercise and you get asthma symptoms, use Bufen Easyhaler as described here. However, do not use Bufen Easyhaler just before exercise to stop asthma symptoms from happening.

### **If you use more Bufen Easyhaler than you should**

It is important that you take your dose as stated on the pharmacist's label or as advised by your doctor. You should not exceed your prescribed dose without seeking medical advice.

The most common symptoms that may occur after if you use more Bufen Easyhaler 80/4.5 than you should are trembling, headache or a rapid heart beat.

### **If you forget to use Bufen Easyhaler**

- If you forget to take a dose, take it as soon as you remember. However, if it is nearly time for your next dose, skip the missed dose.
- Do **not** take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

**The instructions how to use the inhaler are at the end of the leaflet.**

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**If either of the following happen to you, stop using Bufen Easyhaler and talk to your doctor immediately:**

- Swelling of your face, particularly around your mouth (tongue and/or throat and/or difficulty to swallow) or hives together with difficulties to breath (angioedema) and/or sudden feeling of faintness. This may mean that you are having an allergic reaction. This happens rarely, affecting less than 1 in 1 000 people.
- Sudden acute wheezing or shortness of breath immediately after using your inhaler. **If either of these symptoms occur, stop using your Bufen Easyhaler inhaler straightaway and use your 'reliever' inhaler. Contact your doctor immediately as you may need to have your treatment changed.** This happens very rarely, affecting less than 1 in 10 000 people.

**Other possible side effects:**

**Common (may affect up to 1 in 10 people)**

- Palpitations (awareness of your heart beating), trembling or shaking. If these effects occur, they are usually mild and usually disappear as you continue to use Bufen Easyhaler.
- Thrush (a fungal infection) in the mouth. This is less likely if you rinse your mouth out with water after using your Bufen Easyhaler.
- Mild sore throat, coughing and a hoarse voice.
- Headache.

**Uncommon (may affect up to 1 in 100 people)**

- Feeling restless, nervous or agitated.
- Disturbed sleep.
- Feeling dizzy.
- Nausea (feeling sick).
- Fast heart beat.
- Bruising of the skin.
- Muscle cramps.

- Blurred vision.

#### **Rare (may affect up to 1 in 1 000 people)**

- Rash, itching.
- Bronchospasm (tightening of the muscles in the airways which causes wheezing). If the wheezing comes on suddenly after using Bufori Easyhaler stop using Bufori Easyhaler and talk to your doctor immediately.
- Low levels of potassium in your blood.
- Uneven heart beat.

#### **Very rare (may affect up to 1 in 10 000 people)**

- Depression.
- Changes in behaviour, especially in children.
- Chest pain or tightness in the chest (angina pectoris).
- An increase in the amount of sugar (glucose) in your blood.
- Taste changes, such as an unpleasant taste in the mouth.
- Changes in your blood pressure.

Inhaled corticosteroids can affect the normal production of steroid hormones in your body, particularly if you use high doses for a long time. The effects include:

- changes in bone mineral density (thinning of the bones)
- cataract (clouding of the lens in the eye)
- glaucoma (increased pressure in the eye)
- a slowing of the rate of growth of children and adolescents
- an effect on the adrenal gland (a small gland next to the kidney).

These effects are much less likely to happen with inhaled corticosteroids than with corticosteroid tablets.

#### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [\[To be completed nationally\]](#). By reporting side effects, you can help provide more information on the safety of this medicine.

### **5. How to store Bufori Easyhaler 80/4.5**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date that is stated on the carton, foil bag and label of your inhaler after EXP. The expiry date refers to the last day of that month.
- After opening the foil bag do not store above 25°C and store protected from moisture. You may use a protective cover with your Easyhaler.
- If your Bufori Easyhaler gets damp you need to replace it with a new one.
- Replace Bufori Easyhaler 4 months after you opened the foil bag. Write down the date you opened the bag to help you remember.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

### **6. Contents of the pack and other information**

#### **What Bufori Easyhaler 80/4.5 contains**

- The active substances are budesonide and formoterol fumarate dihydrate. Each inhaled dose contains 80 micrograms of budesonide and 4.5 micrograms of formoterol fumarate dihydrate.
- The other ingredient is lactose monohydrate (which contains milk proteins).

**What Bufori Easyhaler 80/4.5 looks like and contents of the pack**

Bufori Easyhaler 80/4.5 is an inhaler containing your medicine. The inhalation powder is white to yellowish in colour. Each inhaler contains 60 or 120 doses and has a white body with red upper part. Bufori Easyhaler 80/4.5 is available in packs of 1 or 3 inhaler(s).

Not all pack sizes may be marketed.

Your pack may contain a protective cover that is available depending on the country. If you need a protective cover, please contact the Marketing Authorisation Holder (details given below).

**Marketing Authorisation Holder**

Orion Corporation

Orionintie 1

FI-02200 Espoo

Finland

**Manufacturer**

Orion Corporation Orion Pharma

Orionintie 1

FI-02200 Espoo

Finland

**This leaflet was last revised on 14/06/2025**

Detailed and updated information on how to use this product is available by scanning this QR code (included also on outer carton and inhaler label) with a smartphone. The same information is also available on the following URL: [www.oeh.fi/xxxx](http://www.oeh.fi/xxxx) [to be completed nationally].

QR code to [www.oeh.fi/xxxx](http://www.oeh.fi/xxxx) [to be completed nationally].

## How to use the Easyhaler inhaler

### About your Easyhaler

Bufori Easyhaler may be different to inhalers you have used in the past. Therefore it is very important that you use it properly as using incorrectly can lead to you not receiving the right amount of medicine. This could make you very unwell or could lead to your asthma not being treated as it should.


Your doctor, nurse or pharmacist will show you how to use your inhaler properly. Make sure you understand the correct way to use the inhaler. If you are unsure contact your doctor, nurse or pharmacist. As with all inhalers, caregivers should ensure that children prescribed Bufori Easyhaler use correct inhalation technique, as described below. You can also use the instruction video at [www.oeh.fi/xxxx](http://www.oeh.fi/xxxx)

### When you first get your Easyhaler

The Easyhaler comes in a foil bag. Do not open the foil bag until you are ready to start using the medicine as it helps to keep the powder dry in the inhaler.

When you are ready to start treatment open the bag and record the date e.g., in your calendar.

Use the inhaler within 4 months of removing from the foil bag.



## HOW TO USE CORRECTLY

### Step 1: SHAKE


- Remove the dustcap
- Shake the inhaler 3 to 5 times holding it in the **upright position**

### SHAKE x 3-5



### Important points to remember

- It is important to keep the inhaler in the upright position
- If you accidentally click while you shake the inhaler, empty the powder from the mouthpiece as shown below

|                                                                                                                                                                                                                                                                                                    |                                                                                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Step 2: CLICK</b></p> <ul style="list-style-type: none"> <li>• Keep holding the inhaler upright between your forefinger and thumb</li> <li>• Press down until you hear a click, and let the inhaler click back again. This releases a dose</li> <li>• <b>Only click down once</b></li> </ul> | <p><b>CLICK x 1</b></p>  | <p><b>Important points to remember</b></p> <ul style="list-style-type: none"> <li>• The inhaler will not click if the dustcap is still on</li> <li>• Only click down once</li> <li>• If you accidentally click more than once, empty the powder from the mouthpiece, see below</li> <li>• Click to release dose before you inhale, not at the same time</li> <li>• Keep the inhaler <b>upright</b> when you click it and when you inhale the dose. If you tip it, the powder could fall out before you are able to inhale it</li> </ul> |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

|                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Step 3: INHALE</b></p> <ul style="list-style-type: none"> <li>• Keep holding the inhaler upright</li> <li>• Breathe out normally</li> <li>• Place the mouthpiece in your mouth between your teeth and close your lips tightly around the mouthpiece</li> <li>• Take a strong and deep breath in</li> <li>• Take the inhaler out of your mouth, then breathe out normally.</li> </ul> | <p><b>INHALE</b></p>  | <p><b>Important points to remember</b></p> <ul style="list-style-type: none"> <li>• Make sure the whole mouthpiece is well inside your mouth, so that the medication gets into your lungs</li> <li>• Make sure your lips make a good seal around the mouthpiece</li> <li>• Do not breathe out into the inhaler. This is important: it could clog up the inhaler. If you breathed out into the inhaler, empty the powder from mouthpiece, see below</li> </ul> |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

**If you need to take another inhalation, please repeat the steps 1–3 Shake-Click-Inhale.**


**After you have used the inhaler:**

- Put the dust cap back on the mouthpiece. It stops the inhaler going off by accident.
- After you have taken the dose, rinse your mouth with water, and spit it out.

### How to empty the powder from the mouthpiece

If you click the inhaler by accident, or if you might have clicked it more than once, or if you breathe out into it, empty the mouthpiece.

- Tap the mouthpiece to empty the powder onto a table top, or the palm of your hand.
- Then start again with steps Shake-Click-Inhale.



### Cleaning the Easyhaler

Keep your inhaler dry and clean. If necessary you may wipe the mouthpiece of your inhaler with a dry cloth or tissue. Do not use water: the powder in the Easyhaler is sensitive to moisture.

### Using the Easyhaler with a protective cover

You may use a protective cover with your Easyhaler. This may help to improve durability of the product. When you first insert your inhaler in the protective cover make sure the dustcap is on the inhaler as this stops it going off by accident. You can use the inhaler without removing it from the protective cover.

Follow the same instructions as above, **1. Shake – 2. Click – 3. Inhale.**

Remember to:

- Keep the inhaler in the upright position when clicking it
- Replace the dustcap after taking the dose as this stops the inhaler going off by accident.



### When to switch to a new Easyhaler

The dose counter shows the number of remaining doses. The counter turns after every 5<sup>th</sup> click. When the dose counter starts turning red, there are 20 doses left.

If you do not already have a new Easyhaler, contact your doctor for a new prescription. When the counter reaches 0 (zero), you need to replace the Easyhaler.

If you use the protective cover, you can keep it and insert your new inhaler into it.



**Remember**

- **1. Shake – 2. Click – 3. Inhale.**
- After you have taken the dose, rinse your mouth with water and spit it out.
- Do not get your inhaler wet, protect it from moisture.

**If you have any further questions on the use of this product, ask your doctor or pharmacist.**