

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

Desloratadin Orion 5 mg orodispersible tablets

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each orodispersible tablet contains 5 mg desloratadine.

#### Excipient with known effect

Each orodispersible tablet contains 3 mg aspartame (E951).

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Orodispersible tablet.

Light red, round flat faced, speckled tablets debossed with “D2” on one side and plain on the other.  
Diameter of approx. 8 mm.

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Desloratadin Orion is indicated in adults and adolescents aged 12 years and older for the relief of symptoms associated with:

- allergic rhinitis (see section 5.1)
- urticaria (see section 5.1).

#### 4.2 Posology and method of administration

##### Posology

*Adults and adolescents (12 years of age and over)*

The recommended dose is one 5 mg orodispersible tablet placed in the mouth once a day.

Intermittent allergic rhinitis (presence of symptoms for less than 4 days per week or for less than 4 weeks) should be managed in accordance with the evaluation of patient's disease history and the treatment could be discontinued after symptoms are resolved and reinitiated upon their reappearance. In persistent allergic rhinitis (presence of symptoms for 4 days or more per week and for more than 4 weeks), continued treatment may be proposed to the patients during the allergen exposure periods.

##### *Paediatric population*

The safety and efficacy of Desloratadin Orion 5 mg orodispersible tablets in children below the age of 12 years have not been established.

There is limited clinical trial efficacy experience with the use of desloratadine in adolescents 12 through 17 years of age (see sections 4.8 and 5.1).

##### Method of administration

Oral use.

The dose can be taken with or without food.

The orodispersible tablet is placed in the mouth where it will disperse rapidly. Water or other liquid is not needed to swallow the tablet.

### **4.3 Contraindications**

Hypersensitivity to the active substance, to any of the excipients listed in section 6.1, or to loratadine.

### **4.4 Special warnings and precautions for use**

#### Renal function impairment

In the case of severe renal insufficiency, desloratadine should be used with caution (see section 5.2).

#### Seizures

Desloratadine should be administered with caution in patients with medical or familial history of seizures, and mainly young children (see section 4.8), being more susceptible to develop new seizures under desloratadine treatment. Healthcare providers may consider discontinuing desloratadine in patients who experience a seizure while on treatment.

#### Excipients

This medicinal product contains 3 mg aspartame per tablet. Aspartame is hydrolysed in the gastrointestinal tract when orally ingested. One of the major hydrolysis products is phenylalanine. It may be harmful if the patient has phenylketonuria (PKU).

This medicinal product contains less than 1 mmol (23 mg) sodium per tablet, that is to say essentially “sodium-free”.

### **4.5 Interaction with other medicinal products and other forms of interaction**

No clinically relevant interactions were observed in clinical trials with desloratadine tablets in which erythromycin or ketoconazole were co-administered (see section 5.1).

#### Paediatric population

Interaction studies have only been performed in adults.

In a clinical pharmacology trial, desloratadine tablets taken concomitantly with alcohol did not potentiate the performance impairing effects of alcohol (see section 5.1). However, cases of alcohol intolerance and intoxication have been reported during post-marketing use. Therefore, caution is recommended if alcohol is taken concomitantly.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

A large amount of data on pregnant women (more than 1 000 pregnancy outcomes) indicates no malformative nor foetal/neonatal toxicity of desloratadine. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). As a precautionary measure, it is preferable to avoid the use of desloratadine during pregnancy.

#### Breast-feeding

Desloratadine has been identified in breastfed newborns/infants of treated women. The effect of desloratadine on newborns/infants is unknown. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from desloratadine therapy taking into account the benefit of breast feeding for the child and the benefit of therapy for the woman.

#### Fertility

There are no data available on male and female fertility.

#### 4.7 Effects on ability to drive and use machines

Desloratadin Orion has no or negligible influence on the ability to drive and use machines based on clinical trials. Patients should be informed that most people do not experience drowsiness. Nevertheless, as there is individual variation in response to all medicinal products, it is recommended that patients are advised not to engage in activities requiring mental alertness, such as driving a car or using machines, until they have established their own response to the medicinal product.

#### 4.8 Undesirable effects

##### Summary of the safety profile

In clinical trials in a range of indications including allergic rhinitis and chronic idiopathic urticaria, at the recommended dose of 5 mg daily, undesirable effects with desloratadine tablets were reported in 3 % of patients in excess of those treated with placebo. The most frequent of adverse reactions reported in excess of placebo were fatigue (1.2 %), dry mouth (0.8 %) and headache (0.6 %).

##### Tabulated list of adverse reactions

The frequency of the clinical trial adverse reactions reported in excess of placebo and other undesirable effects reported during the post-marketing period are listed in the following table.

Frequencies are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ), rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ), very rare ( $< 1/10\ 000$ ) and not known (cannot be estimated from the available data).

| System Organ Class                                          | Frequency | Adverse reactions                                                                                     |
|-------------------------------------------------------------|-----------|-------------------------------------------------------------------------------------------------------|
| <b>Metabolism and nutrition disorders</b>                   | Not known | Increased appetite                                                                                    |
| <b>Psychiatric disorders</b>                                | Very rare | Hallucinations                                                                                        |
|                                                             | Not known | Abnormal behaviour*, aggression*, depressed mood                                                      |
| <b>Nervous system disorders</b>                             | Common    | Headache                                                                                              |
|                                                             | Very rare | Dizziness, somnolence, insomnia, psychomotor hyperactivity, seizures                                  |
| <b>Eye disorders</b>                                        | Not known | Eye dryness                                                                                           |
| <b>Cardiac disorders</b>                                    | Very rare | Tachycardia, palpitations                                                                             |
|                                                             | Not known | QT prolongation*                                                                                      |
| <b>Gastrointestinal disorders</b>                           | Common    | Dry mouth                                                                                             |
|                                                             | Very rare | Abdominal pain, nausea, vomiting, dyspepsia, diarrhoea                                                |
| <b>Hepatobiliary disorders</b>                              | Very rare | Elevations of liver enzymes, increased bilirubin, hepatitis                                           |
|                                                             | Not known | Jaundice                                                                                              |
| <b>Skin and subcutaneous tissue disorders</b>               | Not known | Photosensitivity                                                                                      |
| <b>Musculoskeletal and connective tissue disorders</b>      | Very rare | Myalgia                                                                                               |
| <b>General disorders and administration site conditions</b> | Common    | Fatigue                                                                                               |
|                                                             | Very rare | Hypersensitivity reactions (such as anaphylaxis, angioedema, dyspnoea, pruritus, rash, and urticaria) |
|                                                             | Not known | Asthenia                                                                                              |
| <b>Investigations</b>                                       | Not known | Weight increased                                                                                      |

\* Undesirable effects reported during the post-marketing period also in paediatric patients.

#### Paediatric population

Other undesirable effects reported during the post-marketing period in paediatric patients with an unknown frequency included arrhythmia and bradycardia.

In a clinical trial with 578 adolescent patients, 12 through 17 years of age, the most common adverse event was headache; this occurred in 5.9% of patients treated with desloratadine and 6.9% of patients receiving placebo.

A retrospective observational safety study indicated an increased incidence of new-onset seizure in patients 0 to 19 years of age when receiving desloratadine compared with periods not receiving desloratadine. Among children 0–4 years old, the adjusted absolute increase was 37.5 (95% Confidence Interval (CI) 10.5–64.5) per 100 000 person years (PY) with a background rate of new onset seizure of 80.3 per 100 000 PY. Among patients 5–19 years of age, the adjusted absolute increase was 11.3 (95% CI 2.3–20.2) per 100 000 PY with a background rate of 36.4 per 100 000 PY (see section 4.4).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [the national reporting system listed in Appendix V](#).

### **4.9 Overdose**

The adverse event profile associated with overdosage, as seen during post-marketing use, is similar to that seen with therapeutic doses, but the magnitude of the effects can be higher.

#### Treatment

In the event of overdose, consider standard measures to remove unabsorbed active substance. Symptomatic and supportive treatment is recommended.

Desloratadine is not eliminated by haemodialysis; it is not known if it is eliminated by peritoneal dialysis.

#### Symptoms

Based on a multiple dose clinical trial, in which up to 45 mg of desloratadine was administered (nine times the clinical dose), no clinically relevant effects were observed.

#### Paediatric population

The adverse event profile associated with overdosage, as seen during post-marketing use, is similar to that seen with therapeutic doses, but the magnitude of the effects can be higher.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antihistamines for systemic use, other antihistamines for systemic use; ATC code: R06AX27

#### Mechanism of action

Desloratadine is a non-sedating, long-acting histamine antagonist with selective peripheral H<sub>1</sub>-receptor antagonist activity. After oral administration, desloratadine selectively blocks peripheral histamine H<sub>1</sub>-receptors because the substance is excluded from entry to the central nervous system.

Desloratadine has demonstrated antiallergic properties from *in vitro* studies. These include inhibiting the release of proinflammatory cytokines such as IL-4, IL-6, IL-8, and IL-13 from human mast

cells/basophils, as well as inhibition of the expression of the adhesion molecule P-selectin on endothelial cells. The clinical relevance of these observations remains to be confirmed.

#### Clinical efficacy and safety

In a multiple dose trial, desloratadine orodispersible tablets were well tolerated.

At the recommended dose, desloratadine 5 mg orodispersible tablet was found to be bioequivalent to the desloratadine 5 mg conventional tablet formulation of desloratadine. Therefore, the efficacy of desloratadine orodispersible tablet is expected to be the same as with the desloratadine tablet formulation.

In a multiple dose clinical trial, in which up to 20 mg of desloratadine was administered daily for 14 days, no statistically or clinically relevant cardiovascular effect was observed. In a clinical pharmacology trial, in which desloratadine was administered at a dose of 45 mg daily (nine times the clinical dose) for ten days, no prolongation of QTc interval was seen.

No clinically relevant changes in desloratadine plasma concentrations were observed in multiple-dose, ketoconazole and erythromycin interaction trials.

#### Pharmacodynamic effects

Desloratadine does not readily penetrate the central nervous system. In clinical trials, at the recommended dose of 5 mg daily, there was no excess incidence of somnolence as compared to placebo. Desloratadine tablets given at a single daily dose of 7.5 mg did not affect psychomotor performance in clinical trials. In a single dose study performed in adults, desloratadine 5 mg did not affect standard measures of flight performance including exacerbation of subjective sleepiness or tasks related to flying.

In clinical pharmacology trials, co-administration with alcohol did not increase the alcohol-induced impairment in performance or increase in sleepiness. No significant differences were found in the psychomotor test results between desloratadine and placebo groups, whether administered alone or with alcohol.

In patients with allergic rhinitis, desloratadine tablets were effective in relieving symptoms such as sneezing, nasal discharge and itching, as well as ocular itching, tearing and redness, and itching of palate. Desloratadine tablets effectively controlled symptoms for 24 hours.

#### Paediatric population

The efficacy of desloratadine tablets has not been clearly demonstrated in trials with adolescent patients 12 through 17 years of age.

In addition to the established classifications of seasonal and perennial, allergic rhinitis can alternatively be classified as intermittent allergic rhinitis and persistent allergic rhinitis according to the duration of symptoms. Intermittent allergic rhinitis is defined as the presence of symptoms for less than 4 days per week or for less than 4 weeks. Persistent allergic rhinitis is defined as the presence of symptoms for 4 days or more per week and for more than 4 weeks.

Desloratadine was effective in alleviating the burden of seasonal allergic rhinitis as shown by the total score of the rhino-conjunctivitis quality of life questionnaire. The greatest amelioration was seen in the domains of practical problems and daily activities limited by symptoms.

Chronic idiopathic urticaria was studied as a clinical model for urticarial conditions, since the underlying pathophysiology is similar, regardless of aetiology, and because chronic patients can be more easily recruited prospectively. Since histamine release is a causal factor in all urticarial diseases, desloratadine is expected to be effective in providing symptomatic relief for other urticarial conditions, in addition to chronic idiopathic urticaria, as advised in clinical guidelines.

In two placebo-controlled six-week trials in patients with chronic idiopathic urticaria, desloratadine was effective in relieving pruritus and decreasing the size and number of hives by the end of the first dosing interval. In each trial, the effects were sustained over the 24-hour dosing interval. As with other antihistamine trials in chronic idiopathic urticaria, the minority of patients who were identified as non-responsive to antihistamines was excluded. An improvement in pruritus of more than 50% was observed in 55% of patients treated with desloratadine compared with 19% of patients treated with placebo. Treatment with desloratadine also significantly reduced interference with sleep and daytime function, as measured by a four-point scale used to assess these variables.

## 5.2 Pharmacokinetic properties

### Absorption

Desloratadine plasma concentrations can be detected within 30 minutes of administration. Desloratadine is well absorbed with maximum concentration achieved after approximately 3 hours; the terminal phase half-life is approximately 27 hours. The degree of accumulation of desloratadine was consistent with its half-life (approximately 27 hours) and a once daily dosing frequency. The bioavailability of desloratadine was dose proportional over the range of 5 mg to 20 mg.

In a series of pharmacokinetic and clinical trials, 6% of the subjects reached a higher concentration of desloratadine. The prevalence of this poor metaboliser phenotype was greater among Black adults than Caucasian adults (18% vs. 2%) however the safety profile of these subjects was not different from that of the general population.

In a multiple-dose pharmacokinetic study conducted with the tablet formulation in healthy adult subjects, four subjects were found to be poor metabolisers of desloratadine. These subjects had a  $C_{max}$  concentration about 3-fold higher at approximately 7 hours with a terminal phase half-life of approximately 89 hours.

### Distribution

Desloratadine is moderately bound (83% – 87%) to plasma proteins. There is no evidence of clinically relevant medicine accumulation following once daily dosing of desloratadine (5 mg to 20 mg) for 14 days.

### Biotransformation

The enzyme responsible for the metabolism of desloratadine has not been identified yet, and therefore, some interactions with other medicinal products cannot be fully excluded. Desloratadine does not inhibit CYP3A4 *in vivo*, and *in vitro* studies have shown that the medicinal product does not inhibit CYP2D6 and is neither a substrate nor an inhibitor of P-glycoprotein.

In single-dose crossover studies of desloratadine 5 mg orodispersible tablets with desloratadine 5 mg conventional tablets, the formulations were bioequivalent.

### Elimination

The presence of food prolongs  $T_{max}$  for desloratadine from 2.5 to 4 hours and  $T_{max}$  for 3-OH-desloratadine from 4 to 6 hours. In a separate study, grapefruit juice had no effect on the disposition of desloratadine. Water had no effect on the bioavailability of desloratadine orodispersible tablets.

### Renally impaired patients

The pharmacokinetics of desloratadine in patients with chronic renal insufficiency (CRI) was compared with that of healthy subjects in one single-dose study and one multiple dose study. In the single-dose study, the exposure to desloratadine was approximately 2 and 2.5-fold greater in subjects with mild to moderate and severe CRI, respectively, than in healthy subjects. In the multiple-dose study, steady state was reached after Day 11, and compared to healthy subjects the exposure to desloratadine was ~1.5-fold greater in subjects with mild to moderate CRI and ~2.5-fold greater in subjects with severe CRI. In both studies, changes in exposure (AUC and  $C_{max}$ ) of desloratadine and 3-hydroxydesloratadine were not clinically relevant.

### **5.3 Preclinical safety data**

Desloratadine is the primary active metabolite of loratadine. Non-clinical studies conducted with desloratadine and loratadine demonstrated that there are no qualitative or quantitative differences in the toxicity profile of desloratadine and loratadine at comparable levels of exposure to desloratadine.

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction and development. The collective analysis of preclinical and clinical irritation studies for the orodispersible tablet indicate that this formulation is unlikely to pose risk for local irritation with clinical use. The lack of carcinogenic potential was demonstrated in studies conducted with desloratadine and loratadine.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Mannitol (E421)  
Cellulose, microcrystalline  
Croscarmellose sodium  
Maize starch  
Red iron oxide  
Polacrillin potassium  
Citric acid  
Peppermint flavour (maltodextrin, acacia, flavouring components)  
Aspartame (E951)  
Magnesium stearate

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Aluminium/Aluminium blister.  
Pack sizes: 30, 90 and 100 orodispersible tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Orion Corporation

Orionintie 1  
FI-02200 Espoo  
Finland

**8. MARKETING AUTHORISATION NUMBER(S)**

[To be completed nationally]

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

[To be completed nationally]

**10. DATE OF REVISION OF THE TEXT**

11/06/2026