

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Finpros 5 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 5 mg finasteride.

Excipient with known effect:

Lactose monohydrate (90.95 mg/tablet).

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet.

Blue, round biconvex tablet marked "F5". The diameter is 7 mm.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Finpros is indicated for the treatment and control of benign prostatic hyperplasia (BPH) in patients with an enlarged prostate to:

- cause regression of the enlarged prostate, improve urinary flow and improve the symptoms associated with BPH
- reduce the incidence of acute urinary retention and the need for surgery including transurethral resection of the prostate (TURP) and prostatectomy.

Finpros 5 mg tablets should only be administered in patients with an enlarged prostate (prostate volume above ca. 40 ml).

4.2 Posology and method of administration

Posology

The recommended dosage is one 5 mg tablet daily with or without food.

Finasteride can be administered alone or in combination with the alpha-blocker doxazosin (see section 5.1 'Pharmacodynamic properties').

Even if improvement can be seen within a short time, treatment for at least 6 months may be necessary in order to determine objectively whether a satisfactory response to treatment has been achieved.

Elderly patients

Dosage adjustments are not necessary although pharmacokinetic studies have shown that the elimination rate of finasteride is slightly decreased in patients above 70 years of age.

Patients with renal impairment

Dosage adjustments are not necessary in patients with varying degrees of renal insufficiency (with creatinine clearance down to as low as 9 ml/min) as in pharmacokinetic studies renal insufficiency was

not found to affect the elimination of finasteride. Finasteride has not been studied in patients on haemodialysis.

Patients with hepatic impairment

There are no data available in patients with hepatic insufficiency (see section 4.4).

Paediatric population

Finpros is not indicated for use in children (see section 4.3).

Method of administration

For oral use only.

The tablet should be swallowed whole and must not be divided or crushed (see section 6.6).

4.3 Contraindications

Finasteride is not indicated for use in women or children.

Finasteride is contraindicated in the following:

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Pregnancy - Use in women when they are or may potentially be pregnant (see 4.6 Pregnancy and lactation, Exposure to finasteride - risk to male fetus).

4.4 Special warnings and precautions for use

General:

- To avoid obstructive complications it is important that patients with large residual urine and/or heavily decreased urinary flow are carefully controlled. The possibility of surgery should be an option.

Mood alterations and depression

Mood alterations including depressed mood, depression and, less frequently, suicidal ideation have been reported in patients treated with finasteride 5 mg. Patients should be monitored for psychiatric symptoms and if these occur, the patient should be advised to seek medical advice.

Effects on PSA and prostate cancer detection

No clinical benefit has yet been demonstrated in patients with prostate cancer treated with finasteride 5 mg. Patients with BPH and elevated serum prostate specific antigen (PSA) were monitored in controlled clinical studies with serial PSAs and prostate biopsies. In these BPH studies, finasteride 5 mg did not appear to alter the rate of prostate cancer detection, and the overall incidence of prostate cancer was not significantly different in patients treated with finasteride 5 mg or placebo.

Digital rectal examinations as well as other evaluations for prostate cancer are recommended prior to initiating therapy with finasteride 5 mg and periodically thereafter. Serum PSA is also used for prostate cancer detection. Generally a baseline PSA >10 ng/mL (Hybritech) prompts further evaluation and consideration of biopsy; for PSA levels between 4 and 10 ng/mL, further evaluation is advisable. There is considerable overlap in PSA levels among men with and without prostate cancer. Therefore, in men with BPH, PSA values within the normal reference range do not rule out prostate cancer, regardless of treatment with finasteride 5 mg. A baseline PSA <4 ng/mL does not exclude prostate cancer.

Finasteride 5 mg causes a decrease in serum PSA concentrations by approximately 50% in patients with BPH, even in the presence of prostate cancer. This decrease in serum PSA levels in patients with BPH treated with finasteride 5 mg should be considered when evaluating PSA data and does not rule out concomitant prostate cancer. This decrease is predictable over the entire range of PSA values,

although it may vary in individual patients. Analysis of PSA data from over 3000 patients in the 4-year, double-blind, placebo-controlled finasteride Long-Term Efficacy and Safety Study (PLESS) confirmed that in typical patients treated with finasteride 5 mg for six months or more, PSA values should be doubled for comparison with normal ranges in untreated men. This adjustment preserves the sensitivity and specificity of the PSA assay and maintains its ability to detect prostate cancer.

Any sustained increase in PSA levels of patients treated with finasteride 5 mg should be carefully evaluated, including consideration of non-compliance to therapy with finasteride 5 mg.

Percent free PSA (free to total PSA ratio) is not significantly decreased by finasteride 5 mg. The ratio of free to total PSA remains constant even under the influence of finasteride 5 mg. When percent free PSA is used as an aid in the detection of prostate cancer, no adjustment to its value is necessary.

Drug/laboratory test interactions

Effect on levels of PSA

Serum PSA concentration is correlated with patient age and prostatic volume, and prostatic volume is correlated with patient age. When PSA laboratory determinations are evaluated, consideration should be given to the fact that PSA levels decrease in patients treated with finasteride 5 mg. In most patients, a rapid decrease in PSA is seen within the first months of therapy, after which time PSA levels stabilize to a new baseline. The post-treatment baseline approximates half of the pre-treatment value. Therefore, in typical patients treated with finasteride 5 mg for six months or more, PSA values should be doubled for comparison to normal ranges in untreated men. For clinical interpretation, see 4.4 Special warnings and precautions for use, Effects on PSA and prostate cancer detection.

Breast cancer in men

Breast cancer has been reported in men taking finasteride 5 mg during clinical trials and the post-marketing period. Physicians should instruct their patients to promptly report any changes in their breast tissue such as lumps, pain, gynaecomastia or nipple discharge.

Paediatric use

Finasteride is not indicated for use in children.

Safety and effectiveness in children have not been established.

Hepatic insufficiency

The effect of hepatic insufficiency on the pharmacokinetics of finasteride has not been studied.

Finpros contains lactose and sodium

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

No drug interactions of clinical importance have been identified. Finasteride is metabolized primarily via, but does not appear to affect significantly, the cytochrome P450 3A4 system. Although the risk for finasteride to affect the pharmacokinetics of other drugs is estimated to be small, it is probable that inhibitors and inducers of cytochrome P450 3A4 will affect the plasma concentration of finasteride. However, based on established safety margins, any increase due to concomitant use of such inhibitors is unlikely to be of clinical significance. Compounds which have been tested in man have included propranolol, digoxin, glibenclamide, warfarin, theophylline, and phenazone and no clinically meaningful interactions were found.

4.6 Fertility, pregnancy and lactation

Pregnancy

Finasteride is contraindicated for use in women when they are or may potentially be pregnant (See 4.3 Contraindications).

Because of the ability of type II 5 α -reductase inhibitors to inhibit conversion of testosterone to dihydrotestosterone, these drugs, including finasteride, may cause abnormalities of the external genitalia of a male fetus when administered to a pregnant woman.

Exposure to finasteride – risk to male fetus

Women should not handle crushed or broken tablets of finasteride when they are or may potentially be pregnant because of the possibility of absorption of finasteride and the subsequent potential risk to a male fetus (see 4.6 Pregnancy and lactation *Pregnancy*). Finasteride tablets are coated and will prevent contact with the active ingredient during normal handling, provided that the tablets have not been broken or crushed.

Small amounts of finasteride have been recovered from the semen in subjects receiving finasteride 5 mg/day. It is not known whether a male fetus may be adversely affected if his mother is exposed to the semen of a patient being treated with finasteride. When the patient's sexual partner is or may potentially be pregnant, the patient is recommended to minimise exposure of his partner to semen.

Breast-feeding

Finasteride is not indicated for use in women.

It is not known whether finasteride is excreted in human milk.

4.7 Effects on ability to drive and use machines

There are no data to suggest that finasteride affects the ability to drive or use machines.

4.8 Undesirable effects

The most frequent adverse reactions are impotence and decreased libido. These adverse reactions occur early in the course of therapy and resolve with continued treatment in the majority of patients.

The adverse reactions reported during clinical trials and/or post-marketing use with finasteride 5mg and/or finasteride at lower doses are listed in the table below.

Frequency of adverse reactions is determined as follows:

- Very common ($\geq 1/10$)
- Common ($\geq 1/100$ to $< 1/10$)
- Uncommon ($\geq 1/1,000$ to $< 1/100$)
- Rare ($\geq 1/10,000$ to $< 1/1,000$)
- Very rare ($< 1/10,000$)
- Not known (cannot be estimated from the available data)

The frequency of adverse reactions reported during post-marketing use cannot be determined as they are derived from spontaneous reports.

	Common	Uncommon	Not known
Immune system disorders			Hypersensitivity reactions including angioedema (including swelling of lips, tongue, throat and face)
Psychiatric disorders	decreased libido		depression, decreased libido that continued after discontinuation of treatment, anxiety, suicidal ideation

Cardiac disorders			palpitation
Hepatobiliary disorders			increased hepatic enzymes
Skin and subcutaneous tissue disorders		rash	pruritus, urticaria
Reproductive system and breast disorders	impotence	ejaculation disorder, breast tenderness, breast enlargement	sexual dysfunction (erectile dysfunction and ejaculation disorders) that continued after discontinuation of treatment; testicular pain, haemospermia, male infertility and/or poor seminal quality. Normalization or improvement of seminal quality has been reported after discontinuation of finasteride.
Investigations	decreased volume of ejaculate		

In addition, the following has been reported in clinical trials and post-marketing use: male breast cancer (see 4.4 Special warnings and precautions for use).

Medical Therapy of Prostate Symptoms (MTOPS)

The MTOPS study compared finasteride 5 mg/day (n=768), doxazosin 4 or 8 mg/day (n=756), combination therapy of finasteride 5 mg/day and doxazosin 4 or 8 mg/day (n=786), and placebo (n=737). In this study, the safety and tolerability profile of the combination therapy was generally consistent with the profiles of the individual components. The incidence of ejaculation disorder in patients receiving combination therapy was comparable to the sum of incidences of this adverse experience for the two monotherapies.

Other Long-Term Data

In a 7-year placebo-controlled trial that enrolled 18,882 healthy men, of whom 9060 had prostate needle biopsy data available for analysis, prostate cancer was detected in 803 (18.4%) men receiving finasteride 5 mg and 1147 (24.4%) men receiving placebo. In the finasteride 5 mg group, 280 (6.4%) men had prostate cancer with Gleason scores of 7-10 detected on needle biopsy vs. 237 (5.1%) men in the placebo group. Additional analyses suggest that the increase in the prevalence of high-grade prostate cancer observed in the finasteride 5 mg group may be explained by a detection bias due to the effect of finasteride 5 mg on prostate volume. Of the total cases of prostate cancer diagnosed in this study, approximately 98% were classified as intracapsular (clinical stage T1 or T2) at diagnosis. The clinical significance of the Gleason 7-10 data is unknown.

Laboratory Test Findings

When PSA laboratory determinations are evaluated, consideration should be given to the fact that PSA levels are decreased in patients treated with finasteride (see section 4.4 Special warnings and precautions for use).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

Patients have received single doses of finasteride up to 400 mg and multiple doses up to 80 mg/day for three months without adverse effects.

No specific treatment of overdosage with finasteride is recommended.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Testosterone-5 α -reductase-inhibitors
ATC code: G04CB01

Finasteride is a synthetic 4-azasteroid, a specific competitive inhibitor of the intracellular enzyme Type-II-5 α -reductase. The enzyme converts testosterone into the more potent androgen dihydrotestosterone (DHT). The prostate gland and, consequently, also the hyperplastic prostate tissue are dependent on the conversion of testosterone to DHT for their normal function and growth. Finasteride has no affinity for the androgen receptor.

Clinical studies show a rapid reduction of the serum DHT levels of 70%, which leads to a reduction of prostate volume. After 3 months, a reduction of approx. 20% in the volume of the gland occurs, and the shrinking continues and reaches approx. 27% after 3 years. Marked reduction takes place in the periurethral zone immediately surrounding the urethra. Urodynamic measurements have also confirmed a significant reduction of detrusor pressure as a result of the reduced obstruction.

Significant improvements in maximum urinary flow rate and symptoms have been obtained after a couple of weeks, compared with the start of treatment. Differences from placebo have been documented at 4 and 7 months, respectively.

All efficacy parameters have been maintained over a 3 year follow-up period.

Effects of four years treatment with finasteride on incidence of acute urine retention, need for surgery, symptom score and prostate volume:

In clinical studies of patients with moderate to severe symptoms of BPH, an enlarged prostate on digital rectal examination and low residual urinary volumes, finasteride reduced the incidence of acute retention of urine from 7/100 to 3/100 over four years and the need for surgery (TURP or prostatectomy) from 10/100 to 5/100. These reductions were associated with a 2 point improvement in QUASI-AUA symptom score (range 0–34), a sustained regression in prostate volume of approximately 20% and a sustained increase in urinary flow rate.

Medical therapy of prostatic symptoms

The Medical Therapy of Prostatic Symptoms (MTOPS) Trial was a 4- to 6-year study in 3047 men with symptomatic BPH who were randomised to receive finasteride 5 mg/day, doxazosin 4 or 8 mg/day*, the combination of finasteride 5 mg/day and doxazosin 4 or 8 mg/day*, or placebo. The primary endpoint was time to clinical progression of BPH, defined as a >4 point confirmed increase from baseline in symptom score, acute urinary retention, BPH-related renal insufficiency, recurrent urinary tract infections or urosepsis, or incontinence. Compared to placebo, treatment with finasteride, doxazosin, or combination therapy resulted in a significant reduction in the risk of clinical progression of BPH by 34 (p=0.002), 39 (p<0.001) and 67% (p<0.001), respectively. The majority of events (274 out of 351) that constituted BPH progression were confirmed ≥ 4 point increases in symptom score; the risk of symptom score progression was reduced by 30 (95% CI 6 to 48%), 46 (95% CI 25 to 60%), and 64% (95% CI 48 to 75%) in the finasteride, doxazosin, and combination groups, respectively, compared to placebo. Acute urinary retention accounted for 41 of the 351 events of BPH progression; the risk of developing acute urinary retention was reduced by 67 (p=0.011), 31 (p=0.296), and 79% (p=0.001) in the finasteride, doxazosin, and combination groups, respectively, compared to placebo.

Only the finasteride and combination therapy groups were significantly different from placebo.

* Titrated from 1 mg to 4 or 8 mg as tolerated over a 3-week period.

5.2 Pharmacokinetic properties

Absorption

The bioavailability of finasteride is approx. 80%. Peak plasma concentrations are reached approx. 2 hours after intake, and absorption is complete after 6–8 hours.

Distribution

Binding to plasma proteins is approx. 93%.

Clearance and volume of distribution are approx. 165 ml/min (70–279 ml/min) and 76 l (44–96 l), respectively. Accumulation of small amounts of finasteride is seen on repeated administration. After a daily dose of 5 mg the lowest steady-state concentration of finasteride has been calculated to be 8–10 ng/ml, which remains stable over time.

Biotransformation

Finasteride is metabolised in the liver. Finasteride does not significantly affect the cytochrome P 450 enzyme system. Two metabolites with low 5 α -reductase-inhibiting effects have been identified.

Elimination

The plasma half life is a mean of 6 hours (4–12 hours) (in men > 70 years: 8 hours, range 6–15 hours). Following administration of radioactively labelled finasteride, approx. 39% (32–46%) of the dose was excreted in the urine in the form of metabolites. Virtually no unchanged finasteride was recovered in the urine. Approx. 57% (51–64%) of the total dose was excreted in the faeces.

In the elderly, the elimination rate of finasteride is somewhat decreased. Half-life is prolonged from a mean half-life of approximately 6 hours in men aged 18–60 years to 8 hours in men aged more than 70 years. This is of no clinical significance and does not warrant a reduction in dosage.

In patients with chronic renal impairment, whose creatinine clearance ranged from 9 to 55 ml/min, the disposition of a single dose of ¹⁴C-finasteride was not different from that in healthy volunteers.

Protein binding also did not differ in patients with renal impairment. A portion of the metabolites which normally is excreted renally was excreted in the faeces. It therefore appears that faecal excretion increases commensurate to the decrease in urinary excretion of metabolites. Dosage adjustment in non-dialysed patients with renal impairment is not necessary.

Finasteride has been found to cross the blood-brain barrier. Small amounts of finasteride have been recovered in the seminal fluid of treated . In 2 studies of healthy subjects (n=69) receiving finasteride 5 mg/day for 6–24 weeks, finasteride concentrations in semen ranged from undetectable (<0.1 ng/ml) to 10.54 ng/ml. In an earlier study using a less sensitive assay, finasteride concentrations in the semen of 16 subjects receiving finasteride 5 mg/day ranged from undetectable (<1.0 ng/ml) to 21 ng/ml. Thus, based on a 5-ml ejaculate volume, the amount of finasteride in semen was estimated to be 50- to 100-fold less than the dose of finasteride (5 μ g) that had no effect on circulating DHT levels in men (see also section 5.3.)

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies of repeated dose toxicity, genotoxicity, and carcinogenic potential.

Reproduction toxicology studies in male rats have demonstrated reduced prostate and seminal vesicular weights, reduced secretion from accessory genital glands and reduced fertility index (caused by the primary pharmacological effect of finasteride). The clinical relevance of these findings is unclear.

As with other 5-alpha-reductase inhibitors, feminisation of male rat fetuses has been seen with administration of finasteride in the gestation period. Intravenous administration of finasteride to pregnant rhesus monkeys at doses up to 800 ng/day during the entire period of embryonic and foetal development resulted in no abnormalities in male fetuses. This dose is about 60–120 times higher than the estimated amount in semen of a man who have taken 5 mg finasteride, and to which a woman could be exposed via semen. In confirmation of the relevance of the Rhesus model for human foetal development, oral administration of finasteride 2 mg/kg/day (the systemic exposure (AUC) of monkeys was slightly higher (3 x) than that of men who have taken 5 mg finasteride, or approximately 1–2 million times the estimated amount of finasteride in semen) to pregnant monkeys resulted in external genital abnormalities in male fetuses. No other abnormalities were observed in male fetuses and no finasteride-related abnormalities were observed in female fetuses at any dose.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core:

Lactose monohydrate
Microcrystalline cellulose
Pregelatinised starch (maize)
Lauroyl macrogolglycerides
Sodium starch glycolate (Type A)
Magnesium stearate

Film-coating:

Hypromellose
Macrogol 6000
Titanium dioxide (E171)
Indigo carmine aluminium lake (E132)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

This medicine does not require any special storage conditions.

6.5 Nature and contents of container

Blister packs Aluminium/PVC or Aluminium/Aluminium: 7, 14, 28, 30, 49, 50, 60, 90, 98, 100 and 300 (10x30) tablets.

Plastic bottles (HDPE): 10, 30, 50, 100 and 300 tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Women who are pregnant or may become pregnant should not handle crushed or broken finasteride

tablets because of the possibility of absorption of finasteride and the subsequent potential risk to a male foetus (see section 4.6).

7. MARKETING AUTHORISATION HOLDER

<To be completed nationally>

8. MARKETING AUTHORISATION NUMBER(S)

<To be completed nationally>

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 17 February 2006

Date of latest renewal:

10. DATE OF REVISION OF THE TEXT

2025-09-01

Detailed information on this medicinal product is available on the website of {name of Member State Agency (link)}