

Package leaflet: Information for the patient

Fluoxetine Sandoz 20 mg dispersible tablets

fluoxetine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Fluoxetine Sandoz is and what it is used for
2. What you need to know before you take Fluoxetine Sandoz
3. How to take Fluoxetine Sandoz
4. Possible side effects
5. How to store Fluoxetine Sandoz
6. Contents of the pack and other information

1. What Fluoxetine Sandoz is and what it is used for

Fluoxetine Sandoz contains the active substance fluoxetine which is one of a group of medicines called selective serotonin re-uptake inhibitors (SSRI) antidepressants.

This medicine is used to treat the following conditions:

Adults:

- Major depressive episodes
- Obsessive compulsive disorder
- Bulimia nervosa: Fluoxetine Sandoz is used alongside psychotherapy for the reduction of binge-eating and purging

Children and adolescents aged 8 years and above:

- Moderate to severe major depressive disorder, if the depression does not respond to psychological therapy after 4–6 sessions. Fluoxetine Sandoz should be offered to a child or young person with moderate to severe major depressive disorder **only** in combination with psychological therapy.

How Fluoxetine Sandoz works

Everyone has a substance called serotonin in their brain. It is not fully understood how Fluoxetine Sandoz and other SSRIs work but they may help by increasing the level of serotonin in the brain.

Treating these conditions is important to help you get better. If it's not treated, your condition may not go away and may become more serious and more difficult to treat.

You may need to be treated for a few weeks or months to ensure that you are free from symptoms.

2. What you need to know before you take Fluoxetine Sandoz

Do not take Fluoxetine Sandoz if you are:

- allergic to fluoxetine or any of the other ingredients of this medicine (listed in section 6).
- If you develop a rash or other allergic reactions (like itching, swollen lips or face or shortness of breath), stop taking the tablets straight away and contact your doctor immediately.
- taking other medicines known as irreversible, non-selective monoamine oxidase inhibitors (MAOIs), since serious or even fatal reactions can occur (e.g. iproniazid used to treat depression). Treatment with Fluoxetine Sandoz should only be started at least 2 weeks after discontinuation of an irreversible, non-selective MAOI.

Do not take any irreversible, non-selective MAOIs for at least 5 weeks after you stop taking Fluoxetine Sandoz. If Fluoxetine Sandoz has been prescribed for a long period and/or at a high dose, a longer interval needs to be considered by your doctor.

- taking metoprolol (to treat heart failure) since there is an increased risk of your heart beat becoming too slow.

Warnings and precautions

Talk to your doctor or pharmacist before taking Fluoxetine Sandoz and if any of the following applies to you:

- heart problems;
- appearance of fever, muscle stiffness or tremor, changes in your mental state like confusion, irritability and extreme agitation; you may suffer from the so-called “serotonin syndrome” or “neuroleptic malignant syndrome”. Although this syndrome occurs rarely it may result in potentially life threatening conditions; **contact your doctor immediately**, since Fluoxetine Sandoz might need to be discontinued.
- mania now or in the past; if you have a manic episode, contact your doctor immediately because Fluoxetine Sandoz might need to be discontinued;
- history of bleeding disorders or appearance of bruises or unusual bleeding, or if you are pregnant (see ‘Pregnancy’);
- ongoing treatment with medicines that thin the blood (see ‘Other medicines and Fluoxetine Sandoz’).
- epilepsy or fits. If you have a fit (seizures) or experience an increase in seizure frequency, contact your doctor immediately; Fluoxetine Sandoz might need to be discontinued;
- ongoing ECT (electro-convulsive therapy);
- ongoing treatment with tamoxifen (used to treat breast cancer) (see ‘Other medicines and Fluoxetine Sandoz’);
- starting to feel restless and cannot sit or stand still (akathisia). Increasing your dose of Fluoxetine Sandoz may make this worse;
- diabetes (your doctor may need to adjust your dose of insulin or other antidiabetic treatment);
- liver problems (your doctor may need to adjust your dose);
- low resting heart-rate and/or if you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets);
- ongoing treatment with diuretics (water tablets), especially if you are elderly;
- glaucoma (increased pressure in the eye).

Medicines like Fluoxetine Sandoz (so called SSRIs/SNRIs) may cause symptoms of sexual dysfunction (see section 4). In some cases, these symptoms have continued after stopping treatment.

Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- If you have previously had thoughts about killing or harming yourself.
- If you are a young adult. Information from clinical trials has shown an increased risk of suicidal behaviour in adults aged less than 25 years with psychiatric conditions who were treated with an antidepressant.

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away.**

You may find it helpful to tell a relative or close friend that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

Children and adolescents aged 8 to 18 years:

Patients under 18 have an increased risk of side-effects such as suicide attempt, suicidal thoughts and hostility (predominantly aggression, oppositional behaviour and anger) when they take this class of medicines. Fluoxetine Sandoz should only be used in children and adolescents aged 8 to 18 years for the treatment of moderate to severe major depressive episodes (in combination with psychological therapy) and it should not be used to treat other conditions.

Additionally, only limited information concerning the long-term safety of Fluoxetine Sandoz on growth, puberty, mental, emotional and behavioural development in this age group is available. Despite this, and if you are a patient under 18, your doctor may prescribe Fluoxetine Sandoz for moderate to severe major depressive episodes, in combination with psychological therapy because he/she decides that this is in your best interests. If your doctor has prescribed Fluoxetine Sandoz for a patient under 18 and you want to discuss this, please go back to your doctor. You should inform your doctor if any of the symptoms listed above develop or worsen when patients under 18 are taking Fluoxetine Sandoz.

Fluoxetine Sandoz should not be used in the treatment of children under the age of 8 years.

Other medicines and Fluoxetine Sandoz

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Do not take Fluoxetine Sandoz with:

- Certain **irreversible, non-selective monoamine oxidase inhibitors (MAOIs)**, some used to treat depression. Irreversible, non-selective MAOIs must not be used with Fluoxetine Sandoz as serious or even fatal reactions (serotonin syndrome) can occur (see section “Do not take Fluoxetine Sandoz”). Treatment with Fluoxetine Sandoz should only be started at least 2 weeks after discontinuation of an irreversible, non-selective MAOI (for instance tranylcypromine). **Do not** take any irreversible, non-selective MAOIs for at least 5 weeks after you stop taking Fluoxetine Sandoz. If Fluoxetine Sandoz has been prescribed for a long period and/or at a high dose, a longer interval than 5 weeks may need to be considered by your doctor.
- **metoprolol** when used for heart failure; there is an increased risk of your heart beat becoming too slow.

Fluoxetine Sandoz may affect the way the following medicines work (interaction):

- **tamoxifen** (used to treat breast cancer), because Fluoxetine Sandoz may change the blood levels of this drug, resulting in the possibility of a reduction in the effect of tamoxifen, your doctor may need to consider prescribing a different antidepressant treatment.
- **monoamine oxidase inhibitors A (MAOI-A)** including linezolid (an antibiotic) and methylthioninium chloride (also called methylene blue used to <indication(s) authorised in member state>): due to the risk of serious or even fatal reactions (called serotonin syndrome). Treatment with fluoxetine can be started the day after stopping treatment with reversible MAOIs but the doctor may wish to monitor you carefully and use a lower dose of the MAOI-A drug.
- **mequitazine** (for allergies); because taking this drug with Fluoxetine Sandoz may increase the risk of changes in the electrical activity of the heart.

- **phenytoin** (for epilepsy); because Fluoxetin Sandoz may influence the blood levels of this drug, your doctor may need to introduce phenytoin more carefully and carry out check-ups when given with Fluoxetin Sandoz.
lithium, selegiline, St. John's Wort, tramadol (a painkiller), **buprenorphine, triptans** (for migraine) and **tryptophan**; there is an increased risk of mild serotonin syndrome when these drugs are taken with Fluoxetin Sandoz. Your doctor will carry out more frequent check-ups.
- medicines that may affect the heart's rhythm, e.g. **Class IA and III antiarrhythmics, antipsychotics** (e.g. phenothiazine derivatives, pimozide, haloperidol), **tricyclic antidepressants**, certain **antimicrobial agents** (e.g. sparfloxacin, moxifloxacin, erythromycin IV, pentamidine), **anti-malaria treatment** particularly halofantrine or certain **antihistamines** (astemizole, mizolastine), because taking one or more of these drugs with Fluoxetin Sandoz may increase the risk of changes in the electrical activity of the heart.
- **anti-coagulants** (such as warfarin), **NSAID** (such as ibuprofen, diclofenac), **aspirin** and other medicines which can thin the blood (including clozapine, used to treat certain mental disorders). Fluoxetin Sandoz may alter the effect of these medicines on the blood. If Fluoxetin Sandoz treatment is started or stopped when you are taking warfarin, your doctor will need to perform certain tests, adjust your dose and check on you more frequently.
- **cypheptadine** (for allergies); because it may reduce the effect of Fluoxetin Sandoz.
- **drugs that lower sodium levels in the blood** (including, drug that causes increase in urination, desmopressin, carbamazepine and oxcarbazepine); because these drugs may increase the risk of sodium levels in the blood becoming too low when taken with Fluoxetin Sandoz.
- antidepressants such as tricyclic anti-depressants, other selective serotonin reuptake inhibitors (SSRIs) or bupropion, mefloquine or chloroquine (used to treat malaria), tramadol (used to treat severe pain) or **anti-psychotics** such as phenothiazines or butyrophenones; because Fluoxetin Sandoz may increase the risk of seizures when taken with these medicines.
- flecainide, propafenone, nebivolol or **encainide** (for heart problems), carbamazepine (for epilepsy), **atomoxetine** or tricyclic antidepressants (for example imipramine, desipramine and amitriptyline) or risperidone (for schizophrenia); because Fluoxetin Sandoz may possibly change the blood levels of these medicines, your doctor may need to lower their dose when administered with Fluoxetin Sandoz.

Fluoxetin Sandoz with food, drink and alcohol

You can take Fluoxetin Sandoz with or without food, whatever you prefer.

You should avoid alcohol while you are taking this medicine.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

This medicine contains benzyl alcohol (see **Fluoxetin Sandoz contains lactose, glucose, sodium and benzyl alcohol**).

Pregnancy

Talk to your doctor as soon as possible if you're pregnant, if you might be pregnant, or if you're planning to become pregnant.

In babies whose mothers took fluoxetine during the first few months of pregnancy, there have been some studies describing an increased risk of birth defects affecting the heart. In the general population, about 1 in 100 babies are born with a heart defect. This increased to about 2 in 100 babies in mothers who took fluoxetine.

When taken during pregnancy, particularly in the last 3 months of pregnancy, medicines like fluoxetine may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear bluish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

It is preferable not to use this treatment during pregnancy unless the potential benefit outweighs the potential risk. Thus, you and your doctor may decide to gradually stop taking Fluoxetine Sandoz while you are pregnant or before being pregnant. However, depending on your circumstances, your doctor may suggest that it is better for you to keep taking Fluoxetine Sandoz.

Caution should be exercised when used during pregnancy, especially during late pregnancy or just before giving birth since the following effects have been reported in new born children: irritability, tremor, muscle weakness, persistent crying, and difficulty in sucking or in sleeping.

If you take Fluoxetine Sandoz near the end of your pregnancy there may be an increased risk of heavy vaginal bleeding shortly after birth, especially if you have a history of bleeding disorders. Your doctor or midwife should be aware that you are taking Fluoxetine Sandoz so they can advise you.

Breast-feeding

Fluoxetine is excreted in breast milk and can cause side effects in babies. You should only breast-feed if it is clearly necessary. If breast-feeding is continued, your doctor may prescribe a lower dose of fluoxetine.

Fertility

Fluoxetine has been shown to reduce the quality of sperm in animal studies. Theoretically, this could affect fertility, but impact on human fertility has not been observed as yet.

Driving and using machines

Psychotropic drugs such as Fluoxetine Sandoz may affect your judgment or co-ordination. Do not drive or use machinery until you know how Fluoxetine Sandoz affects you.

Fluoxetine Sandoz contains lactose, glucose, sodium and benzyl alcohol

This medicine contains lactose and glucose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per dispersible tablet, that is to say essentially 'sodium-free'.

This medicine contains 0.027 mg benzyl alcohol in each dispersible tablet. Benzyl alcohol may cause allergic reactions. Ask your doctor or pharmacist for advice if you are pregnant or breast-feeding, if you have a liver or kidney disease. This is because large amounts of benzyl alcohol can build-up in your body and may cause side effects (called "metabolic acidosis").

3. How to use Fluoxetine Sandoz

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. Do not take more tablets than your doctor tells you.

Fluoxetine can be taken as a single dose or divided into several doses.

Fluoxetine Sandoz dispersible tablets can be dissolved in a glass of water, divided into equal doses, or swallowed whole.

Adults:

The recommended dose is:

- **Depression:**

The recommended dose is 1 tablet (20 mg) daily.

Your doctor will review and adjust your dose if necessary within 3 to 4 weeks of the start of treatment. If required, the dose can be gradually increased up to a maximum of 3 tablets (60 mg) daily.

The dose should be increased carefully to ensure that you receive the lowest effective dose.

You may not feel better immediately when you first start taking your medicine for depression. This is usual because an improvement in depressive symptoms may not occur until after the first few weeks.

Patients with depression should be treated for at least 6 months.

- **Bulimia nervosa:** The recommended dose is 3 tablets (60 mg) daily.
- **Obsessive compulsive disorder:**
The recommended dose is 1 tablet (20 mg) daily.

Your doctor will review and adjust your dose if necessary after 2 weeks of treatment. If required, the dose can be gradually increased up to a maximum of 3 tablets (60 mg) daily. If no improvement is noted within 10 weeks, your doctor will reconsider your treatment.

Use in children and adolescents aged 8 to 18 years with depression:

Treatment should be started and be supervised by a specialist. The starting dose is 10 mg/day. After 1 to 2 weeks, your doctor may increase the dose to 20 mg/day. The dose should be increased carefully to ensure that you receive the lowest effective dose.

Lower weight children may need lower doses.

If there is a satisfactory response to treatment, your doctor will review the need for continuing treatment beyond 6 months. If you have not improved within 9 weeks, your doctor will reassess your treatment.

Elderly:

Your doctor will increase the dose with more caution and the daily dose should generally not exceed 2 tablets (40 mg). The maximum dose is 3 tablets (60 mg) daily.

Hepatic impairment

If you have a liver problem or are using other medicine that might affect Fluoxetin Sandoz, your doctor may decide to prescribe a lower dose or tell you to use Fluoxetin Sandoz every other day.

If you take more Fluoxetin Sandoz than you should

- If you take too many tablets, go to your nearest hospital emergency department (or casualty) or tell your doctor straight away.
- Take the pack of Fluoxetin Sandoz with you if you can.

Symptoms of overdose include: nausea, vomiting, seizures, heart problems (like irregular heart beat and cardiac arrest), lung problems and change in mental condition ranging from agitation to coma.

If you forget to take Fluoxetin Sandoz

- If you miss a dose, do not worry. Take your next dose the next day at the usual time. Do not take a double dose to make up for a forgotten dose.
- Taking your medicine at the same time each day may help you to remember to take it regularly.

If you stop taking Fluoxetin Sandoz

Do not stop taking

- without asking your doctor first, even when you start to feel better. It is important that you keep taking your medicine.
- Make sure you do not run out of tablets.

You may notice the following effects (withdrawal effects) when you stop taking Fluoxetine Sandoz: dizziness; tingling feelings like pins and needles; sleep disturbances (vivid dreams, nightmares, inability to sleep); feeling restless or agitated; unusual tiredness or weakness; feeling anxious; nausea/vomiting (feeling sick or being sick); tremor (shakiness); headaches.

Most people find that any symptoms on stopping Fluoxetine Sandoz are mild and disappear within a few weeks. If you experience symptoms when you stop treatment, contact your doctor.

When stopping Fluoxetine Sandoz, your doctor will help you to reduce your dose slowly over one or two weeks - this should help reduce the chance of withdrawal effects.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

- If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away** (see section 2).
- If you get a **rash or allergic reaction** such as itching, swollen lips/tongue or wheezing/shortness of breath, **stop taking the tablets straight away and tell your doctor immediately**.
- If you **feel restless** and cannot sit or stand still you may have akathisia; increasing your dose of Fluoxetine Sandoz may make you feel worse. If you feel like this, **contact your doctor**.
- **Tell your doctor immediately** if your skin starts to turn red or you develop a varied skin reaction or your skin starts to blister or peel. This is very rare.

The most frequent side effects (very common side effects that may affect more than 1 in 10 people) are insomnia, headache, diarrhoea, feeling sick (nausea) and fatigue.

Some patients have had:

- a combination of symptoms (known as “serotonin syndrome”) including unexplained fever with faster breathing or heart rate, sweating, muscle stiffness or tremor, confusion, extreme agitation or sleepiness (only rarely);
- feelings of weakness, drowsiness or confusion mostly elderly people and in (elderly) people taking diuretics (water tablets);
- prolonged and painful erection;
- irritability and extreme agitation.
- heart problems, such as fast or irregular heart rate, fainting, collapsing or dizziness upon standing which may indicate abnormal functioning of the heart rate.

If you have any of the above side effects, you should tell your doctor immediately.

The following side effects have also been reported in patients taking Fluoxetine Sandoz:

Common (may affect up to 1 in 10 people)

- not feeling hungry, weight loss
- nervousness, anxiety
- restlessness, poor concentration
- feeling tense
- decreased sex drive or sexual problems (including difficulty maintaining an erection for sexual activity)
- sleep problems, unusual dreams, tiredness or sleepiness

- dizziness
- change in taste
- uncontrollable shaking movements
- blurred vision
- rapid and irregular heartbeat sensations
- flushing
- yawning
- indigestion, vomiting
- dry mouth
- rash, urticaria, itching
- excessive sweating
- joint pain
- passing urine more frequently- unexplained vaginal bleeding
- feeling shaky or chills

Uncommon (may affect up to 1 in 100 people)

- feeling detached from yourself
- strange thinking
- abnormally high mood
- orgasm problems
- thoughts of suicide or harming yourself
- teeth grinding
- muscle twitching, involuntary movements or problems with balance or co-ordination
- memory impairment
- enlarged (dilated) pupils
- ringing in the ears
- low blood pressure
- shortness of breath
- nose bleeds
- difficulty swallowing
- hair loss
- increased tendency to bruising
- unexplained bruising or bleeding
- cold sweat
- difficulty passing urine
- feeling hot or cold
- abnormal liver test results

Rare (may affect up to 1 in 1,000 people)

- low levels of salt in the blood
- reduction in blood platelets, which increases risk of bleeding or bruising
- reduction in white blood cell count
- untypical wild behaviour
- hallucinations
- agitation
- panic attacks
- confusion
- stuttering
- aggression
- fits
- vasculitis (inflammation of a blood vessel)
- rapid swelling of the tissues around the neck, face, mouth and/or throat
- pain in the tube that takes food or water to your stomach

- hepatitis
- lung problems
- sensitivity to sunlight
- muscle pain
- problems urinating
- producing breast milk

Not known (frequency cannot be estimated from the available data)

- heavy vaginal bleeding shortly after birth (postpartum haemorrhage), see Pregnancy in section 2 for more information

Bone fractures - An increased risk of bone fractures has been observed in patients taking this type of medicines.

Most of these side effects are likely to disappear with continued treatment.

Additionally in children and adolescents (8-18 years) – In addition to the possible side effects listed above, fluoxetine may slow growth or possibly delay sexual maturity. Suicide-related behaviours (suicide attempt and suicidal thoughts), hostility, mania and nose bleeds were also commonly reported in children.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V*. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Fluoxetin Sandoz

Keep this medicine out of the sight and reach of children.

This medicine does not require any special storage conditions.

Do not use this medicine after the expiry date which is stated on the carton and blister after “EXP”. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Fluoxetin Sandoz contains

- The active substance is fluoxetine hydrochloride equivalent to 20 mg fluoxetine.
- The other ingredients are lactose monohydrate, microcrystalline cellulose, pregelatinised maize starch, croscarmellose sodium, colloidal anhydrous silica, sodium saccharin, sodium cyclamate, magnesium stearate, plum flavour (contains glucose, silicon dioxide, 1,2-propylene glycol, benzyl alcohol), peppermint flavour (contains maltodextrin, gum arabic).

What Fluoxetin Sandoz looks like and contents of the pack

Fluoxetin Sandoz 20 mg dispersible tablets are white, round and biconvex tablets, one side with a score line.

The dispersible tablets are packed in aluminium /aluminium blisters and inserted in a carton.

Pack sizes:

7, 10, 14, 20, 28, 30, 60, 70, 100 and 100x1 dispersible tablets

Not all pack sizes may be marketed

Marketing Authorisation Holder and Manufacturer

[To be completed nationally]

This medicine is authorised in the Member States of the European Economic Area under the following names:

[To be completed nationally]

This leaflet was last revised in 2024-01-31.

[To be completed nationally]