

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Gardette mite, 80 micrograms/4.5 micrograms/inhalation, inhalation powder.

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each delivered dose (the dose that leaves the mouthpiece) contains: budesonide 80 micrograms/inhalation and formoterol fumarate dihydrate 4.5 micrograms/inhalation.

Each metered dose contains: budesonide 100 micrograms/inhalation and formoterol fumarate dihydrate 6 micrograms/inhalation.

### Excipient with known effect

Lactose monohydrate 810 micrograms per delivered dose.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Inhalation powder.

White powder.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Gardette mite is indicated in adults, adolescents, and children aged 6 years and older.

Gardette is indicated in the regular treatment of asthma where use of a combination (inhaled corticosteroid and long-acting  $\beta_2$  adrenoceptor agonist) is appropriate:

- patients not adequately controlled with inhaled corticosteroids and “as needed” inhaled short-acting  $\beta_2$  adrenoceptor agonists.
- or
- patients already adequately controlled on both inhaled corticosteroids and long-acting  $\beta_2$  adrenoceptor agonists.

**Note:** Gardette mite (80 micrograms/4.5 micrograms/inhalation) is not appropriate in patients with severe asthma.

### 4.2 Posology and method of administration

Route of administration: For inhalation use

#### Posology

Gardette is not intended for the initial management of asthma.

The dosage of the components of Gardette is individual and should be adjusted to the severity of the disease. This should be considered not only when treatment with combination products is initiated but also when the maintenance dose is adjusted. If an individual patient should require a combination of doses other than those available in the combination inhaler, appropriate doses of  $\beta_2$  adrenoceptor agonists and/or corticosteroids by individual inhalers should be prescribed.

The dose should be titrated to the lowest dose at which effective control of symptoms is maintained. Patients should be regularly reassessed by their prescriber/health care provider so that the dosage of Gardette remains optimal. When long-term control of symptoms is maintained with the lowest recommended dosage, then the next step could include a test of inhaled corticosteroid alone.

For Gardette there are two treatment approaches:

**A. Gardette maintenance therapy:** Gardette is taken as regular maintenance treatment with a separate rapid-acting bronchodilator as rescue.

**B. Gardette maintenance and reliever therapy:** Gardette is taken as regular maintenance treatment and as needed in response to symptoms.

**A. Gardette maintenance therapy**

Patients should be advised to have their separate rapid-acting bronchodilator available for rescue use at all times.

*Recommended doses:*

*Adults (18 years and older):* 1-2 inhalations twice daily. Some patients may require up to a maximum of 4 inhalations twice daily.

*Adolescents (12-17 years):* 1-2 inhalations twice daily.

*Children (6 years and older):* 2 inhalations twice daily.

In usual practice when control of symptoms is achieved with the twice daily regimen, titration to the lowest effective dose could include Gardette mite given once daily, when in the opinion of the prescriber, a long-acting bronchodilator in combination with an inhaled corticosteroid would be required to maintain control.

Increasing use of a separate rapid acting bronchodilator indicates a worsening of the underlying condition and warrants a reassessment of the asthma therapy.

*Children under 6 years:* As only limited data are available, Gardette mite is not recommended for children younger than 6 years.

**B. Gardette maintenance and reliever therapy**

Patients take a daily maintenance dose of Gardette and in addition take Gardette as needed in response to symptoms. Patients should be advised to always have Gardette available for rescue use.

Gardette maintenance and reliever therapy should especially be considered for patients with:

- inadequate asthma control and in frequent need of reliever medication
- asthma exacerbations in the past requiring medical intervention

Close monitoring for dose-related adverse effects is needed in patients who frequently take high numbers of Gardette as-needed inhalations.

*Recommended doses:*

*Adults and adolescents (12 years and older):* The recommended maintenance dose is 2 inhalations per day, given either as one inhalation in the morning and evening or as 2 inhalations in either the morning or evening. Patients should take 1 additional inhalation as needed in response to symptoms. If symptoms persist after a few minutes, an additional inhalation should be taken. Not more than 6 inhalations should be taken on any single occasion.

A total daily dose of more than 8 inhalations is not normally needed; however, a total daily dose of up to 12 inhalations could be used for a limited period. Patients using more than 8 inhalations daily should be strongly recommended to seek medical advice. They should be reassessed and their maintenance therapy should be reconsidered.

*Children under 12 years:* Gardette maintenance and reliever therapy is not recommended for children.

### **General information**

#### *Special patient groups:*

There are no special dosing requirements for elderly patients. There are no data available for use of Gardette in patients with hepatic or renal impairment. As budesonide and formoterol are primarily eliminated via hepatic metabolism, an increased exposure can be expected in patients with severe liver cirrhosis.

#### Method of administration

##### *Instructions for correct use of Gardette mite:*

The inhaler is inspiratory flow-driven, which means that when the patient inhales through the mouthpiece, the substance will follow the inspired air into the airways.

**Note:** It is important to instruct the patient

- to carefully read the instructions for use in the patient information leaflet which is packed together with each Gardette mite Inhaler.
- to breathe in forcefully and deeply through the mouthpiece to ensure that an optimal dose is delivered to the lungs.
- never to breathe out through the mouthpiece.
- to replace the cover of the Gardette mite Inhaler after use.
- to rinse their mouth out with water after inhaling the maintenance dose to minimise the risk of oropharyngeal thrush. If oropharyngeal thrush occurs, patients should also rinse their mouth with water after the as-needed inhalations.

The patient may not taste or feel any medication when using Gardette mite Inhaler due to the small amount of drug dispensed.

### **4.3 Contraindications**

Hypersensitivity to the active substances or to the excipient listed in section 6.1 (lactose, which contains small amounts of milk protein).

### **4.4 Special warnings and precautions for use**

#### Dosing advice

Once asthma symptoms are controlled, consideration may be given to gradually reducing the dose of Gardette. Regular review of patients as treatment is stepped down is important. The lowest effective dose of Gardette should be used (see section 4.2).

Patients should be advised to have their rescue inhaler available at all times, either Gardette (for patients using Gardette as maintenance and reliever therapy) or a separate rapid-acting bronchodilator (for patients using Gardette as maintenance therapy only).

Patients should be reminded to take their Gardette maintenance dose as prescribed, even when asymptomatic. The prophylactic use of Gardette, e.g. before exercise, has not been studied. The reliever inhalations of Gardette should be taken in response to asthma symptoms but are not intended for regular prophylactic use, e.g. before exercise. For such use, a separate rapid-acting bronchodilator should be considered.

To minimise the risk of oropharyngeal candida infection (see section 4.8), the patient should be instructed to rinse their mouth out with water after inhaling the maintenance dose. If oropharyngeal thrush occurs, patients should also rinse their mouth with water after the as-needed inhalations.

It is recommended that the dose is tapered when the treatment is discontinued and should not be stopped abruptly.

### Deterioration of disease

Serious asthma-related adverse events and exacerbations may occur during treatment with Gardette. Patients should be asked to continue treatment but to seek medical advice if asthma symptoms remain uncontrolled or worsen after initiation of Gardette.

If patients find the treatment ineffective, or exceed the highest recommended dose of Gardette, medical attention must be sought (see section 4.2). Sudden and progressive deterioration in control of asthma is potentially life threatening and the patient should undergo urgent medical assessment. In this situation consideration should be given to the need for increased therapy with corticosteroids, e.g. a course of oral corticosteroids, or antibiotic treatment if an infection is present.

Patients should not be initiated on Gardette during an exacerbation, or if they have significantly worsening or acutely deteriorating asthma.

### Transfer from oral therapy

If there is any reason to suppose that adrenal function is impaired from previous systemic steroid therapy, care should be taken when transferring patients to Gardette therapy.

The benefits of inhaled budesonide therapy would normally minimise the need for oral steroids, but patients transferring from oral steroids may remain at risk of impaired adrenal reserve for a considerable time. Recovery may take a considerable amount of time after cessation of oral steroid therapy and hence oral steroid-dependent patients transferred to inhaled budesonide may remain at risk from impaired adrenal function for some considerable time. In such circumstances HPA axis function should be monitored regularly.

During transfer from oral therapy to Gardette mite, a generally lower systemic steroid action will be experienced which may result in the appearance of allergic or arthritic symptoms such as rhinitis, eczema and muscle and joint pain. Specific treatment should be initiated for these conditions. A general insufficient glucocorticosteroid effect should be suspected if, in rare cases, symptoms such as tiredness, headache, nausea and vomiting should occur. In these cases a temporary increase in the dose of oral glucocorticosteroids is sometimes necessary.

### Excipients

Gardette contains lactose monohydrate (< 1 mg/inhalation). This amount does not normally cause problems in lactose intolerant people. The excipient lactose contains small amounts of milk proteins, which may cause allergic reactions.

### Interactions with other medicinal products

Concomitant treatment with itraconazole, ritonavir or other potent CYP3A4 inhibitors should be avoided (see section 4.5). If this is not possible the time interval between administration of the interacting drugs should be as long as possible. In patients using potent CYP3A4 inhibitors, Gardette maintenance and reliever therapy is not recommended.

### Caution with special diseases

Gardette should be administered with caution in patients with thyrotoxicosis, pheochromocytoma, diabetes mellitus, untreated hypokalaemia, hypertrophic obstructive cardiomyopathy, idiopathic subvalvular aortic stenosis, severe hypertension, aneurysm or other severe cardiovascular disorders, such as ischaemic heart disease, tachyarrhythmias or severe heart failure.

Caution should be observed when treating patients with prolongation of the QTc-interval. Formoterol itself may induce prolongation of the QTc-interval.

Potentially serious hypokalaemia may result from high doses of  $\beta_2$  adrenoceptor agonists. Concomitant treatment of  $\beta_2$  adrenoceptor agonists with drugs which can induce hypokalaemia or potentiate a hypokalaemic effect, e.g. xanthine derivatives, steroids and diuretics, may add to a possible hypokalaemic effect of the  $\beta_2$  adrenoceptor agonist. Particular caution is recommended in

unstable asthma with variable use of rescue bronchodilators, in acute severe asthma as the associated risk may be augmented by hypoxia and in other conditions when the likelihood for hypokalaemia is increased. It is recommended that serum potassium levels are monitored during these circumstances.

As for all  $\beta_2$  adrenoceptor agonists, additional blood glucose controls should be considered in diabetic patients.

The need for, and dose of inhaled corticosteroids should be re-evaluated in patients with active or quiescent pulmonary tuberculosis, fungal and viral infections in the airways.

#### Systemic effects

Systemic effects may occur with any inhaled corticosteroid, particularly at high doses prescribed for long periods. These effects are much less likely to occur with inhalation treatment than with oral corticosteroids. Possible systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression, growth retardation in children and adolescents, decrease in bone mineral density, cataract and glaucoma, and more rarely, a range of psychological or behavioural effects including psychomotor hyperactivity, sleep disorders, anxiety, depression or aggression (particularly in children) (see section 4.8).

Long-term studies with inhaled budesonide in children at mean daily doses of 400 micrograms (metered dose) or in adults at daily doses of 800 micrograms (metered dose) have not shown any significant effects on bone mineral density. No information regarding the effect of Gardette at higher doses is available.

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes, which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR), which have been reported after use of systemic and topical corticosteroids.

#### Adrenal function

Treatment with supplementary systemic steroids or inhaled budesonide should not be stopped abruptly.

Prolonged treatment with high doses of inhaled corticosteroids, particularly higher than recommended doses, may also result in clinically significant adrenal suppression. Therefore, additional systemic corticosteroid cover should be considered during periods of stress such as severe infections or elective surgery. Rapid reduction in the dose of steroids can induce acute adrenal crisis. Symptoms and signs which might be seen in acute adrenal crisis may be somewhat vague but may include anorexia, abdominal pain, weight loss, tiredness, headache, nausea, vomiting, decreased level of consciousness, seizures, hypotension and hypoglycaemia.

#### Paradoxical bronchospasm

As with other inhalation therapy, paradoxical bronchospasm may occur, with an immediate increase in wheezing and shortness of breath after dosing. If the patient experiences paradoxical bronchospasm Gardette should be discontinued immediately, the patient should be assessed and an alternative therapy instituted if necessary. Paradoxical bronchospasm responds to a rapid-acting inhaled bronchodilator and should be treated straightaway (see section 4.8).

#### Paediatric population

It is recommended that the height of children receiving prolonged treatment with inhaled corticosteroids is regularly monitored. If growth is slowed, therapy should be re-evaluated with the aim of reducing the dose of inhaled corticosteroid to the lowest dose at which effective control of asthma is maintained, if possible. The benefits of the corticosteroid therapy and the possible risks of growth suppression must be carefully weighed. In addition consideration should be given to referring the patient to a paediatric respiratory specialist.

Limited data from long-term studies suggest that most children and adolescents treated with inhaled budesonide will ultimately achieve their adult target height. However, an initial small but transient reduction in growth (approximately 1 cm) has been observed. This generally occurs within the first year of treatment.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

##### *Pharmacokinetic interactions*

Potent inhibitors of CYP3A4 (e.g. ketoconazole, itraconazole, voriconazole, posaconazole, clarithromycin, telithromycin, nefazodone and HIV protease inhibitors) are likely to markedly increase plasma levels of budesonide and concomitant use should be avoided. If this is not possible the time interval between administration of the inhibitor and budesonide should be as long as possible (see section 4.4). In patients using potent CYP3A4 inhibitors, Gardette maintenance and reliever therapy is not recommended.

The potent CYP3A4 inhibitor ketoconazole, 200 mg once daily, increased plasma levels of concomitantly orally administered budesonide (single dose of 3 mg) on average six-fold. When ketoconazole was administered 12 hours after budesonide the concentration was on average increased only three-fold showing that separation of the administration times can reduce the increase in plasma levels. Limited data about this interaction for high-dose inhaled budesonide indicates that marked increase in plasma levels (on average four fold) may occur if itraconazole, 200 mg once daily, is administered concomitantly with inhaled budesonide (single dose of 1000 µg).

##### *Pharmacodynamic interactions*

Beta-adrenergic blockers can weaken or inhibit the effect of formoterol. Gardette should therefore not be given together with beta-adrenergic blockers (including eye drops) unless there are compelling reasons.

Concomitant treatment with quinidine, disopyramide, procainamide, phenothiazines, antihistamines (terfenadine) and tricyclic antidepressants can prolong the QTc-interval and increase the risk of ventricular arrhythmias.

In addition L-Dopa, L-thyroxine, oxytocin and alcohol can impair cardiac tolerance towards  $\beta_2$ -sympathomimetics.

Concomitant treatment with monoamine oxidase inhibitors including agents with similar properties such as furazolidone and procarbazine may precipitate hypertensive reactions.

There is an elevated risk of arrhythmias in patients receiving concomitant anaesthesia with halogenated hydrocarbons.

Concomitant use of other beta-adrenergic drugs or anticholinergic drugs can have a potentially additive bronchodilating effect.

Hypokalaemia may increase the disposition towards arrhythmias in patients who are treated with digitalis glycosides.

Hypokalaemia may result from beta<sub>2</sub>-agonist therapy and may be potentiated by concomitant treatment with xanthine derivatives, corticosteroids and diuretics (see section 4.4).

Budesonide and formoterol have not been observed to interact with any other drugs used in the treatment of asthma.

##### Paediatric population

Interaction studies have only been performed in adults.

#### **4.6 Fertility, pregnancy and lactation**

### Pregnancy

For Gardette or the concomitant treatment with formoterol and budesonide, no clinical data on exposed pregnancies are available. Data from an embryo-fetal development study in the rat showed no evidence of any additional effect from the combination.

There are no adequate data from use of formoterol in pregnant women. In animal studies formoterol has caused adverse effects in reproduction studies at very high systemic exposure levels (see section 5.3).

Data on approximately 2000 exposed pregnancies indicate no increased teratogenic risk associated with the use of inhaled budesonide. In animal studies glucocorticosteroids have been shown to induce malformations (see section 5.3). This is not likely to be relevant for humans given recommended doses.

Animal studies have also identified an involvement of excess prenatal glucocorticoids in increased risks for intrauterine growth retardation, adult cardiovascular disease and permanent changes in glucocorticoid receptor density, neurotransmitter turnover and behaviour at exposures below the teratogenic dose range.

During pregnancy, Gardette should only be used when the benefits outweigh the potential risks. The lowest effective dose of budesonide needed to maintain adequate asthma control should be used.

### Breastfeeding

Budesonide is excreted in breast milk. However, at therapeutic doses no effects on the suckling child are anticipated. It is not known whether formoterol passes into human breast milk. In rats, small amounts of formoterol have been detected in maternal milk. Administration of Gardette to women who are breastfeeding should only be considered if the expected benefit to the mother is greater than any possible risk to the child.

### Fertility

There is no data available on the potential effect of budesonide on fertility. Animal reproduction studies with formoterol have shown a somewhat reduced fertility in male rats at high systemic exposure (see section 5.3).

## **4.7 Effects on ability to drive and use machines**

Gardette has no or negligible influence on the ability to drive and use machines.

## **4.8 Undesirable effects**

Since Gardette contains both budesonide and formoterol, the same pattern of undesirable effects as reported for these substances may occur. No increased incidence of adverse reactions has been seen following concurrent administration of the two compounds. The most common drug related adverse reactions are pharmacologically predictable side effects of  $\beta_2$  adrenoceptor agonist therapy, such as tremor and palpitations. These tend to be mild and usually disappear within a few days of treatment.

Adverse reactions, which have been associated with budesonide or formoterol, are given below, listed by system organ class and frequency. Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ), rare ( $\geq 1/10\ 000$  to  $< 1/1000$ ) and very rare ( $< 1/10\ 000$ ).

**Table 1**

| <b>SOC</b>                  | <b>Frequency</b> | <b>Adverse Drug reaction</b>                      |
|-----------------------------|------------------|---------------------------------------------------|
| Infections and infestations | Common           | Candida infections in the oropharynx              |
| Immune system disorders     | Rare             | Immediate and delayed hypersensitivity reactions, |



|                                                 |           |                                                                                               |
|-------------------------------------------------|-----------|-----------------------------------------------------------------------------------------------|
|                                                 |           | e.g. exanthema, urticaria, pruritus, dermatitis, angioedema and anaphylactic reaction         |
| Endocrine disorders                             | Very rare | Cushing's syndrome, adrenal suppression, growth retardation, decrease in bone mineral density |
| Metabolism and nutrition disorders              | Rare      | Hypokalaemia                                                                                  |
|                                                 | Very rare | Hyperglycaemia                                                                                |
| Psychiatric disorders                           | Uncommon  | Aggression, psychomotor hyperactivity, anxiety, sleep disorders                               |
|                                                 | Very rare | Depression, behavioural changes (predominantly in children)                                   |
| Nervous system disorders                        | Common    | Headache, tremor                                                                              |
|                                                 | Uncommon  | Dizziness                                                                                     |
|                                                 | Very rare | Taste disturbances                                                                            |
| Eye disorders                                   | Uncommon  | Vision blurred (see also section 4.4)                                                         |
|                                                 | Very rare | Cataract and glaucoma                                                                         |
| Cardiac disorders                               | Common    | Palpitations                                                                                  |
|                                                 | Uncommon  | Tachycardia                                                                                   |
|                                                 | Rare      | Cardiac arrhythmias, e.g. atrial fibrillation, supraventricular tachycardia, extrasystoles    |
|                                                 | Very rare | Angina pectoris. Prolongation of QTc-interval                                                 |
| Vascular disorders                              | Very rare | Variations in blood pressure                                                                  |
| Respiratory, thoracic and mediastinal disorders | Common    | Mild irritation in the throat, coughing, dysphonia including hoarseness                       |
|                                                 | Rare      | Bronchospasm                                                                                  |
| Gastrointestinal disorders                      | Uncommon  | Nausea                                                                                        |
| Skin and subcutaneous tissue disorders          | Uncommon  | Bruises                                                                                       |
| Musculoskeletal and connective tissue disorders | Uncommon  | Muscle cramps                                                                                 |

Candida infection in the oropharynx is due to drug deposition. Advising the patient to rinse the mouth out with water after each maintenance dose will minimise the risk. Oropharyngeal Candida infection usually responds to topical anti-fungal treatment without the need to discontinue the inhaled corticosteroid. If oropharyngeal thrush occurs, patients should also rinse their mouth out with water after the as-needed inhalations.

As with other inhalation therapy, paradoxical bronchospasm may occur very rarely, affecting less than 1 in 10,000 people, with an immediate increase in wheezing and shortness of breath after dosing. Paradoxical bronchospasm responds to a rapid-acting inhaled bronchodilator and should be treated straightaway. Gardette should be discontinued immediately, the patient should be assessed and an alternative therapy instituted if necessary (see section 4.4).

Systemic effects of inhaled corticosteroids may occur, particularly at high doses prescribed for prolonged periods. These effects are much less likely to occur than with oral corticosteroids. Possible systemic effects include Cushing's Syndrome, Cushingoid features, adrenal suppression, growth retardation in children and adolescents, decrease in bone mineral density, cataract and glaucoma. Increased susceptibility to infections and impairment of the ability to adapt to stress may also occur. Effects are probably dependent on dose, exposure time, concomitant and previous steroid exposure and individual sensitivity.

Treatment with  $\beta_2$  adrenoceptor agonists may result in an increase in blood levels of insulin, free fatty acids, glycerol and ketone bodies.

#### Paediatric population

It is recommended that the height of children receiving prolonged treatment with inhaled corticosteroids is regularly monitored (see section 4.4).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

### **4.9 Overdose**

An overdose of formoterol would likely lead to effects that are typical for  $\beta_2$  adrenoceptor agonists: tremor, headache, palpitations. Symptoms reported from isolated cases are tachycardia, hyperglycaemia, hypokalaemia, prolonged QTc-interval, arrhythmia, nausea and vomiting. Supportive and symptomatic treatment may be indicated. A dose of 90 micrograms administered during three hours in patients with acute bronchial obstruction raised no safety concerns.

Acute overdosage with budesonide, even in excessive doses, is not expected to be a clinical problem. When used chronically in excessive doses, systemic glucocorticosteroid effects, such as hypercorticism and adrenal suppression, may appear.

If Gardette therapy has to be withdrawn due to overdose of the formoterol component of the drug, provision of appropriate inhaled corticosteroid therapy must be considered.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs for obstructive airway diseases: Adrenergics, Inhalants.

ATC-code: R03AK07

#### Mechanisms of action and Pharmacodynamic effects

Gardette contains formoterol and budesonide, which have different modes of action and show additive effects in terms of reduction of asthma exacerbations. The specific properties of budesonide and formoterol allow the combination to be used either as maintenance and reliever therapy, or as maintenance treatment of asthma.

#### *Budesonide*

Budesonide is a glucocorticosteroid which when inhaled has a dose-dependent anti-inflammatory action in the airways, resulting in reduced symptoms and fewer asthma exacerbations. Inhaled budesonide has less severe adverse effects than systemic corticosteroids. The exact mechanism responsible for the anti-inflammatory effect of glucocorticosteroids is unknown.

#### *Formoterol*

Formoterol is a selective  $\beta_2$  adrenoceptor agonist that when inhaled results in rapid and long-acting relaxation of bronchial smooth muscle in patients with reversible airways obstruction. The bronchodilating effect is dose dependant, with an onset of effect within 1-3 minutes. The duration of effect is at least 12 hours after a single dose.

#### Clinical efficacy and safety

##### *Clinical efficacy for budesonide/formoterol maintenance therapy*

Clinical studies in adults have shown that the addition of formoterol to budesonide improved asthma symptoms and lung function, and reduced exacerbations. In two 12-week studies the effect on lung function of budesonide/formoterol was equal to that of the free combination of budesonide and formoterol, and exceeded that of budesonide alone. All treatment arms used a short-acting  $\beta_2$  adrenoceptor agonist as needed. There was no sign of attenuation of the anti-asthmatic effect over time.

Two 12-week paediatric studies have been performed in which 265 children aged 6-11 years were treated with a maintenance dose of budesonide/formoterol (2 inhalations of 80 micrograms/4.5 micrograms/inhalation twice daily), and a short-acting  $\beta_2$  adrenoceptor agonist as needed. In both studies, lung function was improved and the treatment was well tolerated compared to the corresponding dose of budesonide alone.

#### *Clinical efficacy for budesonide/formoterol maintenance and reliever therapy*

A total of 12076 asthma patients were included in 5 double-blind efficacy and safety studies (4447 were randomised to budesonide/formoterol maintenance and reliever therapy) for 6 or 12 months. Patients were required to be symptomatic despite use of inhaled glucocorticosteroids.

Budesonide/formoterol maintenance and reliever therapy provided statistically significant and clinically meaningful reductions in severe exacerbations for all comparisons in all 5 studies. This included a comparison with budesonide/formoterol at a higher maintenance dose with terbutaline as reliever (study 735) and budesonide/formoterol at the same maintenance dose with either formoterol or terbutaline as reliever (study 734) (Table 2). In study 735, lung function, symptom control, and reliever use were similar in all treatment groups. In study 734, symptoms and reliever use were reduced and lung function improved, compared with both comparator treatments. In the 5 studies combined, patients receiving budesonide/formoterol maintenance and reliever therapy used, on average, no reliever inhalations on 57% of treatment days. There was no sign of development of tolerance over time.

**Table 2 Overview of severe exacerbations in clinical studies**

| Study No.<br>Duration          | Treatment groups                                                       | n           | Severe exacerbations <sup>a</sup> |                         |
|--------------------------------|------------------------------------------------------------------------|-------------|-----------------------------------|-------------------------|
|                                |                                                                        |             | Events                            | Events/<br>patient-year |
| <b>Study 735<br/>6 months</b>  | <b>Budesonide/formoterol 160/4.5 µg bd + as needed</b>                 | <b>1103</b> | <b>125</b>                        | <b>0.23<sup>b</sup></b> |
|                                | Budesonide/formoterol 320/9 µg bd + terbutaline 0.4 mg as needed       | 1099        | 173                               | 0.32                    |
|                                | Salmeterol/fluticasone 2 x 25/125 µg bd + terbutaline 0.4 mg as needed | 1119        | 208                               | 0.38                    |
| <b>Study 734<br/>12 months</b> | <b>Budesonide/formoterol 160/4.5 µg bd + as needed</b>                 | <b>1107</b> | <b>194</b>                        | <b>0.19<sup>b</sup></b> |
|                                | Budesonide/formoterol 160/4.5 µg bd + formoterol 4.5 µg as needed      | 1137        | 296                               | 0.29                    |
|                                | Budesonide/formoterol 160/4.5 µg bd + terbutaline 0.4 mg as needed     | 1138        | 377                               | 0.37                    |

<sup>a</sup> Hospitalisation/emergency room treatment or treatment with oral steroids

<sup>b</sup> Reduction in exacerbation rate is statistically significant (P value <0.01) for both comparisons

Comparable efficacy and safety in adolescents and adults was demonstrated in 6 double-blind studies, comprising the 5 studies mentioned above and an additional study using a higher maintenance dose of 160/4.5 micrograms, two inhalations twice daily. These assessments were based on a total of 14385 asthma patients of whom 1847 were adolescents. The number of adolescent patients taking more than 8 inhalations on at least one day as part of budesonide/formoterol maintenance and reliever therapy was limited, and such use was infrequent.

In 2 other studies with patients seeking medical attention due to acute asthma symptoms, budesonide/formoterol provided rapid and effective relief of bronchoconstriction similar to salbutamol and formoterol.

## **5.2 Pharmacokinetic properties**

### Absorption

The fixed-dose combination of budesonide and formoterol, and the corresponding monoproducts have been shown to be bioequivalent with regard to systemic exposure of budesonide and formoterol, respectively. In spite of this, a small increase in cortisol suppression was seen after administration of the fixed-dose combination compared to the monoproducts. The difference is considered not to have an impact on clinical safety.

There was no evidence of pharmacokinetic interactions between budesonide and formoterol.

Pharmacokinetic parameters for the respective substances were comparable after the administration of budesonide and formoterol as monoproducts or as the fixed-dose combination. For budesonide, AUC was slightly higher, rate of absorption more rapid and maximal plasma concentration higher after administration of the fixed combination. For formoterol, maximal plasma concentration was similar after administration of the fixed combination. Inhaled budesonide is rapidly absorbed and the maximum plasma concentration is reached within 30 minutes after inhalation. In studies, mean lung deposition of budesonide after inhalation via the powder inhaler ranged from 32% to 44% of the delivered dose. The systemic bioavailability is approximately 49% of the delivered dose. In children 6-16 years of age the lung deposition falls in the same range as in adults for the same given dose. The resulting plasma concentrations were not determined.

Inhaled formoterol is rapidly absorbed and the maximum plasma concentration is reached within 10 minutes after inhalation. In studies the mean lung deposition of formoterol after inhalation via the powder inhaler ranged from 28% to 49% of the delivered dose. The systemic bioavailability is about 61% of the delivered dose.

#### Distribution and biotransformation

Plasma protein binding is approximately 50% for formoterol and 90% for budesonide. Volume of distribution is about 4 l/kg for formoterol and 3 l/kg for budesonide. Formoterol is inactivated via conjugation reactions (active O-demethylated and deformedylated metabolites are formed, but they are seen mainly as inactivated conjugates). Budesonide undergoes an extensive degree (approximately 90%) of biotransformation on first passage through the liver to metabolites of low glucocorticosteroid activity. The glucocorticosteroid activity of the major metabolites, 6-beta-hydroxy-budesonide and 16-alfa-hydroxy-prednisolone, is less than 1% of that of budesonide. There are no indications of any metabolic interactions or any displacement reactions between formoterol and budesonide.

#### Elimination

The major part of a dose of formoterol is transformed by liver metabolism followed by renal elimination. After inhalation, 8% to 13% of the delivered dose of formoterol is excreted unmetabolised in the urine. Formoterol has a high systemic clearance (approximately 1.4 l/min) and the terminal elimination half-life averages 17 hours.

Budesonide is eliminated via metabolism mainly catalysed by the enzyme CYP3A4. The metabolites of budesonide are eliminated in urine as such or in conjugated form. Only negligible amounts of unchanged budesonide have been detected in the urine. Budesonide has a high systemic clearance (approximately 1.2 l/min) and the plasma elimination half-life after i.v. dosing averages 4 hours.

The pharmacokinetics of formoterol in children have not been studied. The pharmacokinetics of budesonide and formoterol in patients with renal failure are unknown. The exposure of budesonide and formoterol may be increased in patients with liver disease.

#### Linearity/non-linearity

Systemic exposure for both budesonide and formoterol correlates in a linear fashion to administered dose.

### **5.3 Preclinical safety data**

The toxicity observed in animal studies with budesonide and formoterol, given in combination or separately, were effects associated with exaggerated pharmacological activity.

In animal reproduction studies, corticosteroids such as budesonide have been shown to induce malformations (cleft palate, skeletal malformations). However, these animal experimental results do not seem to be relevant in humans at the recommended doses. Animal reproduction studies with formoterol have shown a somewhat reduced fertility in male rats at high systemic exposure and implantation losses as well as decreased early postnatal survival and birth weight at considerably higher systemic exposures than those reached during clinical use. However, these animal experimental results do not seem to be relevant in humans.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate (which contains milk proteins).

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Gardette is an inspiratory flow driven, multidose powder inhaler. The inhaler is white with a red turning grip. The inhaler is made of different plastic materials (PP, PC, HDPE, LDPE, LLDPE, PBT). In each secondary package there are 1, 2, 3, 10 or 18 inhaler(s) containing 60 (or 120) doses.

Not all pack-sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements.

## **7. MARKETING AUTHORISATION HOLDER**

[To be completed nationally]

For Sweden (RMS):  
AstraZeneca AB  
151 85 Södertälje  
Sweden

## **8. MARKETING AUTHORISATION NUMBER(S)**

[To be completed nationally]

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

[To be completed nationally]

**10. DATE OF REVISION OF THE TEXT**

2020-12-17