

Package leaflet: Information for the user

Metadon Abcur 5 mg tablets

Metadon Abcur 10 mg tablets

methadone hydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Metadon Abcur is and what it is used for
2. What you need to know before you take Metadon Abcur
3. How to take Metadon Abcur
4. Possible side effects
5. How to store Metadon Abcur
6. Contents of the pack and other information

1. What Metadon Abcur is and what it is used for

Metadon Abcur is a morphine-like medicinal product which is used for:

- treating severe chronic pain which can be adequately managed only with opioid analgesics
- substitution therapy in opiate dependent patients, given together with medical and psychological treatment and social rehabilitation.

2. What you need to know before you take Metadon Abcur

Do not take Metadon Abcur:

- if you are allergic to methadone or any of the other ingredients of this medicine (listed in section 6).
- if you use or have used a monoamino-oxidase inhibitor (MAO-inhibitor) within the last two weeks (medicine used in the treatment of depression and Parkinson's disease).
- if you have respiratory depression, with slow and shallow breathing, or other acute breathing difficulties (acute obstructive airway disease).

Metadon Abcur must NOT be given to children.

Warnings and precautions

Talk to your doctor before taking Metadon Abcur if you:

- get acute asthma attacks
- suffer from lung disease or breathing difficulties
- have a heart disorder or ischemic heart disease
- have a liver disease or impaired liver function (including gallstones)
- have impaired kidney function (including kidney stones)
- have an underactive thyroid gland
- have skin- or tissue diseases
- have an enlarged prostate or narrowing of the urethra

- have been affected by a head injury and the pressure in the brain is higher than it should be (check this with your doctor). You may get severe headache.
- are treated for drug addiction or overdose of opioids
- have difficulties to pass water
- have stomach pains, diarrhoea or constipation
- have a low oxygen level or high carbon dioxide level in the blood
- take other medicines of opioid type (painkillers), such as morphine and pentazocine
- take other medicines which gets you relaxed and help you go to sleep (barbiturates and benzodiazepines), taking benzodiazepines with Metadon Abcur may result in difficulties in breathing (respiratory depression), coma and may be life-threatening. Even if benzodiazepines are prescribed, your doctor may need to change the dose, the duration of treatment or monitor you regularly.
- have low levels of potassium or magnesium in the blood

Metadon Abcur may affect the electrical signals which operate the heart constrictions, especially at high doses. Tell your doctor if you have a history of heart problems.

Talk to your doctor or pharmacist if you experience any of the following symptoms while taking Metadon Abcur:

- Weakness, fatigue, lack of appetite, nausea, vomiting or low blood pressure. This may be a symptom of the adrenals producing too little of the hormone cortisol, and you may need to take hormone supplement.
- Long-term use may cause decreased sex hormone levels and increased levels of the hormone prolactin. Contact your doctor if you experience symptoms such as decreased libido, impotence or absence of menstruation (amenorrhea).
- Contact your doctor if you experience severe upper abdominal pain possibly radiating to the back, nausea, vomiting or fever as this could be symptoms associated with inflammation of the pancreas (pancreatitis) or the biliary tract system.
- Pain or increased sensitivity to pain (hyperalgesia) which does not respond to a higher dosage of your medicine.

Tolerance, dependence, and addiction

This medicine contains methadone which is an opioid medicine. Repeated use of opioids can result in the drug being less effective (you become accustomed to it, known as tolerance). Repeated use of Metadon Abcur can also lead to dependence, abuse, and addiction, which may result in life-threatening overdose. The risk of these side effects can increase with a higher dose and longer duration of use.

Dependence or addiction can make you feel that you are no longer in control of how much medicine you need to take or how often you need to take it. When used for the treatment of pain, you might feel that you need to carry on taking your medicine, even when it doesn't help to relieve your pain.

The risk of becoming dependent or addicted varies from person to person. You may have a greater risk of becoming dependent or addicted on Metadon Abcur if:

- You or anyone in your family have ever abused or been dependent on alcohol, prescription medicines or illegal drugs ("addiction").
- You are a smoker.
- You have ever had problems with your mood (depression, anxiety, or a personality disorder) or have been treated by a psychiatrist for other mental illnesses.

If you notice any of the following signs whilst taking Metadon Abcur, it could be a sign that you have become dependent or addicted.

- You need to take the medicine for longer than advised by your doctor
- You need to take more than the recommended dose

- You are using the medicine for reasons other than prescribed, for instance, 'to stay calm' or 'help you sleep'
- You have made repeated, unsuccessful attempts to quit or control the use of the medicine
- When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again ('withdrawal effects')

If you notice any of these signs, speak to your doctor to discuss the best treatment pathway for you, including when it is appropriate to stop and how to stop safely (See section 3, If you stop taking Metadon Abcur).

Sleep-related breathing disorders

Metadon Abcur can cause sleep-related breathing disorders such as sleep apnoea (breathing pauses during sleep) and sleep related hypoxemia (low oxygen level in the blood). The symptoms can include breathing pauses during sleep, night awakening due to shortness of breath, difficulties to maintain sleep or excessive drowsiness during the day. If you or another person observe these symptoms, contact your doctor. A dose reduction may be considered by your doctor.

Elderly

Elderly patients have an increased risk of low blood pressure and fainting.

Store out of reach of children

Children are more sensitive to methadone than adults. Therefore, may poisoning occur at very low doses. In order to avoid children to take methadone by mistake when you use it at home, the methadone should be stored in a safe place, out of reach of children.

Other medicines and Metadon Abcur

Tell your doctor or pharmacist if you are taking or have recently taken or might take any other medicine.

Tell your doctor if you are taking any of the following medicines:

- medicines that affects your mental state (eg. thioridazine, perphenazine, risperidone, haloperidole and sertindole)
- medicines for the treatment of ADHD (atomoxetine)
- medicines used to treat depression, known as monoamine-oxidase inhibitor (MAO-inhibitor), especially if you have taken them within the last two weeks. Also tell your doctor if you are using other antidepressant medicines (eg. imipramine, nefazodone, fluoxetine, fluvoxamine, paroxetine and sertraline).
- other medicines used to treat depression known as tricyclic antidepressants (eg. clomipramine, nortriptyline, desipramine)
- medicines to treat heart diseases (verapamil, quinidine)
- antiarrhythmic agents (eg. propafenone and flecainide) and beta receptor blocker (metoprolol)
- medicines to treat breast cancer (tamoxifen)
- anti-inflammatory and immunosuppressive medicines (eg. dexamethasone and ciclosporin)
- antiviral medicines, including certain anti-HIV medicines eg. nevirapine, zidovudine, efavirenz, nelfinavir, delavirdin, ritonavir, combination of lopinavir/ritonavir, combination of ritonavir/saquinavir, didanosine and stavudine
- antibiotics of macrolide type (medicines to treat bacterial infections) such as clarithromycin, telithromycin and erythromycin
- rifampicin (a medicine to treat tuberculosis or TBC)
- antifungal medicines such as ketoconazole, itraconazole and fluconazole.
- cimetidine (medicine to treat gastric ulcer)
- naloxone (a medicine to treat breathing difficulties)
- medicines to treat drug addiction eg. naltrexone and buprenorphine
- medicines to treat epilepsy (eg. phenytoin, phenobarbital and carbamazepine)
- vitamin tablets containing C-vitamin
- medicines to treat diarrhoea (loperamide, diphenoxylate)
- medicines and products that make the urine more acidic such as ammonium chloride (sal ammoniac)
- diuretics medicines (spironolactone)

- medicines that make you sleepy or drowsy
- herbal medicines containing St John's wort
- Cannabidiol (a medicine used to treat seizures)
- Gabapentin and pregabalin (medicines used to treat epilepsy, nerve pain or anxiety), can increase the risk of opioid overdose, respiratory depression (breathing difficulties) and may be life-threatening.

Concomitant use of Methadone and sedative medicines such as benzodiazepines or related drugs increases the risk of drowsiness, difficulties in breathing (respiratory depression), coma and may be life-threatening. Because of this, concomitant use should only be considered when other treatment options are not possible. However, if your doctor does prescribe Metadon Abcur together with sedative medicines the dose and duration of concomitant treatment should be limited by your doctor.

The risk of side effects increases, if you use methadone concomitantly with antidepressants (such as citalopram, duloxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline, venlafaxine, amitriptyline, clomipramine, imipramine, nortriptyline). Contact your doctor if you experience symptoms such as:

- mental-status changes (e.g. agitation, hallucinations, coma)
- fast heartbeat, unstable blood pressure, fever
- exaggeration of reflexes, impaired coordination, muscle stiffness
- gastrointestinal symptoms (e.g. nausea, vomiting, diarrhoea)

Please tell your doctor about all sedative medicines you are taking and follow your doctor's dose recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.

Other medicines you may be taken can also affect the heart (eg. sotalol and amiodarone).

You must tell your doctor about all other medicines that you are taking, since they may potentially be dangerous if they are taken with methadone. In such situations, your doctor may decide that your heart should be monitored with ECG in the beginning of the treatment to check that no such effects occur.

Methadone may also affect certain blood- and urine samples. Tell your doctor that you are taking methadone before any test is performed.

Using Metadon Abcur with drink and alcohol

You are not allowed to drink alcohol while taking Metadon Abcur since this can give serious side effects. Do not drink grapefruit juice while being treated with Metadon Abcur, since this may change the effect of the medicine.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Some studies have reported birth defects or neurodevelopmental problems (problems with early childhood development) in children born to mothers who used methadone during pregnancy to treat opioid addiction. However, it is not possible to determine if this is caused by methadone use or other factors such as the health of the mother and social and environmental conditions associated with opioid addiction.

It is not recommended to use Metadon Abcur during labor.

Talk to your doctor if you are breastfeeding or thinking of breast-feeding while you are taking methadone as it may affect your baby. Monitor your baby for abnormal signs and symptoms such as increased drowsiness (more than usual), breathing difficulties or limpness. Consult your doctor immediately if you notice any of these symptoms.

Fertility:

Male impotence has been reported with long term use of methadone.

Driving and using machines

Methadone affects the coordination between the brain and body movements, so the ability to drive or use machines may be severely affected until your medication has been stabilised at a certain dose level. Therefore, you should not drive or use machines during the initial treatment period. How long it takes before you can drive or use machines is to a large extent individual, so you should decide this together with your doctor.

Metadon Abcur contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Metadon Abcur

Always take Metadon Abcur exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Before starting treatment and regularly during treatment, your doctor will discuss with you what you may expect from using Metadon Abcur, when and how long you need to take it, when to contact your doctor, and when you need to stop it (see also if you stop taking Metadon Abcur). It is important that you do not take more than the dose that you and your doctor have agreed upon. The individual need may be different. Modification or discontinuation of treatment should only be done in consultation with your doctor.

This product is for oral use only, and must not be injected.

Drug addiction

Adults:

The doctor adjusts the dosage according to your needs.

The usual initial dose is 10-30 mg. In patients with high opioid tolerance the initial dose is 25-40 mg. The dose is then increased gradually over a period of 3 weeks.

The usual maintenance dose is 60-120 mg daily but some patients need higher doses.

The maximum daily dose for treatment of drug addict should not exceed 150 mg.

Metadon Abcur is normally administered once daily.

Severe pain

Adults:

The usual initial dose is 5 mg, 1-3 times daily. Thereafter the dose can be increased slowly.

Patients who have taken opioids earlier may start the treatment at higher doses, 5-20 mg, 2-3 times daily, since their earlier experience of opioid treatment is taken into consideration. If required, other tablet strength for Metadon Abcur are available.

Continued determination of the daily dose, the dividing of doses and all dose adjustments during the treatment are done by the attending doctor and depends on the previous dose.

The maximum daily dose for treatment of severe pain should not exceed 100 mg.

Elderly:

It may be necessary to decrease the initial dose and you may need closer follow up when you are taking Metadon Abcur. Follow the doctor's advice.

Impaired liver or kidney function:

It may be necessary to decrease the dose or extend the dosage interval. Follow the doctor's advice.

If you take more Metadon Abcur than you should

If you have taken more medicinal product than you should or if for example a child has taken the medicine by mistake, you should immediately seek medical care.

Signs of overdose of Metadon Abcur may be:

- difficulty breathing or short respiratory arrest
- extreme sleepiness, fainting or coma
- pin point pupils (small pupils)
- muscle weakness
- cold, clammy and pale skin
- slow heartbeat, low blood pressure, heart attack or shock
- loss of coordination, difficulty walking, facial drooping, personality changes or trouble speaking (signs of a brain disorder known as toxic leukoencephalopathy)
- in severe cases death may occur
- it can result in low blood sugar

Contact your doctor if any of the above symptoms occur.

Children and elderly are more sensitive to this medicine.

If you forget to take Metadon Abcur

Do not take a double dose to make up for a forgotten tablet.

If you stop taking Metadon Abcur

You should only discontinue or terminate the treatment after talking with your doctor. You should not suddenly discontinue taking Metadon Abcur, since you may experience withdrawal symptoms such as sleeplessness, running nose, tearing, anorexia, diarrhoea and pains.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Metadon Abcur can cause side effects, although not everybody gets them.

Very common side effects (may affect more than 1 in 10 people):

- Nausea, vomiting

Common side effects (may affect up to 1 in 100 people):

- Weight increase
- Accumulation of water in the body
- Constipation
- Excitement (euphoria)
- You see or hear things that are not real (hallucinations)
- A feeling of dizziness or that you are spinning around
- Blurred vision
- Pinpoint pupils
- Drowsiness
- Rash
- Sweating
- Tiredness.

Uncommon side effects (may affect up to 1 in 100 people):

- Loss of appetite
- Breathing difficulties (with cough)
- Mouth dryness
- Inflammation in the tongue
- Depressed mood (dysphoria)

- Anxiety
- Sleep disturbance
- Confusion
- Decreased sexual drive
- Headache
- Fainting
- Itching, urticarial, rash
- Swollen legs
- Weakness
- Accumulation of water in the body
- Seizure in the bile ducts (gives stomach ache)
- Facial flush
- Low blood pressure
- Difficulties to pass water
- Difficulties in achieving or maintaining erection
- Menstrual disorders.

Rare side effects (may affect up to 1 in 1,000 people)

- Heart disorders, slower pulse, palpitation

Not known: frequency cannot be estimated from the available data

- Too low level of potassium or magnesium in the blood
- Decreased number of blood platelets in the blood.
- Changes in your mood
- You can become dependent on Metadon Abcur (for more information see section 2 Warnings and Precautions) Cardiac arrest
- Irregular heartbeats (cardiac arrhythmias), with symptoms such as too fast or too slow heartbeat, a fluttering in the chest, shortness of breath and chest pain.
- A fall in blood pressure on standing up which causes dizziness, light-headedness or fainting
- Sudden death
- Low blood sugar
- Sleep apnoea (breathing pauses during sleep)
- Symptoms associated with inflammation of the pancreas (pancreatitis) and the biliary tract system (a problem affecting a valve in the intestines known as sphincter of Oddi dysfunction), e.g. severe upper abdominal pain possibly radiating to the back, nausea, vomiting or fever.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#).

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Metadon Abcur

Keep this medicine out of the sight and reach of children. Store this medicine in a safe and secure storage space, where other people cannot access it. It can cause serious harm and be fatal to people when it has not been prescribed for them.

This medicinal product does not require any special storage conditions

Do not use this medicine after the expiry date which is stated on the package after EXP. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Contents of the pack and other information

What Metadon Abcur contains

- The active substance is methadone hydrochloride 5 mg or 10 mg.
- The other ingredients are lactose monohydrate, maize starch, magnesium stearate, talc, povidone (PVP K-25), colloidal anhydrous silica.

What Metadon Abcur looks like and contents of the pack

Metadon Abcur 5 mg is a white to off-white, round, flat bevelled tablet, concave with a score line on one side and embossing "M5" on the other side, tablet dimension 7 x 2.5 mm. The tablet can be divided into equal doses.

Metadon Abcur 10 mg is a white to off-white round, flat bevelled tablet, with score line on one side and embossing "M10" on the other side, tablet dimension 9 x 3.1 mm. The tablet can be divided into equal doses.

Metadon Abcur 5 mg and 10 mg is available in:

Child-resistant container: 25, 100, 200 tablets

Blister: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

[To be completed nationally]

This leaflet was last revised in

28 APRIL 2026