

Package leaflet: Information for the user

Mometason ABECE 50 micrograms/actuation, nasal spray, suspension

mometasone furoate

Please read this leaflet carefully before you start using this medicine because it contains important information for you.

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. (See section 4).
- You must talk to a doctor if you do not feel better or if you feel worse after 14 days.

What is in this leaflet:

1. What Mometason ABECE is and what it is used for
2. What you need to know before you use Mometason ABECE
3. How to use Mometason ABECE
4. Possible side effects
5. How to store Mometason ABECE
6. Contents of the pack and other information

1. What Mometason ABECE is and what it is used for

What is Mometason ABECE?

Mometason Abece contains mometasone furoate, one of group of medicines called corticosteroids. When mometasone furoate is sprayed into the nose, it can help to relieve inflammation (swelling and irritation of the nose), sneezing, itching and a blocked up or runny nose.

What is Mometason ABECE used for?

Hay fever and perennial rhinitis (allergies such as dust mites and animal hair)

Mometason Abece is used to treat the symptoms of hayfever (also called seasonal allergic rhinitis) and perennial rhinitis in adults 18 years and older.

Hay fever, which occurs at certain times of the year, is an allergic reaction caused by breathing in pollen from trees, grasses, weeds and also moulds and fungal spores. Perennial rhinitis occurs throughout the year and symptoms can be caused by a sensitivity to a variety of things including house dust mite, animal hair (or dander), feather and certain foods. Mometason Abece reduces the swelling and irritation in your nose and thereby relieving sneezing, itching and a blocked-up or runny nose caused by hay fever or perennial rhinitis.

2. What you need to know before you use Mometason ABECE

Do not use Mometason ABECE

- if you are allergic (hypersensitive) to mometasone furoate or any of the other ingredients of this medicine (listed in Section 6).
- if you have an untreated infection in your nose. Use of Mometason Abece during an untreated infection in your nose, such as herpes, can worsen the infection. You should then wait until the infection is resolved before you start using the nasal spray.
- if you have recently had an operation on your nose or you have injured your nose. You should wait until your nose has healed before you start using the nasal spray.

Warnings and precautions

Talk to your doctor or pharmacist before using Mometason Abece:

- if you have or have ever had tuberculosis.
- if you have any other type of infection.
- if you are taking other corticosteroid medicines, either by mouth or by injection.
- if you have cystic fibrosis.

While you are using Mometason Abece, talk to your doctor

- if your immune system is not functioning well (if you have difficulty in fighting infection) and you come into contact with anyone with measles or chickenpox. You should avoid coming into contact with anyone who has these infections.
- if you have an infection of the nose or throat.
- if you are using the medicine for several months or longer.
- if you have persistent irritation to the nose or throat.

Contact your doctor if you experience blurred vision or other visual disturbances.

When corticosteroid nasal sprays are used at high doses for long periods of time, side effects may occur due to the drug being absorbed in the body.

If your eyes are itching or irritated, your doctor may recommend that you use other treatments with Mometason Abece.

Other medicines and Mometason ABECE

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicines may increase the effects of Mometason Abece and your doctor may wish to monitor you carefully if you are taking these medicines (including some medicines for HIV: ritonavir, cobicistat).

If you are taking other corticosteroid medicines for allergy, either by mouth or injection, your doctor may advise you to stop taking them once you begin using Mometason Abece. A few people may find that once they discontinue oral or injected corticosteroids they suffer from some undesirable effects, such as joint or muscular pain, weakness and depression. You may also seem to develop other allergies, such as itchy, watering eyes or patches of red and itchy skin. If you develop any of these effects and you are worried, you should see your doctor.

Pregnancy and breast-feeding

There is little or no information on the use of Mometason Abece in pregnant women. It is not known if mometasone furoate is found in breast milk

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

Driving and using machines

There is no known information on the effect of Mometason Abece on the ability to drive or use machinery.

Mometason ABECE contains benzalkonium chloride

Mometason Abece contains 0.02 mg benzalkonium chloride per dose. Benzalkonium chloride may cause irritation or swelling inside the nose, especially if used for long time.

3. How to use Mometason ABECE

Dosage in adults aged 18 years and older is 2 sprays in each nostril once daily. When good effect has been achieved, the dose can be reduced to 1 spray in each nostril once daily. Do not exceed this dosage.

In some patients Mometason Abece nasal spray should begin to relieve symptoms 12 hours after the first dose. It is possible, however, that the full effect of the treatment is seen only after two days.

If you suffer badly from hay fever, the use of Mometason Abece may need to start some days before the pollen season begins.

You must talk to a doctor if you do not feel better or if you feel worse after 14 days.

Do not use Mometason Abece nasal spray for more than 3 months without consulting a doctor.

Please note that Mometason Abece should only be given to children at the order of a doctor.

Preparing your nasal spray for use

Your Mometason Abece nasal spray has a dust cap which protects the nozzle and keeps it clean.

Remember to take this off before using the spray and to replace it after use.

If you are using the spray for the first time you need to 'prime' the bottle by pumping the spray 10 times until a fine mist is produced:

1. Shake the bottle well.
2. Put your forefinger and middle finger either side of the nozzle and your thumb underneath the bottle (Figure 1).
3. Point the nozzle away from you and then press down with your fingers to pump the spray 10 times until a fine mist is produced.



Figure 1

If you have not used the spray for 14 days or more, you need to "re-prime" the bottle by pumping the spray 2 times until a fine mist is produced.

How to use your nasal spray

1. Shake the bottle well and remove the dust cap. (Figure 2)




Figure 2

2. Gently blow your nose.
3. Close one nostril and put the nozzle into the other nostril.
4. Tilt your head forward slightly, keeping the bottle upright.

5. Start to breathe in gently or slowly through your nose and whilst you are breathing in squirt a spray of fine mist into your nose by pressing down ONCE with your fingers (Figure 3).



Figure 3

6. Breathe out through your mouth. Repeat step 5 to inhale a second spray in the same nostril.
7. Remove the nozzle from this nostril and breathe out through the mouth.
8. Repeat steps 3 to 6 for the other nostril.

After using the spray, wipe the nozzle carefully with a clean handkerchief or tissue and replace the dust cap.

Cleaning your nasal spray

It is important to clean your nasal spray regularly for it to work properly. Remove the dust cap and gently pull off the nozzle. Wash the nozzle and dust cap in warm water and then rinse under a running tap. Allow the dust cap and nozzle to dry in a warm place. Push the nozzle back onto the bottle and replace the dust cap. The spray will need to be re-primed with 2 sprays when first used after cleaning.

If you use more Mometason ABECE than you should

Tell your doctor if you accidentally use more than you were told.

If you use steroids for a long time or in large amounts they may, rarely, affect some of your hormones.

If you forget to use Mometason ABECE

If you forget to use your nasal spray at the right time, use it as soon as you remember, then carry on as before. Do not take a double dose to make up for a forgotten dose.

If you stop using Mometason ABECE

In some patients Mometason Abece should begin to relieve symptoms 12 hours after the first dose; however, full benefit of treatment may not be seen for up to two days. It is very important that you use your nasal spray regularly.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Immediate hypersensitivity (allergic) reactions may occur after use of this product. These reactions may be severe. You should stop taking Mometason Abece and get immediate medical help if you experience symptoms such as:

- swollen face, tongue or pharynx
- trouble swallowing
- hives
- wheezing or trouble breathing.

When corticosteroid nasal sprays are used at high doses for long periods of time, side effects may occur due to the drug being absorbed in the body.

Other side effects

Most people do not have any problems after using the nasal spray. However, some people, after using Mometason Abeceor other corticosteroid nasal sprays, may find that they suffer from the following side effects:

Common side effects (may affect up to 1 in 10 people):

- headache
- sneezing
- nose bleeds
- sore nose or throat
- ulcers in the nose
- respiratory tract infection.

Not known (frequency cannot be estimated from the available data):

- increase in pressure in the eye (glaucoma) and/or cataracts causing visual disturbances,
- blurred vision
- damage to the partition in the nose which separates the nostrils
- changes in taste and smell
- difficulty in breathing and/or wheezing.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **the national reporting system** listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Mometason ABECE

- Keep this medicine out of the sight and reach of children.
- Do not use Mometason Abece after the expiry date which is stated on the bottle and carton. The expiry date refers to the last day of the month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Mometason ABECE contains

- The active substance is mometasone furoate. Each delivered spray (actuation) contains 50 micrograms of mometasone furoate (as mometasone furoate monohydrate).
- The other ingredients are microcrystalline cellulose and carmellose sodium, glycerol, citric acid monohydrate, sodium citrate, polysorbate 80, benzalkonium chloride, purified water.

What Mometason ABECE looks like and contents of the pack

Mometason Abece nasal spray is a nasal spray suspension packaged in a white plastic bottle with a spray pump..

- 1 bottle contains 10.0 g nasal spray, suspension, which is equivalent to 60 actuations.
- 2 bottles of 10.0 g nasal spray, suspension, which is equivalent to 120 actuations.
- 1 bottle contains 18.0 g nasal spray, suspension, which is equivalent to 140 actuations.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Evolan Pharma AB
Box 120
SE-182 12 Danderyd
Sweden

Manufacturer

FARMEA, 10, rue Bouché Thomas, Z.A.C d'Orgemont, 49000 ANGERS, France

FARMACLAIR, 440 avenue du Général de Gaulle, 14200 Hérouville Saint Clair, France

H2 Pharma, Zac La Croix Bonnet, 21 Rue Jacques Tati, Bois D Arcy, 78390 France

Perfect Care manufacturing S.R.L., 82 Complexului street, 085200, Mihailesti city, Giurgiu county, Romania

This medicinal product is authorised in the Member States of the EEA under the following names:

[To be completed nationally]

This leaflet was last revised in
2026-04-29