

Package leaflet: Information for the user

Pantoprazole Bluefish 40 mg gastro-resistant tablets

pantoprazole

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet, See section 4.

What is in this leaflet:

1. What Pantoprazole Bluefish is and what it is used for
2. What you need to know before you take Pantoprazole Bluefish
3. How to take Pantoprazole Bluefish
4. Possible side effects
5. How to store Pantoprazole Bluefish
6. Contents of the pack and other information

1. What Pantoprazole Bluefish is and what it is used for

Pantoprazole Bluefish contains the active substance pantoprazole. Pantoprazole Bluefish is a selective “proton pump inhibitor”, a medicine which reduces the amount of acid produced in your stomach. It is used for treating acid related diseases of the stomach and intestine.

Pantoprazole Bluefish is used to treat adults and adolescents 12 years of age and above for

- Reflux oesophagitis. An inflammation of your oesophagus (the tube which connects your throat to your stomach) accompanied by the regurgitation of stomach acid.

Pantoprazole Bluefish is used to treat adults for

- An infection with a bacterium called *Helicobacter pylori* in patients with duodenal ulcers and stomach ulcers in combination with two antibiotics (eradication therapy).

The aim is to get rid of the bacteria and so reduce the likelihood of these ulcers returning.

- Stomach and duodenal ulcers.
- Zollinger-Ellison-Syndrome and other conditions producing too much acid in the stomach.

2. What you need to know before you take Pantoprazole Bluefish

Do not take Pantoprazole Bluefish

- If you are allergic to pantoprazole or any of the other ingredients of this medicine (listed in section 6).
- If you are allergic to medicines containing other proton pump inhibitors

Warnings and precautions

Talk to your doctor or pharmacist or nurse before taking Pantoprazole Bluefish

- If you have severe liver problems. Please tell your doctor if you have ever had problems with your liver. He will check your liver enzymes more frequently, especially when you are taking

Pantoprazole Bluefish as a long-term treatment. In the case of a rise of liver enzymes the treatment should be stopped.

- If you have reduced body stores or risk factors for reduced vitamin B12 and receive long-term treatment with pantoprazole. As with all acid reducing agents, pantoprazole may lead to a reduced absorption of vitamin B12.
- If you are taking HIV protease inhibitors such as atazanavir (for the treatment of HIV-infection) at the same time as pantoprazole, ask your doctor for specific advice.
- Taking a proton pump inhibitor like pantoprazole, especially over a period of more than one year, may slightly increase your risk of fracture in the hip, wrist or spine. Tell your doctor if you have osteoporosis or if you are taking corticosteroids (which can increase the risk of osteoporosis).
- If you are on Pantoprazole Bluefish for more than three months it is possible that the levels of magnesium in your blood may fall. Low levels of magnesium can be seen as fatigue, involuntary muscle contractions, disorientation, convulsions, dizziness, increased heart rate. If you get any of these symptoms, please tell your doctor promptly. Low levels of magnesium can also lead to a reduction in potassium or calcium levels in the blood. Your doctor may decide to perform regular blood tests to monitor your levels of magnesium
- If you have ever had a skin reaction after treatment with a medicine similar to Pantoprazole Bluefish that reduces stomach acid.
- If you get a rash on your skin, especially in areas exposed to the sun tell your doctor as soon as you can, as you may need to stop your treatment with Pantoprazole Bluefish. Remember to also mention any other ill-effects like pain in your joints.
- If you are due to have a specific blood test (Chromogranin A).

Tell your doctor immediately, before or after taking this medicine, if you notice any of the following symptoms, which could be a sign of another, more serious, disease:

- An unintentional loss of weight
- Vomiting, particularly if repeated
- Vomiting blood; this may appear as dark coffee grounds in your vomit
- You notice blood in your stools; which may be black or tarry in appearance
- Difficulty in swallowing or pain when swallowing
- You look and feel pale (anaemia)
- Chest pain
- Stomach pain
- Severe and/or persistent diarrhoea, because this medicine has been associated with a small increase in infectious diarrhoea.

Your doctor may decide that you need some tests to rule out malignant disease because pantoprazole also alleviates the symptoms of cancer and could cause delay in diagnosing it. If your symptoms continue in spite of your treatment, further investigations will be considered.

If you take Pantoprazole Bluefish on a long-term basis (longer than 1 year) your doctor will probably keep you under regular surveillance. You should report any new and exceptional symptoms and circumstances whenever you see your doctor.

Children and adolescents

Pantoprazole Bluefish is not recommended for use in children as it has not been proven to work in children below 12 years of age

Other medicines and Pantoprazole Bluefish

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription

This is because Pantoprazole Bluefish may influence the effectiveness of other medicines, so tell your doctor if you are taking:

- Medicines such as ketoconazole, itraconazole and posaconazole (used to treat fungal infections) or erlotinib (used for certain types of cancer) because Pantoprazole Bluefish may stop these and other medicines from working properly.
- Warfarin and phenprocoumon, which affect the thickening, or thinning of the blood. You may need further checks.
- Medicines used to treat HIV-infection, such as atazanavir.
- Methotrexate (used to treat rheumatoid arthritis, psoriasis, and cancer) – if you are taking methotrexate your doctor may temporarily stop your Pantoprazole Bluefish treatment because pantoprazole can increase levels of methotrexate in the blood.
- Fluvoxamine (used to treat depression and other psychiatric diseases – if you are taking fluvoxamine your doctor may reduce the dose.
- Rifampicin (used to treat infections).
- St John's wort (*Hypericum perforatum*) (used to treat mild depression).

Pregnancy and breast-feeding

There are no adequate data from the use of pantoprazole in pregnant women. Excretion into human milk has been reported.

If you are pregnant or **breast-feeding**, or think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine .

You should use this medicine only if your doctor considers the benefit for you greater than the potential risk for your unborn child or baby.

Driving and using machines

Pantoprazole Bluefish has no or negligible influence on the ability to drive and use machines.

If you experience side effects like dizziness or disturbed vision, you should not drive or operate machines.

3. How to take Pantoprazole Bluefish

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Method of administration:

Take the tablets 1 hour before a meal without chewing or breaking them and swallow them whole with some water.

The recommended dose is:

Adults and adolescents 12 years of age and above

-To treat reflux oesophagitis

The usual dose is one tablet a day. Your doctor may tell you to increase to 2 tablets daily. The treatment period for reflux oesophagitis is usually between 4 and 8 weeks. Your doctor will tell you how long to take your medicine.

Adults

-For the treatment of an infection with a bacterium called *Helicobacter pylori* in patients with duodenal ulcers and stomach ulcers in combination with two antibiotics (Eradication therapy). One tablet, two times a day plus two antibiotic tablets of either amoxicillin, clarithromycin and metronidazole (or tinidazole), each to be taken two times a day with your pantoprazole tablet. Take the first pantoprazole tablet 1 hour before breakfast and the second pantoprazole tablet 1 hour before your evening meal. Follow your doctor's instructions and make sure you read the package leaflets for these antibiotics. The usual treatment period is one to two weeks.

-For the treatment of stomach and duodenal ulcers. The usual dose is one tablet a day. After consultation with your doctor, the dose may be doubled. Your doctor will tell you how long to take your medicine. The treatment period for stomach ulcers is usually between 4 and 8 weeks. The treatment period for duodenal ulcers is usually between 2 and 4 weeks.

-For the long-term treatment of Zollinger- Ellison-Syndrome and of other conditions in which too much stomach acid is produced. The recommended starting dose is usually two tablets a day. Take the two tablets 1 hour before a meal. Your doctor may later adjust the dose, depending on the amount of stomach acid you produce. If prescribed more than two tablets a day, the tablets should be taken twice daily.

If your doctor prescribes a daily dose of more than four tablets a day, you will be told exactly when to stop taking the medicine.

Patients with kidney problems

If you have kidney problems, you should not take Pantoprazole Bluefish for eradication of *Helicobacter pylori*.

Patients with liver problems

If you suffer from severe liver problems, you should not take more than one tablet 20 mg pantoprazole a day (for this purpose tablets containing 20 mg pantoprazole are available).

If you suffer from moderate or severe liver problems, you should not take Pantoprazole Bluefish for eradication of *Helicobacter pylori*.

Use in children and adolescents:

These tablets are not recommended for use in children below 12 years.

If you take more Pantoprazole Bluefish than you should

Tell your doctor or pharmacist. There are no known symptoms of overdose.

If you forget to take Pantoprazole Bluefish

Do not take a double dose to make up for a forgotten dose. Take your next normal dose at the usual time.

If you stop taking Pantoprazole Bluefish

Do not stop taking these tablets without first talking to your doctor or pharmacist.

If you have any further questions about the use of this medicine, ask your doctor or pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you get any of the following side effects, stop taking these tablets and tell your doctor immediately or contact the casualty department at your nearest hospital.

- **Serious allergic reactions (frequency rare:** may affect up to 1 in 1,000 people): swelling of the tongue and/or throat, difficulty in swallowing, hives (nettle rash), difficulties in breathing, allergic facial swelling (Quincke's oedema / angioedema), severe dizziness with very fast heartbeat and heavy sweating.
- **Serious skin conditions (frequency not known:** frequency cannot be estimated from the available data): blistering of the skin and rapid deterioration of your general condition, erosion (including slight bleeding) of eyes, nose, mouth/lips or genitals (Stevens-Johnson-Syndrome, Lyell-Syndrome, Erythema multiforme) and sensitivity to light.

Other serious conditions (frequency not known: frequency cannot be estimated from the available data): yellowing of the skin or whites of the eyes (severe damage to liver cells, jaundice) or fever, rash, and enlarged kidneys sometimes with painful urination and lower back pain (serious inflammation of the kidneys) possibly leading to kidney failure.

Other side effects are:

Common: (may affect up to 1 in 10 people):
Benign polyps in the stomach

Uncommon (may affect up to 1 in 100 people):
Headache; dizziness; diarrhoea; feeling sick, vomiting; bloating and flatulence (wind); constipation; dry mouth; abdominal pain and discomfort; skin rash, exanthema, eruption; itching; feeling weak, exhausted or generally unwell; sleep disorders; fracture of the hip, wrist or spine.

Rare (may affect up to 1 in 1,000 people):
Distortion or complete lack of the sense of taste; disturbances in vision such as blurred vision; hives; pain in the joints; muscle pains; weight changes; raised body temperature; high fever; swelling of the extremities (peripheral oedema); allergic reactions; depression; breast enlargement in males;

Very Rare (may affect up to 1 in 10,000 people):
Disorientation.

Not known: (frequency cannot be estimated from the available data):
Hallucination, confusion (especially in patients with a history of these symptoms); decreased sodium level in blood, decreased magnesium level in blood (see section 2), feeling of tingling, prickling, pins and needles, burning sensation or numbness, rash, possibly with pain in the joints.

Side effects identified through blood tests:

Uncommon (may affect up to 1 in 100 people); an increase in liver enzymes.

Rare (may affect up to 1 in 1,000 people); an increase in bilirubin; increased fat levels in the blood; sharp drop in circulating granular white blood cells, associated with high fever.

Very Rare (may affect up to 1 in 10,000 people);
a reduction in the number of blood platelets, which may cause you to bleed or bruise more than normal; a reduction in the number of white blood cells, which may lead to more frequent infections; coexisting abnormal reduction in the number of red and white blood cells, as well as platelets.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Pantoprazole Bluefish

Keep this medicine out of the sight and reach of children.

Store below 30°C.

Do not use this medicine after the expiry date which is stated on the pack after “EXP”. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Pantoprazole Bluefish contains

- The active substance is pantoprazole. Each gastro-resistant tablet contains 40 mg pantoprazole (as pantoprazole sodium sesquihydrate)
- The other ingredients are disodium phosphate anhydrous, mannitol, microcrystalline cellulose, croscarmellose sodium, magnesium stearate, hypromellose, triethyl citrate, sodium starch glycolate (Type A), methacrylic acid-ethyl acrylate copolymer (1:1) and yellow iron oxide (E 172).

What Pantoprazole Bluefish look like and contents of the pack

Pantoprazole Bluefish 40 mg gastro-resistant tablets are yellow, oval, biconvex and smooth tablets.

The tablet dimensions are the following:

- Width: 6.35 mm ± 0.32 mm (6.03 mm – 6.67 mm)
- Length: 12.00 mm ± 0.60 mm (11.40 mm – 12.60 mm)

Pack sizes:

Boxes of 7, 14, 28, 56 and 100 gastro-resistant tablets in blister packs.

Not all pack sizes may be marketed

Marketing Authorisation Holder

Bluefish Pharmaceuticals AB
P.O. Box 49013
100 28 Stockholm
Sweden

Manufacturer

Bluefish Pharmaceuticals AB
Gävlegatan 22
113 30 Stockholm
Sweden

Teva Pharma, S.L.U.,

C/C, no 4, Poligono Industrial Malpica
50016 Zaragoza,
Spain

This medicinal product is authorized in the Member States of the EEA under the following names:

Name of the Member State	Name of the medicinal product
Austria	Pantoprazole Bluefish 40 mg magensaftresistente tabletten
Denmark	Pantoprazole Bluefish 40mg
Ireland	Pantoprazole Bluefish 40 mg gastro-resistant tablet
Italy	TECNOZOL 40 mg gastroresistenti
Netherland	Pantoprazol Bluefish 40 mg maagsapresistente tabletten
Poland	Pantoprazole Bluefish 40mg enterotabletter
Portugal	Pantoprazole Bluefish
Spain	Pantoprazole Bluefish 40 mg comprimidos gastroresistentes
Sweden (RMS)	Pantoprazole Bluefish 40 mg enterotabletter

This leaflet was last revised in 2018-07-27