

Package leaflet: Information for the patient

Pirfenidon Medical Valley 267 mg film-coated tablets
Pirfenidon Medical Valley 534 mg film-coated tablets
Pirfenidon Medical Valley 801 mg film-coated tablets
pirfenidone

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Keep this leaflet. You may need to read it again.

- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What <invented name> is and what it is used for
2. What you need to know before you take <invented name>
3. How to take <invented name>
4. Possible side effects
5. How to store <invented name>
6. Contents of the pack and other information

1. What <invented name> is and what it is used for

<invented name> contains the active substance pirfenidone and it is used for the treatment of Idiopathic Pulmonary Fibrosis (IPF) in adults.

IPF is a condition in which the tissues in your lungs become swollen and scarred over time, and as a result makes it difficult to breathe deeply. This makes it hard for your lungs to work properly. <invented name> helps to reduce scarring and swelling in the lungs, and helps you breathe better.

2. What you need to know before you take <invented name>

Do not take <invented name>

- if you are allergic to pirfenidone or any of the other ingredients of this medicine (listed in section 6)
- if you have previously experienced angioedema with pirfenidone, including symptoms such as swelling of the face, lips and/or tongue which may be associated with difficulty breathing or wheezing
- if you are taking a medicine called fluvoxamine (used to treat depression and obsessive compulsive disorder [OCD])
- if you have severe or end stage liver disease
- if you have severe or end stage kidney disease requiring dialysis.

If any of the above affects you, do not take <invented name>. If you are unsure ask your doctor or pharmacist.

Warnings and precautions

Talk to your doctor or pharmacist before taking <invented name>

- You may become more sensitive to sunlight (photosensitivity reaction) when taking <invented name>. Avoid the sun (including sunlamps) whilst taking <invented name>. Wear sunblock daily and cover your arms, legs and head to reduce exposure to sunlight (see section 4: Possible side effects).
- You should not take other medicines, such as tetracycline antibiotics (such as doxycycline), which may make you more sensitive to sunlight.
- You should tell your doctor if you suffer from kidney problems.

- You should tell your doctor if you suffer from mild to moderate liver problems.
- You should stop smoking before and during treatment with <invented name>. Cigarette smoking can reduce the effect of <invented name>
- <invented name> may cause dizziness and tiredness. Be careful if you have to take part in activities where you have to be alert and co-ordinated.
- <invented name> can cause weight loss. Your doctor will monitor your weight whilst you are taking this medicine.
- Stevens-Johnson syndrome, toxic epidermal necrolysis, and drug reaction with eosinophilia and systemic symptoms (DRESS), have been reported in association with pifrenidone treatment. Stop using <invented name> and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

<invented name> may cause serious liver problems and some cases have been fatal. You will need a blood test before you start taking <invented name> and at monthly intervals for the first 6 months and then every 3 months thereafter whilst you are taking this medicine to check whether your liver is working properly. It is important that you have these regular blood tests for as long as you are taking <invented name>.

Children and adolescents

Do not give <invented name> to children and adolescents under the age of 18.

Other medicines and <invented name>

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines.

This is especially important if you are taking the following medicines, as they may change the effect of <invented name>.

Medicines that may increase side effects of <invented name>:

- enoxacin (a type of antibiotic)
- ciprofloxacin (a type of antibiotic)
- amiodarone (used to treat some types of heart disease)
- propafenone (used to treat some types of heart disease)
- fluvoxamine (used to treat depression and obsessive compulsive disorder (OCD)).

Medicines that may reduce how well <invented name> works:

- omeprazole (used in the treatment of conditions such as indigestion, gastroesophageal reflux disease)
- rifampicin (a type of antibiotic).

<invented name> with food and drink

Do not drink grapefruit juice whilst taking this medicine. Grapefruit may prevent <invented name> from working properly.

Pregnancy and breast-feeding

As a precautionary measure, it is preferable to avoid the use of <invented name> if you are pregnant, planning to become pregnant, or think you might be pregnant as the potential risks to the unborn child are unknown.

If you are breast-feeding or plan to breast-feed speak to your doctor or pharmacist before taking <invented name>. As it is unknown whether pifrenidone passes into breast milk, your doctor will discuss the risks and benefits of taking this medicine while breast-feeding if you decide to do so.

Driving and using machines

Do not drive or use machines if you feel dizzy or tired after taking <invented name>.

<invented name> contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take <invented name>

Treatment with <invented name> should be started and overseen by a specialist doctor experienced in the diagnosis and treatment of IPF.

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Your medicine will usually be given to you in increasing doses as follows:

- for the first 7 days take a dose of 267 mg (1 yellow tablet), 3 times a day with food (a total of 801 mg/day)
- from day 8 to 14 take a dose of 534 mg (2 yellow tablets or 1 orange tablet), 3 times a day with food (a total of 1,602 mg/day)
- from day 15 onwards (maintenance), take a dose of 801 mg (3 yellow tablets or 1 brown tablet), 3 times a day with food (a total of 2,403 mg/day).

The recommended maintenance daily dose of <invented name> is 801 mg (3 yellow tablets or 1 brown tablet) three times a day with food, for a total of 2403 mg/day.

Swallow the tablets whole with a drink of water, during or after a meal to reduce the risk of side effects such as nausea (feeling sick) and dizziness. If symptoms continue, see your doctor.

Dose reduction due to side effects

Your doctor may reduce your dose if you suffer from side effects such as, stomach problems, any skin reactions to sunlight or sun lamps, or significant changes to your liver enzymes.

If you take more <invented name> than you should

Contact your doctor, pharmacist or nearest hospital casualty department immediately if you have taken more tablets than you should and take your medicine with you.

If you forget to take <invented name>

If you forget a dose, take it as soon as you remember. Do not take a double dose to make up for a forgotten dose. Each dose should be separated by at least 3 hours. Do not take more tablets each day than your prescribed daily dose.

If you stop taking <invented name>

In some situations, your doctor may advise you to stop taking <invented name>. If for any reason you have to stop taking <invented name> for more than 14 consecutive days, your doctor will restart your treatment with a dose of 267 mg 3 times a day, gradually increasing this to a dose of 801 mg 3 times a day.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking <invented name> and seek medical attention immediately if you notice any of the following symptoms or signs

- Swelling of the face, lips and/or tongue, itching, hives, difficulty breathing or wheezing, or feeling faint, which are signs of angioedema, a serious allergic reaction or anaphylaxis.
- Yellowing of the eyes or skin, or dark urine, potentially accompanied by itching of the skin, pain on the upper right side of your stomach area (abdomen), loss of appetite, bleeding or bruising more easily than normal, or feeling tired. These may be signs of abnormal liver function and could indicate liver injury, which is an uncommon side effect of <invented name>.
- Reddish non-elevated, or circular patches on the trunk, often with central blisters, skin peeling,

ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (Stevens-Johnson syndrome or toxic epidermal necrolysis).

- Widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome or drug hypersensitivity syndrome).

Other side effects may include

Talk to your doctor if you get any side effects.

Very common side effects (may affect more than 1 in 10 people):

- infections of the throat or the airways going into the lungs and/or sinusitis
- feeling sick (nausea)
- stomach problems such as acid reflux, vomiting, and feeling constipated
- diarrhoea
- indigestion or stomach upset
- weight loss
- decreased appetite
- difficulty sleeping
- tiredness
- dizziness
- headache
- shortness of breath
- cough
- aching joints/joint pains.

Common side effects (may affect up to 1 in 10 people):

- bladder infections
- feeling sleepy
- changes in taste
- hot flushes
- stomach problems such as feeling bloated, abdominal pain and discomfort, heart burn and passing wind
- blood tests may show increased levels of liver enzymes
- skin reactions after going out in the sun or using sunlamps
- skin problems such as itchy skin, skin redness or red skin, dry skin, skin rash
- muscle pain
- feeling weak or feeling low in energy
- chest pain
- sunburn.

Uncommon side effects (may affect up to 1 in 100 people):

- Low levels of sodium in the blood. This may cause headache, dizziness, confusion, weakness, muscle cramps or nausea and vomiting.
- blood tests may show decrease in white blood cells.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

[To be completed nationally]

5. How to store <invented name>

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and the carton after EXP. The

expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What <invented name> contains

- The active substance is pirfenidone.
<invented name> 267 mg film-coated tablets: Each film-coated tablet contains 267 mg of pirfenidone.
<invented name> 534 mg film-coated tablets: Each film-coated tablet contains 534 mg of pirfenidone.
<invented name> 801 mg film-coated tablets: Each film-coated tablet contains 801 mg of pirfenidone.
- The other ingredients are:
Tablet core: Microcrystalline cellulose (E460), Povidone (E1201), Croscarmellose Sodium (E468), Silica colloidal anhydrous and Magnesium Stearate (E470b).
Film-coating: Polyvinyl alcohol (E1203), Titanium Dioxide (E171), Macrogol (E1521), Talc (E553b) and Iron oxide Yellow (E172) (<invented name> 267 mg and 534 mg film-coated tablets), Iron oxide red (E172) (<invented name> 534 mg and 801 mg film-coated tablets), Iron oxide black (E172) (<invented name> 801 mg film-coated tablets)

What <invented name> looks like and contents of the pack

<invented name> 267 mg film-coated tablets: yellow, oval-shaped, approximately 13.2 x 6.2 mm biconvex film coated tablets, plain on both sides.

<invented name> 534 mg film-coated tablets: orange, oval-shaped, approximately 16.3 x 8.5 mm biconvex film coated tablets, plain on both sides.

<invented name> 801 mg film-coated tablets: brown, oval-shaped, approximately 20.2 x 9.5 mm biconvex film coated tablets, plain on both sides.

Blisters composed by PVC/PCTFE–Alu

Pack sizes:

<invented name> 267 mg film-coated tablets are packaged in Blisters containing 21, 63, 252 film-coated tablets.

<invented name> 534 mg film-coated tablets are packaged in Blisters containing 21 film-coated tablets.

<invented name> 801 mg film-coated tablets are packaged in Blisters containing 84 tablets and 252 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer
[To be completed nationally]

This leaflet was last revised in 2024-06-20.