

Package leaflet: Information for the patient

Pramipexol Teva Pharma 0.26 mg prolonged-release tablets
Pramipexol Teva Pharma 0.52 mg prolonged-release tablets
Pramipexol Teva Pharma 1.05 mg prolonged-release tablets
Pramipexol Teva Pharma 1.57 mg prolonged-release tablets
Pramipexol Teva Pharma 2.1 mg prolonged-release tablets
Pramipexol Teva Pharma 2.62 mg prolonged-release tablets
Pramipexol Teva Pharma 3.15 mg prolonged-release tablets
pramipexole

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4

What is in this leaflet

1. What Pramipexol Teva Pharma is and what it is used for
2. What you need to know before you take Pramipexol Teva Pharma
3. How to take Pramipexol Teva Pharma
4. Possible side effects
5. How to store Pramipexol Teva Pharma
6. Contents of the pack and other information

1. What Pramipexol Teva Pharma is and what it is used for

Pramipexol Teva Pharma contains the active substance pramipexole and belongs to a group of medicines known as dopamine agonists, which stimulate dopamine receptors in the brain. Stimulation of the dopamine receptors triggers nerve impulses in the brain that help to control body movements.

Pramipexol Teva Pharma is used to treat the symptoms of primary Parkinson's disease in adults. It can be used alone or in combination with levodopa (another medicine for Parkinson's disease).

2. What you need to know before you take Pramipexol Teva Pharma

Do not take Pramipexol Teva Pharma

- if you are allergic to pramipexole or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor before taking Pramipexol Teva Pharma. Tell your doctor if you have (had) or develop any medical conditions or symptoms, especially any of the following:

- Kidney disease.
- Hallucinations (seeing, hearing or feeling things that are not there). Most hallucinations are visual.
- Dyskinesia (e.g. abnormal, uncontrolled movements of the limbs). If you have advanced Parkinson's disease and are also taking levodopa, you might develop dyskinesia during the up-titration of Pramipexol Teva Pharma.
- Dystonia (inability of keeping your body and neck straight and upright (axial dystonia)). In particular, you may experience forward flexion of the head and neck (also called antecollis), forward bending of the lower back (also called camptocormia) or sideways bending of the back

(also called pleurothotonus or Pisa Syndrome). If this happens, your doctor may want to change your medication.

- Sleepiness and episodes of suddenly falling asleep.
- Psychosis (e.g. comparable with symptoms of schizophrenia).
- Vision impairment. You should have regular eye examinations during treatment with Pramipexol Teva Pharma.
- Severe heart or blood vessels disease. You will need to have your blood pressure checked regularly, especially at the beginning of treatment. This is to avoid postural hypotension (a fall in blood pressure on standing up).

Tell your doctor if you or your family/carer notices that you are developing urges or cravings to behave in ways that are unusual for you and you cannot resist the impulse, drive or temptation to carry out certain activities that could harm yourself or others. These are called impulse control disorders and can include behaviours such as addictive gambling, excessive eating or spending, an abnormally high sex drive or preoccupation with an increase in sexual thoughts or feelings. Your doctor may need to adjust or stop your dose.

Tell your doctor if you or your family/carer notices that you are developing mania (agitation, feeling elated or over-excited) or delirium (decreased awareness, confusion, loss of reality). Your doctor may need to adjust or stop your dose.

Tell your doctor if you experience symptoms such as depression, apathy, anxiety, fatigue, sweating or pain after stopping or reducing your Pramipexol Teva Pharma treatment. If the problems persist more than a few weeks, your doctor may need to adjust your treatment.

Tell your doctor if you are developing an inability of keeping your body and neck straight and upright (axial dystonia). If this happens, your doctor may want to adjust or change your medication.

Pramipexol Teva Pharma is a specially designed tablet from which the active ingredient is gradually released, once the tablet has been ingested. Parts of tablets may occasionally be passed and seen in the stool (faeces) and may look like whole tablets. Inform your doctor if you find tablet pieces in your faeces.

Children and adolescents

Pramipexol Teva Pharma is not recommended for use in children or adolescents under 18 years.

Other medicines and Pramipexol Teva Pharma

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

You should avoid taking Pramipexol Teva Pharma together with antipsychotic medicines.

Take care if you are taking the following medicines:

- cimetidine (to treat excess stomach acid and stomach ulcers);
- amantadine (which can be used to treat Parkinson's disease);
- mexiletine (to treat irregular heartbeats, a condition known as ventricular arrhythmia);
- zidovudine (which can be used to treat the acquired immune deficiency syndrome (AIDS), a disease of the human immune system);
- cisplatin (to treat various types of cancers);
- quinine (which can be used for the prevention of painful night-time leg cramps and for the treatment of a type of malaria known as falciparum malaria (malignant malaria));
- procainamide (to treat irregular heart beat).

If you are taking levodopa, the dose of levodopa is recommended to be reduced when you start treatment with Pramipexol Teva Pharma.

Take care if you are using any medicines that calm you down (have a sedative effect) or if you are drinking alcohol. In these cases Pramipexol Teva Pharma may affect your ability to drive and operate machinery.

Pramipexol Teva Pharma with food, drink and alcohol

You should be cautious while drinking alcohol during treatment with Pramipexol Teva Pharma.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will then discuss with you if you should continue to take Pramipexol Teva Pharma.

The effect of Pramipexol Teva Pharma on the unborn child is not known. Therefore, do not take Pramipexol Teva Pharma if you are pregnant unless your doctor tells you to do so.

Pramipexol Teva Pharma should not be used during breast-feeding. Pramipexol Teva Pharma can reduce the production of breast milk. Also, it can pass into the breast milk and can reach your baby. If use of Pramipexol Teva Pharma is unavoidable, breast-feeding should be stopped.

Driving and using machines

Pramipexol Teva Pharma can cause hallucinations (seeing, hearing or feeling things that are not there). If affected, do not drive or use machines.


Pramipexol Teva Pharma has been associated with sleepiness and episodes of suddenly falling asleep, particularly in patients with Parkinson's disease. If you experience these side effects, you must not drive or operate machinery. You should tell your doctor if this occurs.

3. How to take Pramipexol Teva Pharma

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure. The doctor will advise you on the right dosing.

Take Pramipexol Teva Pharma prolonged-release tablets only once a day and each day at about the same time. You can take Pramipexol Teva Pharma with or without food. Swallow the tablets whole with water.

Do not chew, divide or crush the prolonged-release tablets. If you do, there is a danger you could overdose, because the medicine may be released into your body too quickly.



During the first week, the recommended daily dose is 0.26 mg pramipexole. The dose will be increased every 5-7 days as directed by your doctor until your symptoms are controlled (maintenance dose).

Ascending dose schedule of Pramipexol Teva Pharma prolonged-release tablets		
Week	Daily dose (mg)	Number of tablets
1	0.26	One Pramipexol Teva Pharma 0.26 mg prolonged-release tablet.
2	0.52	One Pramipexol Teva Pharma 0.52 mg prolonged-release tablet, OR two Pramipexol Teva Pharma 0.26 mg prolonged-release tablets.

3	1.05	One Pramipexol Teva Pharma 1.05 mg prolonged-release tablet, OR two Pramipexol Teva Pharma 0.52 mg prolonged-release tablets, OR four Pramipexol Teva Pharma 0.26 mg prolonged-release tablets.
---	------	---

The recommended maintenance dose is 1.05 mg per day. However, your dose may have to be increased even further. If necessary, your doctor may increase your dose up to a maximum of 3.15 mg of pramipexole a day. A lower maintenance dose of one Pramipexol Teva Pharma 0.26 mg prolonged-release tablet a day is also possible.

Patients with kidney disease

If you have kidney disease, your doctor may advise you to take the usual starting dose of 0.26 mg prolonged-release tablets only every other day for the first week. After that, your doctor may increase the dosing frequency to one 0.26 mg prolonged-release tablet every day. If a further dose increase is necessary, your doctor may adjust it in steps of 0.26 mg pramipexole.

If you have serious kidney problems, your doctor may need to switch you to a different pramipexole medicine. If during treatment your kidney problems get worse, you should contact your doctor as soon as possible.

If you are switching from pramipexole (immediate release) tablets

Your doctor will base your dose of pramipexole prolonged-release tablets on the dose of pramipexole (immediate release) tablets you were taking.

Take your pramipexole (immediate release) tablets as normal the day before you switch. Then take your pramipexole prolonged-release tablets next morning and do not take any more pramipexole (immediate release) tablets.

If you take more Pramipexol Teva Pharma than you should

If you accidentally take too many tablets,

- Contact your doctor or nearest hospital casualty department immediately for advice.
- You may experience vomiting, restlessness, or any of the side effects as described in section 4 "Possible side effects".

If you forget to take Pramipexol Teva Pharma

If you forget to take a dose of Pramipexol Teva Pharma, but remember within 12 hours of your usual time, take your tablet straightaway and then take your next tablet at the usual time.

If you forget for more than 12 hours, simply take the next single dose at the usual time. Do not take a double dose to make up for a forgotten tablet dose.

If you stop taking Pramipexol Teva Pharma

Do not stop taking Pramipexol Teva Pharma without first talking to your doctor. If you have to stop taking this medicine, your doctor will reduce the dose gradually. This reduces the risk of worsening symptoms.

If you suffer from Parkinson's disease you should not stop treatment with Pramipexol Teva Pharma abruptly. A sudden stop could cause you to develop a medical condition called neuroleptic malignant syndrome which may represent a major health risk. The symptoms include:

- akinesia (loss of muscle movement),
- rigid muscles,
- fever,
- unstable blood pressure,
- tachycardia (increased heart rate),
- confusion,
- depressed level of consciousness (e.g. coma).

If you stop or reduce Pramipexol Teva Pharma you may also develop a medical condition called dopamine agonist withdrawal syndrome. The symptoms include depression, apathy, anxiety, fatigue, sweating or pain. **If you experience these symptoms you should contact your physician.**

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

You may experience the following side effects:

Very common: may affect more than 1 in 10 people

- Dyskinesia (e.g. abnormal, uncontrolled movements of the limbs)
- Sleepiness
- Dizziness
- Nausea (sickness)

Common: may affect up to 1 in 10 people

- Urge to behave in an unusual way
- Hallucinations (seeing, hearing or feeling things that are not there)
- Confusion
- Tiredness (fatigue)
- Sleeplessness (insomnia)
- Excess of fluid, usually in the legs (peripheral oedema)
- Headache
- Hypotension (low blood pressure)
- Abnormal dreams
- Constipation
- Visual impairment
- Vomiting (being sick)
- Weight loss including decreased appetite

Uncommon: may affect up to 1 in 100 people

- Paranoia (e.g. excessive fear for one's own well-being)
- Delusion
- Excessive daytime sleepiness and suddenly falling asleep
- Amnesia (memory disturbance)
- Hyperkinesia (increased movements and inability to keep still)
- Weight increase
- Allergic reactions (e.g. rash, itching, hypersensitivity)
- Fainting
- Cardiac failure (heart problems which can cause shortness of breath or ankle swelling)*
- Inappropriate antidiuretic hormone secretion*
- Restlessness
- Dyspnoea (difficulties to breathe)
- Hiccups
- Pneumonia (infection of the lungs)
- Inability to resist the impulse, drive or temptation to perform an action that could be harmful to you or others, which may include:
 - Strong impulse to gamble excessively despite serious personal or family consequences.
 - Altered or increased sexual interest and behaviour of significant concern to you or to others, for example, an increased sexual drive.
 - Uncontrollable excessive shopping or spending
 - Binge eating (eating large amounts of food in a short time period) or compulsive eating (eating more food than normal and more than is needed to satisfy your hunger)*

- Delirium (decreased awareness, confusion, loss of reality)

Rare: may affect up to 1 in 1,000 people

- Mania (agitation, feeling elated or over-excited)
- Spontaneous penile erection

Not known: frequency cannot be estimated from the available data

- After stopping or reducing your Pramipexol Teva Pharma treatment: Depression, apathy, anxiety, fatigue, sweating or pain may occur (called dopamine agonist withdrawal syndrome or DAWS).

Tell your doctor if you experience any of these behaviours; he will discuss ways of managing or reducing the symptoms.

For the side effects marked with * a precise frequency estimation is not possible, since these side effects were not observed in clinical studies among 2,762 patients treated with pramipexole. The frequency category is probably not greater than “uncommon”.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Pramipexol Teva Pharma

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Pramipexol Teva Pharma prolonged-release tablets contains

The active substance is pramipexole.

Each tablet contains 0.26 mg, 0.52 mg, 1.05 mg, 1.57 mg, 2.1 mg, 2.62 mg, or 3.15 mg pramipexole as 0.375 mg, 0.75 mg, 1.5 mg, 2.25 mg, 3 mg, 3.75 mg, or 4.5 mg pramipexole dihydrochloride monohydrate, respectively.

The other excipients are hypromellose, calcium hydrogen phosphate, magnesium stearate and colloidal anhydrous silica.

What Pramipexol Teva Pharma prolonged-release tablets looks like and contents of the pack

Pramipexol Teva Pharma 0.26 mg prolonged-release tablets: The round tablets of 9 mm diameter are white or nearly white, have a flat surface with beveled edges and are marked with 026 on one side

Pramipexol Teva Pharma 0.52 mg prolonged-release tablets: The round tablets of 10 mm diameter are white or nearly white, biconvex and are marked with 052 on one side

Pramipexol Teva Pharma 1.05 mg prolonged-release tablets: The round tablets of 10 mm diameter are white or nearly white, biconvex, and are marked with 105 on one side

Pramipexol Teva Pharma 1.57 mg prolonged-release tablets: The round tablets of 10 mm diameter are white or nearly white, biconvex and are marked with 157 on one side

Pramipexol Teva Pharma 2.1 mg prolonged-release tablets: The round tablets of 10 mm diameter are white or nearly white, biconvex and are marked with 210 on one side

Pramipexol Teva Pharma 2.62 mg prolonged-release tablets: The round tablets of 10 mm diameter are white or nearly white, biconvex and are marked with 262 on one side

Pramipexol Teva Pharma 3.15 mg prolonged-release tablets: The round tablets of 11 mm diameter are white or nearly white, have a flat surface with beveled edges and are marked with 315 on one side

Pramipexol Teva Pharma is available in packs containing 10, 30 and 100 prolonged-release tablets in aluminium blisters.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

[To be completed nationally]

This medicine is authorised in the Member States of the European Economic Area and in the United Kingdom (Northern Ireland) under the following names:

[To be completed nationally]

This leaflet was last revised in <{MM/YYYY}> <{month YYYY}>.

[To be completed nationally]24 October 2024