

## Package leaflet: Information for the user

|                                                     |
|-----------------------------------------------------|
| <b>Rosuvastatin STADA 5 mg film-coated tablets</b>  |
| <b>Rosuvastatin STADA 10 mg film-coated tablets</b> |
| <b>Rosuvastatin STADA 20 mg film-coated tablets</b> |
| <b>Rosuvastatin STADA 40 mg film-coated tablets</b> |

rosuvastatin

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

1. What Rosuvastatin STADA is and what it is used for
2. What you need to know before you take Rosuvastatin STADA
3. How to take Rosuvastatin STADA
4. Possible side effects
5. How to store Rosuvastatin STADA
6. Contents of the pack and other information

### **1. What Rosuvastatin STADA is and what it is used for**

Rosuvastatin STADA contains the active ingredient rosuvastatin, which belongs to a group of medicines called statins.

#### **You have been prescribed Rosuvastatin STADA because:**

- You have a high cholesterol level. This means you are at risk from a heart attack or stroke. Rosuvastatin STADA is used in adults, adolescents and children 6 years or older to treat high cholesterol.

You have been advised to take a statin, because changing your diet and doing more exercise were not enough to correct your cholesterol levels. You should continue with your cholesterol-lowering diet and exercise while you are taking Rosuvastatin STADA.

Or

- You have other factors that increase your risk of having a heart attack, stroke or related health problems.

Heart attack, stroke and other problems can be caused by a disease called atherosclerosis. Atherosclerosis is due to build-up of fatty deposits in your arteries.

### **Why it is important to keep taking Rosuvastatin STADA**

Rosuvastatin STADA is used to correct the levels of fatty substances in the blood called lipids, the most common of which is cholesterol.

There are different types of cholesterol found in the blood - 'bad' cholesterol (LDL-C) and 'good' cholesterol (HDL-C).

- Rosuvastatin STADA can reduce the 'bad' cholesterol and increase the 'good' cholesterol.
- It works by helping to block your body's production of 'bad' cholesterol. It also improves your body's ability to remove it from your blood.

For most people, high cholesterol does not affect the way they feel because it does not produce any symptoms. However, if it is left untreated, fatty deposits can build up in the walls of your blood vessels causing them to narrow.

Sometimes, these narrowed blood vessels can get blocked which can cut off the blood supply to the heart or brain leading to a heart attack or a stroke. By lowering your cholesterol levels, you can reduce your risk of having a heart attack, a stroke or related health problems.

You need to **keep taking Rosuvastatin STADA**, even if it has got your cholesterol to the right level, because **it prevents your cholesterol levels from creeping up again** and causing build-up of fatty deposits. However, you should stop if your doctor tells you to do so, or you have become pregnant.

## 2. What you need to know before you take Rosuvastatin STADA

### DO NOT take Rosuvastatin STADA

- **If you have ever had an allergic reaction to rosuvastatin** or to any of the other ingredients of this medicine (listed in section 6).
  - **If you are pregnant** or breast-feeding. If you become pregnant while taking Rosuvastatin STADA **stop taking it immediately and tell your doctor**. Women should avoid becoming pregnant while taking Rosuvastatin STADA by using suitable contraception.
  - **If you have liver disease.**
  - **If you have severe kidney problems.**
  - **If you have repeated or unexplained muscle aches or pains.**
  - **If you take a drug combination of sofosbuvir/velpatasvir/voxilaprevir** (used for viral infection of the liver called hepatitis C).
  - **If you take a drug called ciclosporin** (used, for example, after organ transplants)
- If any of the above applies to you (or you are in doubt), please go back and see your doctor.

### In addition, do not take 40 mg of rosuvastatin (the highest dose):

- **If you have moderate kidney problems** (if in doubt, please ask your doctor).
- **If your thyroid gland** is not working properly.
- **If you have had any repeated or unexplained muscle aches or pains**, a personal or family history of muscle problems, or a previous history of muscle problems when taking other cholesterol-lowering medicines.
- **If you regularly drink large amounts of alcohol.**
- **If you are of Asian origin** (Japanese, Chinese, Filipino, Vietnamese, Korean and Indian).
- **If you take other medicines called fibrates** to lower your cholesterol.

If any of the above applies to you (or you are in doubt), **please go back and see your doctor**.

### Warnings and precautions

Talk to your doctor or pharmacist before taking this medicine.

- **If you have problems with your kidneys.**
- **If you have problems with your liver.**
- **If you have had repeated or unexplained muscle aches or pains**, a personal or family history of muscle problems, or a previous history of muscle problems when taking other cholesterol-lowering medicines. Tell your doctor immediately if you have unexplained muscle aches or pains especially if you feel unwell or have a fever. Also tell your doctor or pharmacist if you have a muscle weakness that is constant.

- If you have or have had **myasthenia** (a disease with general muscle weakness including in some cases muscles used when breathing), or **ocular myasthenia** (a disease causing eye muscle weakness) as statins may sometimes aggravate the condition or lead to the occurrence of myasthenia (see section 4).
- **If you have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after taking Rosuvastatin STADA or other related medicines.**
- **If you regularly drink large amounts of alcohol.**
- **If your thyroid gland** is not working properly.
- **If you take other medicines called fibrates** to lower your cholesterol. Please read this leaflet carefully, even if you have taken other medicines for high cholesterol before.
- **If you take medicines used to treat the HIV infection** e.g. ritonavir with lopinavir and/or atazanavir, please see "Other medicines and Rosuvastatin STADA".
- If you are taking or have taken in the last 7 days a medicine called fusidic acid (a medicine for bacterial infection), orally or by injection. The combination of fusidic acid and Rosuvastatin STADA can lead to serious muscle problems (rhabdomyolysis), please see "Other medicines and Rosuvastatin STADA".
- **If you are over 70** (as your doctor needs to choose the right start dose of Rosuvastatin STADA to suit you).
- **If you have severe respiratory failure.**
- **If you are of Asian origin** - that is Japanese, Chinese, Filipino, Vietnamese, Korean and Indian. Your doctor needs to choose the right start dose of Rosuvastatin STADA to suit you.

If any of the above applies to you (or if you are not sure):

- **Do not take 40 mg of rosuvastatin (the highest dose) and check with your doctor or pharmacist before you actually start taking any dose of Rosuvastatin STADA.**

Serious skin reactions including Stevens-Johnson syndrome and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in association with rosuvastatin treatment. Stop using Rosuvastatin STADA and seek medical attention immediately if you notice any of the symptoms described in section 4.

In a small number of people, statins can affect the liver. This is identified by a simple test which looks for increased levels of liver enzymes in the blood. For this reason, your doctor will usually carry out this blood test (liver function test) before and during treatment with Rosuvastatin STADA.

While you are on this medicine your doctor will monitor you closely if you have diabetes or are at risk of developing diabetes. You are likely to be at risk of developing diabetes if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure.

### **Children and adolescents**

- **If the patient is under 6 years old:** Rosuvastatin STADA should not be given to children younger than 6 years.
- **If the patient is below 18 years of age:** The Rosuvastatin STADA 40 mg tablet is not suitable for use in children and adolescents below 18 years of age.

### **Other medicines and Rosuvastatin STADA**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Tell your doctor if you are taking any of the following:

- ciclosporin (used for example, after organ transplants)
- warfarin, clopidogrel or ticagrelor (or any other drug used for thinning the blood)
- fibrates (such as gemfibrozil, fenofibrate) or any other medicine used to lower cholesterol (such as ezetimibe)

- indigestion remedies (used to neutralise acid in your stomach)
- erythromycin (an antibiotic), fusidic acid (an antibiotic – please see below and “Warnings and precautions”)
- an oral contraceptive (the pill), regorafenib (used to treat cancer)
- darolutamide (used to treat cancer)
- capmatinib (used to treat cancer)
- hormone replacement therapy
- fostamatinib (used to treat low platelet counts)
- febuxostat (used to treat and prevent high blood levels of uric acid)
- teriflunomide (used to treat multiple sclerosis)
- any of the following drugs used to treat viral infections, including HIV or hepatitis C infection, alone or in combination (please see Warnings and precautions): ritonavir, lopinavir, atazanavir, sofosbuvir, voxilaprevir, ombitasvir, paritaprevir, dasabuvir, velpatasvir, grazoprevir, elbasvir, glecaprevir, pibrentasvir.

The effects of these medicines could be changed by Rosuvastatin STADA or they could change the effect of Rosuvastatin STADA.

If you need to take oral fusidic acid to treat a bacterial infection you will need to temporarily stop using this medicine. Your doctor will tell you when it is safe to restart Rosuvastatin STADA. Taking Rosuvastatin STADA with fusidic acid may rarely lead to muscle weakness, tenderness or pain (rhabdomyolysis). See more information regarding rhabdomyolysis in section 4.

### **Pregnancy and breast-feeding**

**Do not take Rosuvastatin STADA** if you are pregnant or breast-feeding. If you become pregnant while taking Rosuvastatin STADA **stop taking it immediately** and tell your doctor. Women should avoid becoming pregnant while taking Rosuvastatin STADA by using suitable contraception.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

Most people can drive a car and operate machinery while using Rosuvastatin STADA - it will not affect their ability. However, some people feel dizzy during treatment with Rosuvastatin STADA. If you feel dizzy, consult your doctor before attempting to drive or use machines.

### **Rosuvastatin STADA contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars (lactose or milk sugar), contact your doctor before taking this medicinal product.

For the full list of ingredients please see “Contents of the pack and other information”.

## **3. How to take Rosuvastatin STADA**

Always take this medicine as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

### **Usual doses in adults**

**If you are taking Rosuvastatin STADA for high cholesterol:**

#### **Starting dose**

Your treatment with Rosuvastatin STADA must start with the **5 mg or the 10 mg dose**, even if you have taken a higher dose of a different statin before. The choice of your start dose will depend upon:

- Your cholesterol levels.
- The level of risk you have of experiencing a heart attack or stroke.
- Whether you have a factor that may make you more sensitive to possible side effects.

Please check with your doctor or pharmacist which start dose of Rosuvastatin STADA will best suit you.

Your doctor may decide to give you the lowest dose (5 mg) if:

- You are of **Asian origin** (Japanese, Chinese, Filipino, Vietnamese, Korean and Indian).
- You are **over 70 years** of age.
- You have moderate kidney problems.
- You are at risk of muscle aches and pains (myopathy).

### **Increasing the dose and maximum daily dose**

Your doctor may decide to increase your dose. This is so that you are taking the amount of Rosuvastatin STADA that is right for you. If you started with a 5 mg dose, your doctor may decide to double this to 10 mg, then 20 mg and then 40 mg if necessary. If you started on 10 mg, your doctor may decide to double this to 20 mg and then 40 mg if necessary. There will be a gap of four weeks between every dose adjustment.

The maximum daily dose of Rosuvastatin STADA is 40 mg. It is only for patients with high cholesterol levels and a high risk of heart attacks or stroke whose cholesterol levels are not lowered enough with 20 mg.

### **If you are taking Rosuvastatin STADA to reduce your risk of having a heart attack, stroke or related health problems:**

The recommended dose is 20 mg daily. However, your doctor may decide to use a lower dose if you have any of the factors mentioned above.

### **Use in children and adolescents aged 6-17 years**

The dose range in children and adolescents aged 6 to 17 years is 5 to 20 mg once daily. The usual start dose is 5 mg per day and your doctor may gradually increase your dose to find the right amount of Rosuvastatin STADA for you. The maximum daily dose of Rosuvastatin STADA is 10 or 20 mg for children aged 6 to 17 years depending on your underlying condition being treated. Take your dose once a day. Rosuvastatin STADA **40 mg** tablet should **not** be used by children.

### **Taking your tablets**

Swallow each tablet whole with a drink of water.

**Take Rosuvastatin STADA once daily.** You can take it at any time of the day with or without food.

Try to take your tablet at the same time every day to help you to remember it.

### **Regular cholesterol checks**

It is important to go back to your doctor for regular cholesterol checks, to make sure your cholesterol has reached and is staying at the correct level.

Your doctor may decide to increase your dose so that you are taking the amount of Rosuvastatin STADA that is right for you.

### **If you take more Rosuvastatin STADA than you should**

Contact your doctor or nearest hospital for advice.

If you go into hospital or receive treatment for another condition, tell the medical staff that you're taking Rosuvastatin STADA.

### **If you forget to take Rosuvastatin STADA**

Don't worry, just take your next scheduled dose at the correct time. Do not take a double dose to make up for a forgotten dose.

### **If you stop taking Rosuvastatin STADA**

Talk to your doctor if you want to stop taking Rosuvastatin STADA. Your cholesterol levels

might increase again if you stop taking Rosuvastatin STADA.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

It is important that you are aware of what these side effects may be. They are usually mild and disappear after a short time.

**Stop taking Rosuvastatin STADA and seek medical help immediately** if you have any of the following symptoms:

- Difficulty in breathing, with or without swelling of the face, lips, tongue and/or throat.
- Swelling of the face, lips, tongue and/or throat, which may cause difficulty in swallowing.
- Severe itching of the skin (with raised lumps).
- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (Stevens-Johnson syndrome)
- Widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome or drug hypersensitivity syndrome).

**Also, stop taking Rosuvastatin STADA and talk to your doctor immediately**

- **if you have any unusual aches or pains** in your muscles which go on for longer than you might expect. Muscle symptoms are more common in children and adolescents than in adults. As with other statins, a very small number of people have experienced unpleasant muscle effects and rarely these have gone on to become a potentially life threatening muscle damage known as *rhabdomyolysis*.
- **if you experience muscle rupture**
- **if you have lupus-like disease syndrome** (including rash, joint disorders and effects on blood cells)

**Common possible side effects (may affect up to 1 in 10 people):**

- Headache, stomach pain, constipation, feeling sick, muscle pain, feeling weak, dizziness
- An increase in the amount of protein in the urine - this usually returns to normal on its own without having to stop taking your Rosuvastatin STADA tablets (only for 40 mg of rosuvastatin)
- Diabetes. This is more likely if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure. Your doctor will monitor you while you are taking this medicine

**Uncommon possible side effects (may affect up to 1 in 100 people):**

- Rash, itching or other skin reactions
- An increase in the amount of protein in the urine - this usually returns to normal on its own without having to stop taking your Rosuvastatin STADA tablets (only for 5 mg, 10 mg and 20 mg of rosuvastatin)

**Rare possible side effects (may affect up to 1 in 1 000 people):**

- Severe allergic reaction - signs include swelling of the face, lips, tongue and/or throat, difficulty in swallowing and breathing, a severe itching of the skin (with raised lumps). If you think you are having an allergic reaction, then stop taking Rosuvastatin STADA and seek medical help immediately
- Muscle damage in adults - as a precaution, stop taking Rosuvastatin STADA and talk to your doctor immediately if you have any unusual aches or pains in your muscles which go

- on for longer than expected
- A severe stomach pain (inflamed pancreas)
- Increase in liver enzymes in the blood
- Bleeding or bruising more easily than normal due to low level of blood platelets
- Lupus-like disease syndrome (including rash, joint disorders and effects on blood cells).

**Very rare possible side effects (may affect up to 1 in 10 000 people):**

- Jaundice (yellowing of the skin and eyes), hepatitis (an inflamed liver), traces of blood in your urine, damage to the nerves of your legs and arms (such as numbness), joint pain, memory loss and breast enlargement in men (gynaecomastia)

**Side effects of not known frequency (frequency cannot be estimated from the available data):**

- Diarrhoea (loose stools), cough, shortness of breath, oedema (swelling), sleep disturbances, including insomnia and nightmares, sexual difficulties, depression, breathing problems, including persistent cough and/or shortness of breath or fever, tendon injury and muscle weakness that is constant
- myasthenia gravis (a disease causing general muscle weakness including in some cases muscles used when breathing), ocular myasthenia (a disease causing eye muscle weakness). Talk to your doctor if you experience weakness in your arms or legs that worsens after periods of activity, double vision or drooping of your eyelids, difficulty swallowing, or shortness of breath.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Rosuvastatin STADA

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the box/blisters/label after EXP. The expiry date refers to the last day of the month.

Store in the original package in order to protect the product from light.

This medicinal product does not require any special temperature storage conditions.

Do not throw away any medicines via wastewater <or household waste>. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

**What Rosuvastatin STADA contains:**

The active substance in Rosuvastatin STADA is rosuvastatin.

5 mg: Each film-coated tablet contains 5 mg rosuvastatin (as rosuvastatin calcium).

10 mg: Each film-coated tablet contains 10 mg rosuvastatin (as rosuvastatin calcium).

20 mg: Each film-coated tablet contains 20 mg rosuvastatin (as rosuvastatin calcium).

40 mg: Each film-coated tablet contains 40 mg rosuvastatin (as rosuvastatin calcium).

The other ingredients are:

Tablet core:

Lactose monohydrate  
Povidone K 30  
Crospovidone type A  
Cellulose, powdered  
Copovidone  
Silica, colloidal anhydrous  
Magnesium stearate

Tablet coat:  
Polyvinyl alcohol  
Macrogol (MW 3350)  
Titanium dioxide (E 171)  
Talc.

What Rosuvastatin STADA looks like and contents of the pack

Rosuvastatin STADA 5 mg film-coated tablets  
White, round, biconvex film-coated tablets with a diameter of 5 mm.

Rosuvastatin STADA 10 mg film-coated tablets  
White, round, biconvex film-coated tablets with a diameter of 7 mm.

Rosuvastatin STADA 20 mg film-coated tablets  
White, round, biconvex film-coated tablets with a diameter of 9 mm.

Rosuvastatin STADA 40 mg film-coated tablets  
White, oval, biconvex film-coated tablets with a length of 16.3 and a width of 7.6 mm.

The tablets are supplied in blisters manufactured of a combined oPA-Alu-PVC form foil sealed with an aluminium foil.

Package size(s):

|       |                                                                                                              |
|-------|--------------------------------------------------------------------------------------------------------------|
| 5 mg  | 7, 14, 15, 20, 28, 28 x1(single dose units), 30, 50x1(single dose units), 50, 60, 90, 98, 100, 154, 196, 252 |
| 10 mg | 7, 14, 15, 20, 28, 28 x1(single dose units), 30, 50x1(single dose units), 50, 60, 90, 98, 100, 154, 196, 252 |
| 20 mg | 7, 14, 15, 20, 28, 28 x1(single dose units), 30, 50x1(single dose units), 50, 60, 90, 98, 100, 154, 196, 252 |
| 40 mg | 28, 28 x1(single dose units), 30, 50x1(single dose units), 90, 98, 100, 154, 196, 252                        |

film-coated tablets.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer**

<To be completed nationally>

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

[To be completed nationally]



'This medicinal product is authorised in the Member States of the EEA under the following names:

|    |                                                                             |
|----|-----------------------------------------------------------------------------|
| AT | Rosuvastatin STADA 5 / 10 / 20 / 40 mg Filmtabletten                        |
| BE | Rosuvastatine EG 5 / 10 / 20 / 40 mg filmomhulde tabletten                  |
| DK | Rosuvastatin STADA                                                          |
| ES | Rosuvastatina STADA 5 / 10 / 20 mg comprimidos recubiertos con película EFG |
| FI | Rosuvastatin STADA 5 / 10 / 20 / 40 mg tabletti, kalvopäällysteinen         |
| FR | ROSUVASTATINE EG 5 / 10 / 20 mg, comprimé pelliculé                         |
| IE | Rosuvastatin Clonmel 5 / 10 / 20 / 40 mg film-coated tablets                |
| IT | ROSUVASTATINA EG                                                            |
| LU | Rosuvastatine EG 5 / 10 / 20 / 40 mg comprimés pelliculés                   |
| PT | Rosuvastatina Ciclum                                                        |
| SE | Rosuvastatin STADA 5 / 10 / 20 / 40 mg filmdragerad tablett                 |

**This leaflet was last revised in 2024-02-22.**