

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Sitagliptin/Dapagliflozin Teva 100 mg/10 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 100 mg sitagliptin (as sitagliptin hydrochloride monohydrate) and 10 mg dapagliflozin (as dapagliflozin amorphous base).

Excipient(s) with known effect

Each film-coated tablet contains 91 mg of lactose (as lactose monohydrate).

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Light brown, round film-coated tablet with “101” engraved on one side and with diameter 9.9 mm.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

[Product name], fixed dose combination of sitagliptin and dapagliflozin, is indicated as an adjunct to diet and exercise in adult patients aged 18 years and older with type 2 diabetes mellitus:

- to improve glycaemic control when metformin and sitagliptin do not provide adequate glycaemic control,
- when already being treated with the free combination of sitagliptin and dapagliflozin given concurrently, at the same dose level as in the fixed dose combination, but as separate products.

4.2 Posology and method of administration

Posology

The recommended dose of [Product name] is one tablet once daily.

Missed dose

If a dose is missed and it is ≥ 12 hours until the next dose, the dose should be taken. If a dose is missed and it is < 12 hours until the next dose, the missed dose should be skipped and the next dose taken at the usual time.

Special populations

Renal impairment

Because dosage adjustment should be made for sitagliptin in patients with renal impairment, assessment of renal function is recommended prior to initiation of [Product

name] and periodically thereafter.

For patients with mild renal impairment (glomerular filtration rate [GFR] ≥ 60 to < 90 mL/min) or patients with moderate renal impairment (GFR ≥ 45 to < 60 mL/min) no dose adjustment is required.

[Product name] should not be used in patients with a glomerular filtration rate (GFR) persistently < 45 mL/min, or in patients with end-stage renal disease (ESRD).

Hepatic impairment

No dose adjustment is necessary for patients with mild to moderate hepatic impairment.

[Product name] should not be used in patients with severe hepatic impairment.

Elderly

No dose adjustment is necessary based on age (see section 4.4)

Paediatric population

The safety and efficacy of this medicinal product in children and adolescents aged 0 to < 18 years have not yet been established. No data are available.

Method of administration

[Product name] is taken orally once daily.

To ensure accurate dosing and avoid unpleasant taste, the film-coated tablets should be swallowed whole and can be taken with or without food.

4.3 Contraindications

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

General

[Product name] should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis (see “Diabetic ketoacidosis”).

Acute pancreatitis

Use of DPP-4 inhibitors has been associated with a risk of developing acute pancreatitis. Patients should be informed of the characteristic symptom of acute pancreatitis: persistent, severe abdominal pain. Resolution of pancreatitis has been observed after discontinuation of this medicinal product (with or without supportive treatment), but very rare cases of necrotising or haemorrhagic pancreatitis and/or death have been reported. If pancreatitis is suspected, this medicinal product and other potentially suspect medicinal products should be discontinued; if acute pancreatitis is confirmed, it should not be restarted. Caution should be exercised in patients with a history of pancreatitis.

Hypoglycaemia when used in combination with other anti-hyperglycaemic medicinal products

In clinical trials of sitagliptin as monotherapy and as part of combination therapy with medicinal products not known to cause hypoglycaemia (i.e. metformin and/or a PPAR γ

agonist), rates of hypoglycaemia reported with sitagliptin were similar to rates in patients taking placebo. Hypoglycaemia has been observed when sitagliptin was used in combination with insulin or a sulphonylurea. Therefore, to reduce the risk of hypoglycaemia, a lower dose of sulphonylurea or insulin may be considered.

Renal impairment

[Product name] should not be used in patients with a glomerular filtration rate (GFR) persistently < 45 mL/min, or in patients with end-stage renal disease (ESRD).

The glucose lowering efficacy of dapagliflozin is dependent on renal function, and is reduced in patients with GFR < 45 mL/min and is likely absent in patients with severe renal impairment (see sections 4.2, 5.1 and 5.2). In one study in patients with type 2 diabetes mellitus with moderate renal impairment (GFR < 60 mL/min), a higher proportion of patients treated with dapagliflozin had adverse reactions of increase in creatinine, phosphorus, parathyroid hormone (PTH) and hypotension, compared with placebo.

Hypersensitivity reactions

Post-marketing reports of serious hypersensitivity reactions in patients treated with sitagliptin have been reported. These reactions include anaphylaxis, angioedema, and exfoliative skin conditions including Stevens-Johnson syndrome. Onset of these reactions occurred within the first 3 months after initiation of treatment, with some reports occurring after the first dose. If a hypersensitivity reaction is suspected, [Product name] should be discontinued. Other potential causes for the event should be assessed, and alternative treatment for diabetes initiated (see section 4.8).

Bullous pemphigoid

There have been post-marketing reports of bullous pemphigoid in patients taking DPP-4 inhibitors including sitagliptin. If bullous pemphigoid is suspected, [Product name] should be discontinued.

Hepatic impairment

[Product name] should not be used in patients with severe hepatic impairment.

Use in patients at risk for volume depletion and/or hypotension

Due to dapagliflozin's mechanism of action, this medicinal product increases diuresis which may lead to the modest decrease in blood pressure observed in clinical studies (see section 5.1). It may be more pronounced in patients with very high blood glucose concentrations.

Caution should be exercised in patients for whom a dapagliflozin-induced drop in blood pressure could pose a risk, such as patients on anti-hypertensive therapy with a history of hypotension or elderly patients.

In case of intercurrent conditions that may lead to volume depletion (e.g. gastrointestinal illness), careful monitoring of volume status (e.g. physical examination, blood pressure measurements, laboratory tests including haematocrit and electrolytes) is recommended. Temporary interruption of treatment with this medicinal product is recommended for patients who develop volume depletion until the depletion is corrected (see section 4.8).

Diabetic ketoacidosis

Rare cases of diabetic ketoacidosis (DKA), including life-threatening and fatal cases, have been reported in patients treated with SGLT2 inhibitors, including dapagliflozin. In a number of cases, the presentation of the condition was atypical with only moderately increased blood glucose values, below 14 mmol/L (250 mg/dL).

The risk of DKA must be considered in the event of non-specific symptoms such as nausea,

vomiting, anorexia, abdominal pain, excessive thirst, difficulty breathing, confusion, unusual fatigue or sleepiness. Patients should be assessed for ketoacidosis immediately if these symptoms occur, regardless of blood glucose level.

In patients where DKA is suspected or diagnosed, this medicinal product should be discontinued immediately.

Treatment should be interrupted in patients who are hospitalised for major surgical procedures or acute serious medical illnesses. Monitoring of ketones is recommended in these patients. Measurement of blood ketone levels is preferred to urine. Treatment with this medicinal product may be restarted when the ketone values are normal and the patient's condition has stabilised.

Before initiating treatment with this medicinal product, factors in the patient history that may predispose to ketoacidosis should be considered.

Patients who may be at higher risk of DKA include patients with a low beta-cell function reserve (e.g. type 2 diabetes patients with low C-peptide or latent autoimmune diabetes in adults (LADA) or patients with a history of pancreatitis), patients with conditions that lead to restricted food intake or severe dehydration, patients for whom insulin doses are reduced and patients with increased insulin requirements due to acute medical illness, surgery or alcohol abuse. SGLT2 inhibitors should be used with caution in these patients.

Restarting SGLT2 inhibitor treatment in patients experiencing a DKA while on SGLT2 inhibitor treatment is not recommended, unless another clear precipitating factor is identified and resolved.

The safety and efficacy of the sitagliptin/dapagliflozin fixed dose combination in patients with type 1 diabetes have not been established and it should not be used for treatment of patients with type 1 diabetes. In type 1 diabetes mellitus studies with dapagliflozin, DKA was reported with common frequency.

Necrotising fasciitis of the perineum (Fournier's gangrene)

Postmarketing cases of necrotising fasciitis of the perineum (also known as Fournier's gangrene) have been reported in female and male patients taking SGLT2 inhibitors (see section 4.8). This is a rare but serious and potentially life-threatening event that requires urgent surgical intervention and antibiotic treatment.

Patients should be advised to seek medical attention if they experience a combination of symptoms of pain, tenderness, erythema, or swelling in the genital or perineal area, with fever or malaise. Be aware that either uro-genital infection or perineal abscess may precede necrotising fasciitis. If Fournier's gangrene is suspected, [Product name] should be discontinued and prompt treatment (including antibiotics and surgical debridement) should be instituted.

Urinary tract infections

Urinary glucose excretion may be associated with an increased risk of urinary tract infection; therefore, temporary interruption of this medicinal product should be considered when treating pyelonephritis or urosepsis.

Elderly (≥ 65 years)

Elderly patients may be at a greater risk for volume depletion and are more likely to be treated with diuretics.

Elderly patients are more likely to have impaired renal function, and/or to be treated with

anti-hypertensive medicinal products that may cause changes in renal function such as angiotensin-converting enzyme inhibitors (ACE-I) and angiotensin II type 1 receptor blockers (ARB). The same recommendations for renal function apply to elderly patients as to all patients (see sections 4.2, 4.4, 4.8 and 5.1).

Cardiac failure

Experience with dapagliflozin in NYHA class IV is limited.

Infiltrative cardiomyopathy

Administration of dapagliflozin has not been studied in patients with infiltrative cardiomyopathy.

Increased haematocrit

Increased haematocrit has been observed with dapagliflozin treatment (see section 4.8). Patients with pronounced elevations in haematocrit should be monitored and investigated for underlying haematological disease.

Lower limb amputations

An increase in cases of lower limb amputation (primarily of the toe) has been observed in long-term, clinical studies in type 2 diabetes mellitus with SGLT2 inhibitors. It is unknown whether this constitutes a class effect. It is important to counsel patients with diabetes on routine preventative foot care.

Urine laboratory assessments

Due to the mechanism of action of dapagliflozin, patients taking this medicinal product will test positive for glucose in their urine.

Lactose

[Product name] film-coated tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Sodium

[Product name] contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially “sodium-free”.

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacodynamic interactions

Diuretics

Dapagliflozin may add to the diuretic effect of thiazide and loop diuretics and may increase the risk of dehydration and hypotension (see section 4.4).

Insulin and insulin secretagogues

Insulin and insulin secretagogues, such as sulphonylureas, cause hypoglycaemia. Therefore, a lower dose of insulin or an insulin secretagogue may be required to reduce the risk of hypoglycaemia when used in combination with this medicinal product in patients with type 2 diabetes mellitus (see sections 4.2 and 4.8).

Pharmacokinetic interactions

Lack of a pharmacokinetic interaction between dapagliflozin and sitagliptin has been demonstrated in healthy subjects after a single dose of dapagliflozin and sitagliptin co-administered.

Pharmacokinetic drug interaction studies with [Product name] have not been performed; however, such studies have been conducted with the individual active substances, sitagliptin and dapagliflozin.

Sitagliptin

In vitro studies indicated that the primary enzyme responsible for the limited metabolism of sitagliptin is CYP3A4, with contribution from CYP2C8.

Dapagliflozin

The metabolism of dapagliflozin is primarily via glucuronide conjugation mediated by UDP glucuronosyltransferase 1A9 (UGT1A9).

In *in vitro* studies, dapagliflozin neither inhibited cytochrome P450 (CYP) 1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP3A4, nor induced CYP1A2, CYP2B6 or CYP3A4. Therefore, dapagliflozin is not expected to alter the metabolic clearance of coadministered medicinal products that are metabolised by these enzymes.

Effects of other medicinal products on sitagliptin

Clinical data described below suggest that the risk for clinically meaningful interactions by co-administered medicinal products is low.

In vitro studies indicated that the primary enzyme responsible for the limited metabolism of sitagliptin is CYP3A4, with contribution from CYP2C8. In patients with normal renal function, metabolism, including via CYP3A4, plays only a small role in the clearance of sitagliptin. Metabolism may play a more significant role in the elimination of sitagliptin in the setting of severe renal impairment or ESRD. For this reason, it is possible that potent CYP3A4 inhibitors (i.e. ketoconazole, itraconazole, ritonavir, clarithromycin) could alter the pharmacokinetics of sitagliptin in patients with severe renal impairment or ESRD. The effect of potent CYP3A4 inhibitors in the setting of renal impairment has not been assessed in a clinical study.

In vitro transport studies showed that sitagliptin is a substrate for p-glycoprotein and organic anion transporter-3 (OAT3). OAT3 mediated transport of sitagliptin was inhibited *in vitro* by probenecid, although the risk of clinically meaningful interactions is considered to be low. Concomitant administration of OAT3 inhibitors has not been evaluated *in vivo*.

Metformin: Co-administration of multiple twice-daily doses of 1 000 mg metformin with 50 mg sitagliptin did not meaningfully alter the pharmacokinetics of sitagliptin in patients with type 2 diabetes.

Ciclosporin: A study was conducted to assess the effect of ciclosporin, a potent inhibitor of p-glycoprotein, on the pharmacokinetics of sitagliptin. Co-administration of a single 100 mg oral dose of sitagliptin and a single 600 mg oral dose of ciclosporin increased the AUC and C_{max} of sitagliptin by approximately 29% and 68%, respectively. These changes in sitagliptin pharmacokinetics were not considered to be clinically meaningful. The renal clearance of sitagliptin was not meaningfully altered. Therefore, meaningful interactions would not be expected with other p-glycoprotein inhibitors.

Effect of other medicinal products on dapagliflozin

Interaction studies conducted in healthy subjects, using mainly a single-dose design, suggest that the pharmacokinetics of dapagliflozin are not altered by metformin, pioglitazone, glimepiride, voglibose, hydrochlorothiazide, bumetanide, valsartan, or simvastatin.

Following coadministration of dapagliflozin with rifampicin (an inducer of various active transporters and drug-metabolising enzymes) a 22% decrease in dapagliflozin systemic exposure (AUC) was observed, but with no clinically meaningful effect on 24-hour urinary glucose excretion. No dose adjustment is recommended. A clinically relevant effect with other inducers (e.g. carbamazepine, phenytoin, phenobarbital) is not expected.

Following coadministration of dapagliflozin with mefenamic acid (an inhibitor of UGT1A9), a 55% increase in dapagliflozin systemic exposure was seen, but with no clinically meaningful effect on 24-hour urinary glucose excretion. No dose adjustment is recommended.

Effects of sitagliptin on other medicinal products

Digoxin: Sitagliptin had a small effect on plasma digoxin concentrations. Following administration of 0.25 mg digoxin concomitantly with 100 mg of sitagliptin daily for 10 days, the plasma AUC of digoxin was increased on average by 11%, and the plasma C_{max} on average by 18%. No dose adjustment of digoxin is recommended. However, patients at risk of digoxin toxicity should be monitored for this when sitagliptin and digoxin are administered concomitantly.

In vitro data suggest that sitagliptin does not inhibit nor induce CYP450 isoenzymes. In clinical studies, sitagliptin did not meaningfully alter the pharmacokinetics of metformin, glyburide, simvastatin, rosiglitazone, warfarin, or oral contraceptives, providing *in vivo* evidence of a low propensity for causing interactions with substrates of CYP3A4, CYP2C8, CYP2C9, and organic cationic transporter (OCT). Sitagliptin may be a mild inhibitor of p-glycoprotein *in vivo*.

Effect of dapagliflozin on other medicinal products

Dapagliflozin may increase renal lithium excretion and the blood lithium levels may be decreased. Serum concentration of lithium should be monitored more frequently after dapagliflozin initiation and dose changes. Please refer the patient to the lithium prescribing doctor in order to monitor serum concentration of lithium.

In interaction studies conducted in healthy subjects, using mainly a single-dose design, dapagliflozin did not alter the pharmacokinetics of metformin, pioglitazone, sitagliptin, glimepiride, hydrochlorothiazide, bumetanide, valsartan, digoxin (a P-gp substrate) or warfarin (S-warfarin, a CYP2C9 substrate), or the anticoagulatory effects of warfarin as measured by INR. Combination of a single dose of dapagliflozin 20 mg and simvastatin (a CYP3A4 substrate) resulted in a 19% increase in AUC of simvastatin and 31% increase in AUC of simvastatin acid. The increase in simvastatin and simvastatin acid exposures are not considered clinically relevant.

Interference with 1,5-anhydroglucitol (1,5-AG) assay

Monitoring glycaemic control with 1,5-AG assay is not recommended as measurements of 1,5-AG are unreliable in assessing glycaemic control in patients taking SGLT2 inhibitors. Use of alternative methods to monitor glycaemic control is advised.

Paediatric population

Interaction studies have only been performed in adults.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no adequate data from the use of sitagliptin and dapagliflozin in pregnant women.

Studies in animals with sitagliptin have shown reproductive toxicity at high doses (see section 5.3). The potential risk for humans is unknown. Studies with dapagliflozin in rats have shown toxicity to the developing kidney in the time period corresponding to the second and third trimesters of human pregnancy (see section 5.3). Due to lack of human data, the use of [Product name] is not recommended during pregnancy.

When pregnancy is detected, treatment with [Product name] should be discontinued.

Breast-feeding

It is unknown whether sitagliptin and dapagliflozin and/or its metabolites are excreted in human breast milk. Animal studies have shown excretion of sitagliptin in breast milk. Available pharmacodynamic/toxicological data in animals have shown excretion of dapagliflozin/metabolites in milk, as well as pharmacologically-mediated effects in nursing offspring (see section 5.3). A risk to the newborns/infants cannot be excluded. [Product name] should not be used during breast-feeding.

Fertility

The effect of sitagliptin and dapagliflozin on fertility in humans has not been studied. Animal data do not suggest an effect of treatment with sitagliptin or dapagliflozin on male and female fertility.

4.7 Effects on ability to drive and use machines

[Product name] has no or negligible influence on the ability to drive and use machines. However, when driving or using machines, it should be taken into account that dizziness and somnolence have been reported with the use of sitagliptin. In addition, patients should be alerted to the risk of hypoglycaemia if used in combination with a sulphonylurea or with insulin.

4.8 Undesirable effects

Summary of the safety profile

Sitagliptin/dapagliflozin combination

[Product name] has been demonstrated to be bioequivalent with co-administered sitagliptin and dapagliflozin (see section 2.5.2). There have been no therapeutic clinical trials conducted with [Product name] film-coated tablets.

Sitagliptin: Serious adverse reactions including pancreatitis and hypersensitivity reactions have been reported. Hypoglycaemia has been reported in combination with sulphonylurea (4.7%-13.8%) and insulin (9.6%) (see section 4.4).

Dapagliflozin: In the clinical studies in type 2 diabetes, more than 15,000 patients have been treated with dapagliflozin.

The primary assessment of safety and tolerability was conducted in a pre-specified pooled analysis of 13 short-term (up to 24 weeks) placebo-controlled studies with 2,360 subjects treated with dapagliflozin 10 mg and 2,295 treated with placebo.

In the dapagliflozin cardiovascular outcomes study in type 2 diabetes mellitus (DECLARE study, see section 5.1), 8,574 patients received dapagliflozin 10 mg and 8,569 received

placebo for a median exposure time of 48 months. In total, there were 30,623 patient-years of exposure to dapagliflozin.

The most frequently reported adverse reactions across the clinical studies were genital infections.

Chronic kidney disease

In the dapagliflozin renal outcome study in patients with chronic kidney disease (DAPA-CKD), 2,149 patients were treated with dapagliflozin 10 mg and 2,149 patients with placebo for a median exposure time of 27 months. The patient population included patients with type 2 diabetes mellitus and without diabetes, with eGFR ≥ 25 to ≤ 75 mL/min/1.73 m², and albuminuria (urine albumin creatinine ratio [UACR] ≥ 200 and ≤ 5000 mg/g). Treatment was continued if eGFR fell to levels below 25 mL/min/1.73 m². The overall safety profile of dapagliflozin in patients with chronic kidney disease was consistent with the known safety profile of dapagliflozin.

Tabulated list of adverse reactions

The safety profile is based on clinical trials and post-marketing experience with the mono-components. Adverse reactions are listed below (Table 1) by system organ class and frequency. Frequencies are defined as: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1\,000$ to $< 1/100$); rare ($\geq 1/10\,000$ to $< 1/1\,000$); very rare ($< 1/10\,000$) and not known (cannot be estimated from the available data).

Table1. Adverse reactions identified from placebo-controlled clinical studies^a and post-marketing experience with the mono-components

System Organ Class	Frequency of adverse reaction					
	Very common	Common	Uncommon	Rare	Very Rare	Not known
Infections and infestations		Vulvovaginitis, balanitis and related genital infections ^{1,*} , ^b Urinary tract infection ^{*,c}	Fungal infection ^{1, **}		Necrotising fasciitis of the perineum (Fournier's gangrene) ¹	
Blood and lymphatic system disorders				Thrombocytopenia ²		
Immune system disorders						Hypersensitivity reactions including anaphylactic responses ^{2,3}
Metabolism and nutrition disorders	Hypoglycaemia (when used with SU or INS) ^{1,2}		Volume depletion ^{1, d} Thirst ^{1, **}	Diabetic ketoacidosis (when used in T2DM) ^{1, i}		

System Organ Class	Frequency of adverse reaction					
	Very common	Common	Uncommon	Rare	Very Rare	Not known
Nervous system disorders		Headache ² , dizziness ^{1,2}				
Respiratory, thoracic and mediastinal disorders						Interstitial lung disease ³
Gastrointestinal disorders			Constipation ^{1,2,**} Dry mouth ^{1,**}			Vomiting ^{2,3} Acute pancreatitis ^{2,3} Fatal and non-fatal haemorrhagic and necrotizing pancreatitis ^{2,3}
Skin and subcutaneous tissue disorders		Rash ^{1,h,2,3}	Pruritus ^{2,3}		Angioedema ¹	Urticaria ^{2,3} Cutaneous vasculitis ^{2,3} Exfoliative skin conditions including Stevens-Johnson syndrome ^{2,3} Bullous pemphigoid ^{2,3}
Musculoskeletal and connective tissue disorders		Back pain ^{1,2,3,*}				Arthralgia ^{2,3} Myalgia ^{2,3} Arthropathy ^{2,3}
Renal and urinary disorders		Dysuria Polyuria ^{1,*,e}	Nocturia ^{1,**}		Tubulointerstitial nephritis ¹	Impaired renal function ^{2,3} Acute renal failure ^{2,3}
Reproductive system and breast disorders			Vulvovaginal pruritus ^{1,*}			

System Organ Class	Frequency of adverse reaction					
	Very common	Common	Uncom mon	Rare	Very Rare	Not known
			Pruritus genital ^{1,**}			
Investigati ons		Haematocrit increased ^{1,f} Creatinine renal clearance decreased during initial treatment ¹ Dyslipidaemia ^{1,g}	Blood creatinin e increased during initial treatment ^{1,**} Blood urea increased ^{1,**} Weight decrease d ^{1,**}			

¹ Related to DAPA

² Related to SITA

³ Adverse reactions were identified through post-marketing surveillance for SITA.

^a The table shows up to 24-week (short-term) data regardless of glycaemic rescue for DAPA adverse reactions.

^b Vulvovaginitis, balanitis and related genital infections includes, e.g. the predefined preferred terms: vulvovaginal mycotic infection, vaginal infection, balanitis, genital infection fungal, vulvovaginal candidiasis, vulvovaginitis, balanitis candida, genital candidiasis, genital infection, genital infection male, penile infection, vulvitis, vaginitis bacterial, vulval abscess.

^c UTI includes the following preferred terms, listed in order of frequency reported: urinary tract infection, cystitis, Escherichia urinary tract infection, genitourinary tract infection, pyelonephritis, trigonitis, urethritis, kidney infection and prostatitis.

^d Volume depletion includes, e.g. the predefined preferred terms: dehydration, hypovolaemia, hypotension.

^e Polyuria includes the preferred terms: pollakiuria, polyuria, urine output increased.

^f Mean changes from baseline in haematocrit were 2.30% for DAPA 10 mg versus -0.33% for placebo. Haematocrit values >55% were reported in 1.3% of the subjects treated with DAPA 10 mg versus 0.4% of placebo subjects.

^g Mean percent change from baseline for DAPA 10 mg versus placebo, respectively, was: total cholesterol 2.5% versus 0.0%; HDL cholesterol 6.0% versus 2.7%; LDL cholesterol 2.9% versus -1.0%; triglycerides -2.7% versus -0.7%.

^h Adverse reaction was identified through postmarketing surveillance. Rash includes the following preferred terms, listed in order of frequency in clinical studies: rash, rash generalised, rash pruritic, rash macular, rash maculo-papular, rash pustular, rash vesicular, and rash erythematous. In active- and placebo-controlled clinical studies (DAPA, N=5936, All control, N=3403), the frequency of rash was similar for DAPA (1.4%) and all control (1.4%), respectively.

ⁱ Reported in the cardiovascular outcomes study in patients with type 2 diabetes (DECLARE). Frequency is based on annual rate.

* Reported in ≥ 2% of subjects and ≥ 1% more and at least 3 more subjects treated with DAPA 10 mg compared to placebo

** Reported by the investigator as possibly related, probably related or related to study treatment and reported in ≥ 0.2% of subjects and ≥ 0.1% more and at least 3 more subjects treated with DAPA 10 mg compared to placebo.

Description of selected adverse reactions

Sitagliptin

In addition to the drug-related adverse experiences described above, adverse experiences reported regardless of causal relationship to medication and occurring in at least 5% and more commonly in patients treated with sitagliptin included upper respiratory tract infection and nasopharyngitis. Additional adverse experiences reported regardless of causal relationship to medication that occurred more frequently in patients treated with sitagliptin (not reaching the 5% level, but occurring with an incidence of > 0.5% higher with sitagliptin than that in the control group) included osteoarthritis and pain in extremity.

Some adverse reactions were observed more frequently in studies of combination use of sitagliptin with other anti-diabetic medicinal products than in studies of sitagliptin monotherapy. These included hypoglycaemia (frequency very common with the combination of sulphonylurea and metformin), influenza (common with insulin (with or without metformin)), nausea and vomiting (common with metformin), flatulence (common with metformin or pioglitazone), constipation (common with the combination of sulphonylurea and metformin), peripheral oedema (common with pioglitazone or the combination of pioglitazone and metformin), somnolence and diarrhoea (uncommon with metformin), and dry mouth (uncommon with insulin (with or without metformin)).

TECOS Cardiovascular Safety Study

The Trial Evaluating Cardiovascular Outcomes with Sitagliptin (TECOS) included 7,332 patients treated with sitagliptin, 100 mg daily (or 50 mg daily if the baseline eGFR was ≥ 30 and < 50 mL/min/1.73 m²), and 7,339 patients treated with placebo in the intention-to-treat population.

Both treatments were added to usual care targeting regional standards for HbA1c and CV risk factors. The overall incidence of serious adverse events in patients receiving sitagliptin was similar to that in patients receiving placebo.

In the intention-to-treat population, among patients who were using insulin and/or a sulphonylurea at baseline, the incidence of severe hypoglycaemia was 2.7% in sitagliptin-treated patients and 2.5% in placebo-treated patients; among patients who were not using insulin and/or a sulphonylurea at baseline, the incidence of severe hypoglycaemia was 1.0% in sitagliptin-treated patients and 0.7% in placebo-treated patients. The incidence of adjudication-confirmed pancreatitis events was 0.3% in sitagliptin-treated patients and 0.2% in placebo-treated patients.

Dapagliflozin

Vulvovaginitis, balanitis and related genital infections

In the 13-study safety pool, vulvovaginitis, balanitis and related genital infections were reported in 5.5% and 0.6% of subjects who received dapagliflozin 10 mg and placebo, respectively. Most infections were mild to moderate, and subjects responded to an initial course of standard treatment and rarely resulted in discontinuation from dapagliflozin treatment. These infections were more frequent in females (8.4% and 1.2% for dapagliflozin and placebo, respectively), and subjects with a prior history were more likely to have a recurrent infection.

In the DECLARE study, the numbers of patients with serious adverse events of genital infections were few and balanced: 2 patients in each of the dapagliflozin and placebo groups.

In the DAPA-CKD study, there were 3 (0.1%) patients with serious adverse events of genital

infections in the dapagliflozin group and none in the placebo group. There were 3 (0.1%) patients with adverse events leading to discontinuation due to genital infections in the dapagliflozin group and none in the placebo group. Serious adverse events of genital infections or adverse events leading to discontinuation due to genital infections were not reported for any patients without diabetes

Cases of phimosis/acquired phimosis have been reported with dapagliflozin concurrent with genital infections and in some cases, circumcision was required.

Necrotising fasciitis of the perineum (Fournier's gangrene)

Cases of Fournier's gangrene have been reported postmarketing in patients taking SGLT2 inhibitors, including dapagliflozin (see section 4.4).

In the DECLARE study with 17,160 type 2 diabetes mellitus patients and a median exposure time of 48 months, a total of 6 cases of Fournier's gangrene were reported, one in the dapagliflozin-treated group and 5 in the placebo group.

Hypoglycaemia

The frequency of hypoglycaemia depended on the type of background therapy used in the clinical studies in diabetes mellitus.

For studies of dapagliflozin in monotherapy, as add-on to metformin or as add-on to sitagliptin (with or without metformin), the frequency of minor episodes of hypoglycaemia was similar (< 5%) between treatment groups, including placebo up to 102 weeks of treatment. Across all studies, major events of hypoglycaemia were uncommon and comparable between the groups treated with dapagliflozin or placebo. Studies with add-on sulphonylurea and add-on insulin therapies had higher rates of hypoglycaemia (see section 4.5).

In an add-on to glimepiride study, at weeks 24 and 48, minor episodes of hypoglycaemia were reported more frequently in the group treated with dapagliflozin 10 mg plus glimepiride (6.0% and 7.9%, respectively) than in the placebo plus glimepiride group (2.1% and 2.1%, respectively).

In an add-on to insulin study, episodes of major hypoglycaemia were reported in 0.5% and 1.0% of subjects treated with dapagliflozin 10 mg plus insulin at weeks 24 and 104, respectively, and in 0.5% of subjects treated with placebo plus insulin groups at weeks 24 and 104. At weeks 24 and 104, minor episodes of hypoglycaemia were reported, respectively, in 40.3% and 53.1% of subjects who received dapagliflozin 10 mg plus insulin and in 34.0% and 41.6% of the subjects who received placebo plus insulin.

In an add-on to metformin and a sulphonylurea study, up to 24 weeks, no episodes of major hypoglycaemia were reported. Minor episodes of hypoglycaemia were reported in 12.8% of subjects who received dapagliflozin 10 mg plus metformin and a sulphonylurea and in 3.7% of subjects who received placebo plus metformin and a sulphonylurea.

In the DECLARE study, no increased risk of major hypoglycaemia was observed with dapagliflozin therapy compared with placebo. Major events of hypoglycaemia were reported in 58 (0.7%) patients treated with dapagliflozin and 83 (1.0%) patients treated with placebo.

In the DAPA-CKD study, major events of hypoglycaemia were reported in 14 (0.7%) patients in the dapagliflozin group and 28 (1.3%) patients in the placebo group and observed only in patients with type 2 diabetes mellitus.

Volume depletion

In the 13-study safety pool, reactions suggestive of volume depletion (including, reports of dehydration, hypovolaemia or hypotension) were reported in 1.1% and 0.7% of subjects who received dapagliflozin 10 mg and placebo, respectively; serious reactions occurred in < 0.2% of subjects balanced between dapagliflozin 10 mg and placebo (see section 4.4).

In the DECLARE study, the numbers of patients with events suggestive of volume depletion were balanced between treatment groups: 213 (2.5%) and 207 (2.4%) in the dapagliflozin and placebo groups, respectively. Serious adverse events were reported in 81 (0.9%) and 70 (0.8%) in the dapagliflozin and placebo group, respectively. Events were generally balanced between treatment groups across subgroups of age, diuretic use, blood pressure and angiotensin converting enzyme inhibitors (ACE-I)/angiotensin II type 1 receptor blockers (ARB) use. In patients with eGFR < 60 mL/min/1.73 m² at baseline, there were 19 events of serious adverse events suggestive of volume depletion in the dapagliflozin group and 13 events in the placebo group.

In the DAPA-CKD study, the numbers of patients with events suggestive of volume depletion were 120 (5.6%) in the dapagliflozin group and 84 (3.9%) in the placebo group. There were 16 (0.7%) patients with serious events of symptoms suggestive of volume depletion in the dapagliflozin group and 15 (0.7%) patients in the placebo group.

Diabetic ketoacidosis in type 2 diabetes mellitus

In the DECLARE study, with a median exposure time of 48 months, events of DKA were reported in 27 patients in the dapagliflozin 10 mg group and 12 patients in the placebo group. The events occurred evenly distributed over the study period. Of the 27 patients with DKA events in the dapagliflozin group, 22 had concomitant insulin treatment at the time of the event. Precipitating factors for DKA were as expected in a type 2 diabetes mellitus population (see section 4.4).

In the DAPA-CKD study, events of DKA were not reported in any patient in the dapagliflozin group and in 2 patients with type 2 diabetes mellitus in the placebo group.

Urinary tract infections

In the 13-study safety pool, urinary tract infections were more frequently reported for dapagliflozin 10 mg compared to placebo (4.7% versus 3.5%, respectively; see section 4.4). Most infections were mild to moderate, and subjects responded to an initial course of standard treatment and rarely resulted in discontinuation from dapagliflozin treatment. These infections were more frequent in females, and subjects with a prior history were more likely to have a recurrent infection.

In the DECLARE study, serious events of urinary tract infections were reported less frequently for dapagliflozin 10 mg compared with placebo, 79 (0.9%) events versus 109 (1.3%) events, respectively.

In the DAPA-CKD study, the numbers of patients with serious adverse events of urinary tract infections were 29 (1.3%) in the dapagliflozin group and 18 (0.8%) in the placebo group. There were 8 (0.4%) patients with adverse events leading to discontinuations due to urinary tract infections in the dapagliflozin group and 3 (0.1%) in the placebo group. The numbers of patients without diabetes reporting serious adverse events of urinary tract infections or adverse events leading to discontinuation due to urinary tract infections were similar between treatment groups (6 [0.9%] versus 4 [0.6%] for serious adverse events, and 1 [0.1%] versus 0 for adverse events leading to discontinuation, in the dapagliflozin and placebo groups, respectively).

Increased creatinine

Adverse reactions related to increased creatinine were grouped (e.g. decreased renal

creatinine clearance, renal impairment, increased blood creatinine and decreased glomerular filtration rate). In the 13-study safety pool, this grouping of reactions was reported in 3.2% and 1.8% of patients who received dapagliflozin 10 mg and placebo, respectively. In patients with normal renal function or mild renal impairment (baseline eGFR ≥ 60 mL/min/1.73 m²) this grouping of reactions were reported in 1.3% and 0.8% of patients who received dapagliflozin 10 mg and placebo, respectively. These reactions were more common in patients with baseline eGFR ≥ 30 and < 60 mL/min/1.73 m² (18.5% dapagliflozin 10 mg versus 9.3% placebo).

Further evaluation of patients who had renal-related adverse events showed that most had serum creatinine changes of ≤ 44 micromoles/L (≤ 0.5 mg/dL) from baseline. The increases in creatinine were generally transient during continuous treatment or reversible after discontinuation of treatment.

In the DECLARE study, including elderly patients and patients with renal impairment (eGFR less than 60 mL/min/1.73 m²), eGFR decreased over time in both treatment groups. At 1 year, mean eGFR was slightly lower, and at 4 years, mean eGFR was slightly higher in the dapagliflozin group compared with the placebo group.

In the DAPA-CKD study, eGFR decreased over time in both the dapagliflozin group and the placebo group. The initial (day 14) decrease in mean eGFR was -4.0 mL/min/1.73 m² in the dapagliflozin group and -0.8 mL/min/1.73 m² in the placebo group. At 28 months, change from baseline in eGFR was -7.4 mL/min/1.73 m² in the dapagliflozin group and -8.6 mL/min/1.73 m² in the placebo group

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in
[To be completed nationally]

4.9 Overdose

There is no information available on overdose with the sitagliptin/dapagliflozin fixed dose combination. In the event of an overdose, appropriate supportive treatment should be initiated as dictated by the patient's clinical status. It is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring (including obtaining an electrocardiogram), and institute supportive therapy if required.

Sitagliptin

During controlled clinical trials in healthy subjects, single doses of up to 800 mg sitagliptin were administered. Minimal increases in QTc, not considered to be clinically relevant, were observed in one study at a dose of 800 mg sitagliptin. There is no experience with doses above 800 mg in clinical studies. In Phase I multiple-dose studies, there were no dose-related clinical adverse reactions observed with sitagliptin with doses of up to 600 mg per day for periods of up to 10 days and 400 mg per day for periods of up to 28 days.

Sitagliptin is modestly dialysable. In clinical studies, approximately 13.5% of the dose was removed over a 3- to 4-hour haemodialysis session. Prolonged haemodialysis may be considered if clinically appropriate. It is not known if sitagliptin is dialysable by

peritoneal dialysis.

Dapagliflozin

Dapagliflozin did not show any toxicity in healthy subjects at single oral doses up to 500 mg (50 times the maximum recommended human dose). These subjects had detectable glucose in the urine for a dose-related period of time (at least 5 days for the 500 mg dose), with no reports of dehydration, hypotension or electrolyte imbalance, and with no clinically meaningful effect on QTc interval. The incidence of hypoglycaemia was similar to placebo. In clinical studies where once-daily doses of up to 100 mg (10 times the maximum recommended human dose) were administered for 2 weeks in healthy subjects and type 2 diabetes subjects, the incidence of hypoglycaemia was slightly higher than placebo and was not dose-related. Rates of adverse events including dehydration or hypotension were similar to placebo, and there were no clinically meaningful dose-related changes in laboratory parameters, including serum electrolytes and biomarkers of renal function. The removal of dapagliflozin by haemodialysis has not been studied.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes, combinations of oral blood glucose lowering drugs, ATC code: A10BD29

Mechanism of action

[Product name] combines sitagliptin and dapagliflozin with complementary mechanisms of action to improve glycaemic control. Sitagliptin, through the selective inhibition of dipeptidyl peptidase-4 (DPP-4), enhances glucose-mediated insulin secretion (incretin effect). Dapagliflozin, a selective inhibitor of sodium-glucose co-transporter 2 (SGLT2), inhibits renal glucose reabsorption independently of insulin. Actions of both medicinal products are regulated by the plasma glucose level.

Sitagliptin is a member of a class of oral anti-hyperglycaemic agents called DPP-4 inhibitors. The improvement in glycaemic control observed with sitagliptin may be mediated by enhancing the levels of active incretin hormones. Incretin hormones, including glucagon-like peptide-1 (GLP-1) and glucose-dependent insulinotropic polypeptide (GIP), are released by the intestine throughout the day, and levels are increased in response to a meal. The incretins are part of an endogenous system involved in the physiologic regulation of glucose homeostasis. When blood glucose concentrations are normal or elevated, GLP-1 and GIP increase insulin synthesis and release from pancreatic beta cells by intracellular signaling pathways involving cyclic AMP.

Dapagliflozin is a highly potent (K_i : 0.55 nM), selective and reversible inhibitor of sodium-glucose co-transporter 2 (SGLT2). Dapagliflozin blocks reabsorption of filtered glucose from the proximal renal tubule, effectively lowering blood glucose in a glucose dependent and insulin-independent manner. Dapagliflozin improves both fasting and post-prandial plasma glucose levels by reducing renal glucose reabsorption leading to urinary glucose excretion. This glucose excretion (glucuretic effect) is observed after the first dose, is continuous over the 24-hour dosing interval and is sustained for the duration of treatment. The amount of glucose removed by the kidney through this mechanism is dependent upon the blood glucose concentration and GFR. Thus, in subjects with normal blood glucose, dapagliflozin has a low propensity to cause hypoglycaemia. Dapagliflozin does not impair normal endogenous glucose production in response to hypoglycaemia. Dapagliflozin acts independently of insulin

secretion and insulin action. Improvement in homeostasis model assessment for beta cell function (HOMA beta-cell) has been observed in clinical studies with dapagliflozin. The SGLT2 is selectively expressed in the kidney. Dapagliflozin does not inhibit other glucose transporters important for glucose transport into peripheral tissues and is > 1,400 times more selective for SGLT2 versus SGLT1, the major transporter in the gut responsible for glucose absorption.

Inhibition of SGLT2 by dapagliflozin reduces reabsorption of glucose from the glomerular filtrate in the proximal renal tubule with a concomitant reduction in sodium reabsorption leading to urinary excretion of glucose and osmotic diuresis. Dapagliflozin therefore increases the delivery of sodium to the distal tubule which increases tubuloglomerular feedback and reduces intraglomerular pressure. This combined with osmotic diuresis leads to a reduction in volume overload, reduced blood pressure, and lower preload and afterload, which may have beneficial effects on cardiac remodelling and diastolic function, and preserve renal function. The cardiac and renal benefits of dapagliflozin are not solely dependent on the blood glucose-lowering effect. Other effects include an increase in haematocrit and reduction in body weight.

Pharmacodynamic effects

Treatment with GLP-1 or with DPP-4 inhibitors in animal models of type 2 diabetes has been demonstrated to improve beta cell responsiveness to glucose and stimulate insulin biosynthesis and release. With higher insulin levels, tissue glucose uptake is enhanced. In addition, GLP-1 lowers glucagon secretion from pancreatic alpha cells. Decreased glucagon concentrations, along with higher insulin levels, lead to reduced hepatic glucose production, resulting in a decrease in blood glucose levels. The effects of GLP-1 and GIP are glucose-dependent such that when blood glucose concentrations are low, stimulation of insulin release and suppression of glucagon secretion by GLP-1 are not observed. For both GLP-1 and GIP, stimulation of insulin release is enhanced as glucose rises above normal concentrations. Further, GLP-1 does not impair the normal glucagon response to hypoglycaemia. The activity of GLP-1 and GIP is limited by the DPP-4 enzyme, which rapidly hydrolyzes the incretin hormones to produce inactive products. Sitagliptin prevents the hydrolysis of incretin hormones by DPP-4, thereby increasing plasma concentrations of the active forms of GLP-1 and GIP. By enhancing active incretin levels, sitagliptin increases insulin release and decreases glucagon levels in a glucose-dependent manner. In patients with type 2 diabetes with hyperglycaemia, these changes in insulin and glucagon levels lead to lower haemoglobin A1c (HbA1c) and lower fasting and postprandial glucose concentrations. The glucose-dependent mechanism of sitagliptin is distinct from the mechanism of sulphonylureas, which increase insulin secretion even when glucose levels are low and can lead to hypoglycaemia in patients with type 2 diabetes and in normal subjects. Sitagliptin is a potent and highly selective inhibitor of the enzyme DPP-4 and does not inhibit the closely-related enzymes DPP-8 or DPP-9 at therapeutic concentrations.

Dapagliflozin's glucuretic effect is observed after the first dose, is continuous over the 24-hour dosing interval, and is sustained for the duration of treatment. Increases in the amount of glucose excreted in the urine were observed in healthy subjects and in subjects with type 2 diabetes mellitus following the administration of dapagliflozin. Approximately 70 g of glucose was excreted in the urine per day (corresponding to 280 kcal/day) at a dapagliflozin dose of 10 mg/day in subjects with type 2 diabetes mellitus for 12 weeks. Evidence of sustained glucose excretion was seen in subjects with type 2 diabetes mellitus given dapagliflozin 10 mg/day for up to 2 years. Urinary uric acid excretion was also increased transiently (for 3-7 days) and accompanied by a sustained reduction in serum uric acid concentration. At 24 weeks, reductions in serum uric acid concentrations ranged from -48.3 to -18.3 micromoles/L (-0.87 to -0.33 mg/dL).

Clinical efficacy and safety

Type 2 diabetes mellitus

The efficacy and safety of dapagliflozin as add-on therapy in patients with type 2 diabetes who were inadequately controlled with a dipeptidyl peptidase-4 inhibitor with or without metformin is supported by a published 24-week, multicenter, randomized, double-blind, placebo-controlled, parallel-group, phase 3 study with a 24-week blinded extension period. Four hundred and fifty-two patients were randomized¹ to receive dapagliflozin 10mg/day or placebo added to sitagliptin (100 mg/day) ± metformin (≥1,500 mg/day). At week 24, dapagliflozin significantly reduced mean HbA1c levels (−0.5% [−4.9 mmol/mol]) versus placebo (0.0% [+0.4 mmol/mol]). Dapagliflozin reduced body weight versus placebo (−2.1 vs. −0.3 kg) and reduced HbA1c levels in patients with baseline values ≥8.0% (−0.8% [−8.7 mmol/mol] vs. 0.0% [0.3 mmol/mol]) and fasting plasma glucose levels (−24.1 mg/dL [−1.3 mmol/L] 1 vs. 3.8 mg/dL [0.2 mmol/L]). Similar results were observed when data were analyzed separately for each stratum (by metformin background therapy). Glycemic and weight benefits observed at week 24 were maintained through week 48. Changes from baseline in systolic blood pressure at week 8 were not significantly different between treatment groups. Over 48 weeks, fewer patients receiving dapagliflozin were discontinued or rescued for failing to achieve glycemic targets compared with placebo. Adverse events were balanced between groups, and discontinuation rates were low. At week 48, signs and symptoms suggestive of genital infection were more frequent with dapagliflozin (9.8%) than with placebo (0.4%). Signs and symptoms suggestive of urinary tract infection were balanced between dapagliflozin (6.7%) and placebo (6.2%).

TECOS Cardiovascular Safety Study

The TECOS was a randomised study in 14,671 patients in the intention-to-treat population with an HbA1c of ≥ 6.5 to 8.0% with established CV disease who received sitagliptin (7,332) 100 mg daily (or 50 mg daily if the baseline eGFR was ≥30 and <50 mL/min/1.73 m²) or placebo (7,339) added to usual care targeting regional standards for HbA1c and CV risk factors. Patients with an eGFR <30 mL/min/1.73 m² were not to be enrolled in the study. The study population included 2,004 patients ≥75 years of age and 3,324 patients with renal impairment (eGFR <60 mL/min/1.73 m²).

Over the course of the study, the overall estimated mean (SD) difference in HbA1c between the sitagliptin and placebo groups was 0.29 % (0.01), 95 % CI (-0.32, -0.27); p <0.001.

The primary cardiovascular endpoint was a composite of the first occurrence of cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, or hospitalisation for unstable angina. Secondary cardiovascular endpoints included the first occurrence of cardiovascular death, nonfatal myocardial infarction, or nonfatal stroke; first occurrence of the individual components of the primary composite; all-cause mortality; and hospital admissions for congestive heart failure.

After a median follow up of 3 years, sitagliptin, when added to usual care, did not increase the risk of major adverse cardiovascular events or the risk of hospitalisation for heart failure compared to usual care without sitagliptin in patients with type 2 diabetes (Table 2).

¹ Among the 452 patients randomized, 451 were treated and 447 were included in the full analysis test

Table 2. Rates of Composite Cardiovascular Outcomes and Key Secondary Outcomes.

	Sitagliptin 100 mg		Placebo		Hazard Ratio (95% CI)	p-value [†]
	N (%)	Inciden ce rate per 100 patient years*	N (%)	Inciden ce rate per 100 patient years*		
Analysis in the Intention-to-Treat Population						
Number of patients	7,332		7,339		0.98 (0.89–1.08)	<0.001
Primary Composite Endpoint (Cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, or hospitalisation for unstable angina)	839 (11.4)	4.1	851 (11.6)	4.2		
Secondary Composite Endpoint (Cardiovascular death, nonfatal myocardial infarction, or nonfatal stroke)	745 (10.2)	3.6	746 (10.2)	3.6		
Secondary Outcome						
Cardiovascular death	380 (5.2)	1.7	366 (5.0)	1.7	1.03 (0.89-1.19)	0.711
All myocardial infarction (fatal and non-fatal)	300 (4.1)	1.4	316 (4.3)	1.5	0.95 (0.81–1.11)	0.487
All stroke (fatal and non-fatal)	178 (2.4)	0.8	183 (2.5)	0.9	0.97 (0.79–1.19)	0.760
Hospitalisation for unstable angina	116 (1.6)	0.5	129 (1.8)	0.6	0.90 (0.70–1.16)	0.419
Death from any cause	547 (7.5)	2.5	537 (7.3)	2.5	1.01 (0.90–1.14)	0.875
Hospitalisation for heart failure [‡]	228 (3.1)	1.1	229 (3.1)	1.1	1.00 (0.83–1.20)	0.983

*Incidence rate per 100 patient-years is calculated as $100 \times (\text{total number of patients with } \geq 1 \text{ event during eligible exposure period per total patient-years of follow-up})$.

[†] Based on a Cox model stratified by region. For composite endpoints, the p-values correspond to a test of non-inferiority seeking to show that the hazard ratio is less than 1.3. For all other endpoints, the p-values correspond to a test of differences in hazard rates.

[‡] The analysis of hospitalisation for heart failure was adjusted for a history of heart failure at baseline.

Dapagliflozin Effect on Cardiovascular Events (DECLARE)

Dapagliflozin Effect on Cardiovascular Events (DECLARE) was an international, multicentre, randomised, double-blind, placebo-controlled clinical study conducted to determine the effect of dapagliflozin compared with placebo on cardiovascular outcomes when added to current background therapy. All patients had type 2 diabetes mellitus and either at least two additional cardiovascular risk factors (age ≥ 55 years in men or ≥ 60 years in women and one or more of dyslipidaemia, hypertension or current tobacco use) or

established cardiovascular disease.

Of 17,160 randomised patients, 6,974 (40.6%) had established cardiovascular disease and 10,186 (59.4%) did not have established cardiovascular disease. 8,582 patients were randomised to dapagliflozin 10 mg and 8,578 to placebo, and were followed for a median of 4.2 years.

The mean age of the study population was 63.9 years, 37.4% were female. In total, 22.4% had had diabetes for ≤ 5 years, mean duration of diabetes was 11.9 years. Mean HbA1c was 8.3% and mean BMI was 32.1 kg/m².

At baseline, 10.0% of patients had a history of heart failure. Mean eGFR was 85.2 mL/min/1.73 m², 7.4% of patients had eGFR < 60 mL/min/1.73 m², and 30.3% of patients had micro- or macroalbuminuria (UACR ≥ 30 to ≤ 300 mg/g or > 300 mg/g, respectively).

Most patients (98%) used one or more diabetic medicinal products at baseline, including metformin (82%), insulin (41%) and sulfonylurea (43%).

The primary endpoints were time to first event of the composite of cardiovascular death, myocardial infarction or ischaemic stroke (MACE) and time to first event of the composite of hospitalisation for heart failure or cardiovascular death. The secondary endpoints were a renal composite endpoint and all-cause mortality.

Major adverse cardiovascular events


Dapagliflozin 10 mg demonstrated non-inferiority versus placebo for the composite of cardiovascular death, myocardial infarction or ischaemic stroke (one-sided $p < 0.001$).

Heart failure or cardiovascular death

Dapagliflozin 10 mg demonstrated superiority versus placebo in preventing the composite of hospitalisation for heart failure or cardiovascular death (Figure 1). The difference in treatment effect was driven by hospitalisation for heart failure, with no difference in cardiovascular death (Figure 2).

The treatment benefit of dapagliflozin over placebo was observed both in patients with and without established cardiovascular disease, with and without heart failure at baseline, and was consistent across key subgroups, including age, gender, renal function (eGFR) and region.

Figure 1: Time to first occurrence of hospitalisation for heart failure or cardiovascular death.



Patients at risk


Dapagliflozin:	8582	8517	8415	8322	8224	8110	7970	7497	5445	1626
Placebo:	8578	8485	8387	8259	8127	8003	7880	7367	5362	1573

Patients at risk is the number of patients at risk at the beginning of the period.

HR=Hazard ratio CI=Confidence interval.

Results on primary and secondary endpoints are displayed in Figure 2. Superiority of dapagliflozin over placebo was not demonstrated for MACE ($p=0.172$). The renal composite endpoint and all-cause mortality were therefore not tested as part of the confirmatory testing procedure.

Figure 2: Treatment effects for the primary composite endpoints and their components, and the secondary endpoints and components.



Renal composite endpoint defined as: sustained confirmed $\geq 40\%$ decrease in eGFR to eGFR < 60 mL/min/1.73 m² and/or end-stage kidney disease (dialysis ≥ 90 days or kidney transplantation, sustained confirmed eGFR < 15 mL/min/1.73 m²) and/or renal or cardiovascular death. p-values are two-sided. p-values for the secondary endpoints and for single components are nominal. Time to first event was analysed in a Cox proportional hazards model. The number of first events for the single components are the actual number of first events for each component and does not add up to the number of events in the composite endpoint. CI=confidence interval.

Nephropathy

Dapagliflozin reduced the incidence of events of the composite of confirmed sustained eGFR decrease, end-stage kidney disease, renal or cardiovascular death. The difference between groups was driven by reductions in events of the renal components; sustained eGFR decrease, end-stage kidney disease and renal death (Figure 2).

The hazard ratio (HR) for time to nephropathy (sustained eGFR decrease, end-stage kidney disease and renal death) was 0.53 (95% CI 0.43, 0.66) for dapagliflozin versus placebo.

In addition, dapagliflozin reduced the new onset of sustained albuminuria (HR 0.79 [95% CI 0.72, 0.87]) and led to greater regression of macroalbuminuria (HR 1.82 [95% CI 1.51, 2.20]) compared with placebo.

Renal impairment

Moderate renal impairment CKD 3A (eGFR ≥ 45 to < 60 mL/min/1.73 m²)

Dapagliflozin

The efficacy of dapagliflozin was assessed in a dedicated study in diabetic patients with an eGFR ≥ 45 to < 60 mL/min/1.73 m² who had inadequate glycaemic control on usual care. Treatment with dapagliflozin resulted in reductions in HbA1c and body weight compared with placebo (Table 3).

Table 3. Results at week 24 of a placebo-controlled study of dapagliflozin in diabetic patients with an eGFR ≥ 45 to < 60 mL/min/1.73 m².

	Dapagliflozin^a 10 mg	Placebo^a
N^b	159	161
HbA1c (%)		
Baseline (mean)	8.35	8.03
Change from baseline ^b	-0.37	-0.03
Difference from placebo ^b	-0.34*	
(95% CI)	(-0.53, -0.15)	
Body weight (kg)		
Baseline (mean)	92.51	88.30
Percent change from baseline ^c	-3.42	-2.02
Difference in percent change from placebo ^c	-1.43*	
(95% CI)	(-2.15, -0.69)	

^a Metformin or metformin hydrochloride were part of the usual care in 69.4% and 64.0% of the patients for the dapagliflozin and placebo groups, respectively.

^b Least squares mean adjusted for baseline value.

^c Derived from least squares mean adjusted for baseline value

* p<0.001

Effects from dapagliflozin on renal and cardiovascular outcomes in patients with chronic kidney disease


The Study to Evaluate the Effect of Dapagliflozin on Renal Outcomes and Cardiovascular Mortality in Patients with Chronic Kidney Disease (DAPA-CKD) was an international, multicentre, randomised, double-blind, placebo-controlled study in patients with chronic kidney disease (CKD) with eGFR ≥ 25 to ≤ 75 mL/min/1.73 m² and albuminuria (UACR ≥ 200 and ≤ 5000 mg/g) to determine the effect of dapagliflozin compared with placebo, when added to background standard of care therapy, on the incidence of the composite endpoint of $\geq 50\%$ sustained decline in eGFR, end-stage kidney disease (ESKD) (defined as sustained eGFR < 15 mL/min/1.73 m², chronic dialysis treatment or receiving a renal transplant), cardiovascular or renal death.

Of 4,304 patients, 2,152 were randomised to dapagliflozin 10 mg and 2,152 to placebo and followed for a median of 28.5 months. Treatment was continued if eGFR fell to levels below 25 mL/min/1.73 m² during the study and could be continued in cases when dialysis was needed.

The mean age of the study population was 61.8 years, 66.9% were male. At baseline, mean eGFR was 43.1 mL/min/1.73 m² and median UACR was 949.3 mg/g, 44.1% of patients had eGFR 30 to < 45 mL/min/1.73 m² and 14.5% had eGFR < 30 mL/min/1.73 m². 67.5% of the patients had type 2 diabetes mellitus. Patients were on standard of care (SOC) therapy; 97.0% of patients were treated with an angiotensin-converting enzyme inhibitor (ACEi) or angiotensin receptor blocker (ARB).

The study was stopped early for efficacy prior to the planned analysis based on a recommendation by the independent Data Monitoring Committee. Dapagliflozin was superior to placebo in preventing the primary composite endpoint of $\geq 50\%$ sustained decline in eGFR, reaching end-stage kidney disease, cardiovascular or renal death. Based on the Kaplan-Meier plot for the time to first occurrence of the primary composite endpoint, the treatment effect was evident beginning at 4 months and was maintained through the end of study (Figure 3).


Figure 3: Time to first occurrence of the primary composite endpoint, $\geq 50\%$ sustained decline in eGFR, end-stage kidney disease, cardiovascular or renal death



Patients at risk is the number of patients at risk at the beginning of the period.

All four components of the primary composite endpoint individually contributed to the treatment effect. Dapagliflozin also reduced the incidence of the composite endpoint of $\geq 50\%$ sustained decline in eGFR, end-stage kidney disease or renal death and the composite endpoint of cardiovascular death and hospitalisation for heart failure. Treatment with dapagliflozin improved overall survival in chronic kidney disease patients with a significant reduction in all-cause mortality (Figure 4).

Figure 4: Treatment effects for the primary and secondary composite endpoints, their individual components, and all-cause mortality



The number of first events for the single components are the actual number of first events for each component and does not add up to the number of events in the composite endpoint.

Event rates are presented as the number of subjects with event per 100 patient years of follow-up.

Hazard ratio estimates are not presented for subgroups with less than 15 events in total, both arms combined.

The treatment benefit of dapagliflozin was consistent in chronic kidney disease patients with type 2 diabetes mellitus and without diabetes. Dapagliflozin reduced the primary composite endpoint of $\geq 50\%$ sustained decline in eGFR, reaching end-stage kidney disease, cardiovascular or renal death with a HR of 0.64 (95% CI 0.52, 0.79) in patients with type 2 diabetes mellitus and 0.50 (95% CI 0.35, 0.72) in patients without diabetes.

The treatment benefit of dapagliflozin over placebo on the primary endpoint was also consistent across other key subgroups, including eGFR, age, gender, and region.

Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with [Product name] in all subsets of the paediatric population in the treatment of type 2 diabetes (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

Absorption

Bioequivalence has been demonstrated between [Product name] and the individual sitagliptin and dapagliflozin tablets after single dose administration in the fasted state in healthy subjects.

The following statements reflect the pharmacokinetic properties of the individual active substances of [Product name].

Sitagliptin

Following oral administration of a 100-mg dose to healthy subjects, sitagliptin was rapidly absorbed, with peak plasma concentrations (median T_{max}) occurring 1 to 4 hours post-dose, mean plasma AUC of sitagliptin was 8.52 $\mu\text{M}\cdot\text{hr}$, C_{max} was 950 nM. The absolute bioavailability of sitagliptin is approximately 87 %. Since co-administration of a high-fat meal with sitagliptin had no effect on the pharmacokinetics, sitagliptin may be administered with or without food.

Dapagliflozin

Dapagliflozin was rapidly and well absorbed after oral administration. Maximum dapagliflozin plasma concentrations (C_{max}) were usually attained within 2 hours after administration in the fasted state. Geometric mean steady-state dapagliflozin C_{max} and AUC_{τ} values following once daily 10 mg doses of dapagliflozin were 158 ng/mL and 628 ng h/mL, respectively. The absolute oral bioavailability of dapagliflozin following the administration of a 10 mg dose is 78%. Administration with a high-fat meal decreased dapagliflozin C_{max} by up to 50% and prolonged T_{max} by approximately 1 hour, but did not alter AUC as compared with the fasted state. These changes are not considered to be clinically meaningful. Hence, dapagliflozin can be administered with or without food.

Distribution

Sitagliptin

The mean volume of distribution at steady state following a single 100-mg intravenous dose of sitagliptin to healthy subjects is approximately 198 litres. The fraction of sitagliptin reversibly bound to plasma proteins is low (38 %).

Dapagliflozin

Dapagliflozin is approximately 91% protein bound. Protein binding was not altered in various disease states (e.g. renal or hepatic impairment). The mean steady-state volume of distribution of dapagliflozin was 118 liters.

Biotransformation

Sitagliptin

Sitagliptin is primarily eliminated unchanged in urine, and metabolism is a minor pathway. Approximately 79% of sitagliptin is excreted unchanged in the urine.

Following a [^{14}C]sitagliptin oral dose, approximately 16% of the radioactivity was excreted as metabolites of sitagliptin. Six metabolites were detected at trace levels and are not expected to contribute to the plasma DPP-4 inhibitory activity of sitagliptin. *In vitro* studies indicated that

the primary enzyme responsible for the limited metabolism of sitagliptin was CYP3A4, with contribution from CYP2C8.

In vitro data showed that sitagliptin is not an inhibitor of CYP isozymes CYP3A4, 2C8, 2C9, 2D6, 1A2, 2C19 or 2B6, and is not an inducer of CYP3A4 and CYP1A2.

Dapagliflozin

Dapagliflozin is extensively metabolised, primarily to yield dapagliflozin 3-O-glucuronide, which is an inactive metabolite. Dapagliflozin 3-O-glucuronide or other metabolites do not contribute to the glucose-lowering effects. The formation of dapagliflozin 3-O-glucuronide is mediated by UGT1A9, an enzyme present in the liver and kidney, and CYP-mediated metabolism was a minor clearance pathway in humans.

Elimination

Sitagliptin

Following administration of an oral [¹⁴C]sitagliptin dose to healthy subjects, approximately 100% of the administered radioactivity was eliminated in faeces (13%) or urine (87%) within one week of dosing. The apparent terminal $t_{1/2}$ following a 100-mg oral dose of sitagliptin was approximately 12.4 hours. Sitagliptin accumulates only minimally with multiple doses. The renal clearance was approximately 350 mL/min.

Elimination of sitagliptin occurs primarily via renal excretion and involves active tubular secretion. Sitagliptin is a substrate for human organic anion transporter-3 (hOAT-3), which may be involved in the renal elimination of sitagliptin. The clinical relevance of hOAT-3 in sitagliptin transport has not been established. Sitagliptin is also a substrate of p-glycoprotein, which may also be involved in mediating the renal elimination of sitagliptin. However, ciclosporin, a p-glycoprotein inhibitor, did not reduce the renal clearance of sitagliptin. Sitagliptin is not a substrate for OCT2 or OAT1 or PEPT1/2 transporters. *In vitro*, sitagliptin did not inhibit OAT3 (IC₅₀=160 µM) or p-glycoprotein (up to 250 µM) mediated transport at therapeutically relevant plasma concentrations. In a clinical study sitagliptin had a small effect on plasma digoxin concentrations indicating that sitagliptin may be a mild inhibitor of p-glycoprotein.

Dapagliflozin

The mean plasma terminal half-life ($t_{1/2}$) for dapagliflozin was 12.9 hours following a single oral dose of dapagliflozin 10 mg to healthy subjects. The mean total systemic clearance of dapagliflozin administered intravenously was 207 mL/min. Dapagliflozin and related metabolites are primarily eliminated via urinary excretion with less than 2% as unchanged dapagliflozin. After administration of a 50 mg [¹⁴C]-dapagliflozin dose, 96% was recovered, 75% in urine and 21% in faeces. In faeces, approximately 15% of the dose was excreted as parent drug.

Linearity

Sitagliptin

Plasma AUC of sitagliptin increased in a dose-proportional manner. Dose-proportionality was not established for C_{max} and C_{24hr} (C_{max} increased in a greater than dose-proportional manner and C_{24hr} increased in a less than dose-proportional manner).

Dapagliflozin

Dapagliflozin exposure increased proportional to the increment in dapagliflozin dose over the range of 0.1 to 500 mg and its pharmacokinetics did not change with time upon repeated daily dosing for up to 24 weeks.

Special populations

Sitagliptin

The pharmacokinetics of sitagliptin were generally similar in healthy subjects and in patients with type 2 diabetes.

Renal impairment

Sitagliptin

A single-dose, open-label study was conducted to evaluate the pharmacokinetics of a reduced dose of sitagliptin (50 mg) in patients with varying degrees of chronic renal impairment compared to normal healthy control subjects. The study included patients with mild, moderate, and severe renal impairment, as well as patients with ESRD on haemodialysis. In addition, the effects of renal impairment on sitagliptin pharmacokinetics in patients with type 2 diabetes and mild, moderate, or severe renal impairment (including ESRD) were assessed using population pharmacokinetic analyses.

Compared to normal healthy control subjects, plasma AUC of sitagliptin was increased by approximately 1.2-fold and 1.6-fold in patients with mild renal impairment ($\text{GFR} \geq 60$ to < 90 mL/min) and patients with moderate renal impairment ($\text{GFR} \geq 45$ to < 60 mL/min), respectively. Because increases of this magnitude are not clinically relevant, dosage adjustment in these patients is not necessary.

Plasma AUC of sitagliptin was increased approximately 2-fold in patients with moderate renal impairment ($\text{GFR} \geq 30$ to < 45 mL/min), and approximately 4-fold in patients with severe renal impairment ($\text{GFR} < 30$ mL/min), including in patients with ESRD on haemodialysis. Sitagliptin was modestly removed by haemodialysis (13.5 % over a 3- to 4-hour haemodialysis session starting 4 hours postdose). To achieve plasma concentrations of sitagliptin similar to those in patients with normal renal function, lower dosages are recommended in patients with $\text{GFR} < 45$ mL/min (see section 4.2).

Dapagliflozin

At steady-state (20 mg once-daily dapagliflozin for 7 days), subjects with type 2 diabetes mellitus and mild, moderate or severe renal impairment (as determined by iohexol plasma clearance) had mean systemic exposures of dapagliflozin of 32%, 60% and 87% higher, respectively, than those of subjects with type 2 diabetes mellitus and normal renal function. The steady-state 24-hour urinary glucose excretion was highly dependent on renal function and 85, 52, 18 and 11 g of glucose/day was excreted by subjects with type 2 diabetes mellitus and normal renal function or mild, moderate or severe renal impairment, respectively. The impact of haemodialysis on dapagliflozin exposure is not known. The effect of reduced renal function on systemic exposure was evaluated in a population pharmacokinetic model. Consistent with previous results, model predicted AUC was higher in patients with chronic kidney disease compared with patients with normal renal function, and was not meaningfully different in chronic kidney disease patients with type 2 diabetes mellitus and without diabetes.

Hepatic impairment

Sitagliptin

No dose adjustment for sitagliptin is necessary for patients with mild or moderate hepatic impairment (Child-Pugh score ≤ 9). There is no clinical experience in patients with severe hepatic impairment (Child-Pugh score > 9). However, because sitagliptin is primarily renally eliminated, severe hepatic impairment is not expected to affect the pharmacokinetics of sitagliptin.

Dapagliflozin

In subjects with mild or moderate hepatic impairment (Child-Pugh classes A and B), mean $C_{\max}$ and AUC of dapagliflozin were up to 12% and 36% higher, respectively, compared to healthy matched control subjects. These differences were not considered to be clinically meaningful. In subjects with severe hepatic impairment (Child-Pugh class C) mean $C_{\max}$ and AUC of dapagliflozin were 40% and 67% higher than matched healthy controls, respectively.

Elderly

Sitagliptin

No dose adjustment is required based on age. Age did not have a clinically meaningful impact on the pharmacokinetics of sitagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data. Elderly subjects (65 to 80 years) had approximately 19 % higher plasma concentrations of sitagliptin compared to younger subjects.

Dapagliflozin

There is no clinically meaningful increase in exposure based on age alone in subjects up to 70 years old. However, an increased exposure due to age-related decrease in renal function can be expected. There are insufficient data to draw conclusions regarding exposure in patients >70 years old.

Other patient characteristics

Sitagliptin

No dose adjustment is necessary based on gender, race, or body mass index (BMI). These characteristics had no clinically meaningful effect on the pharmacokinetics of sitagliptin based on a composite analysis of Phase I pharmacokinetic data and on a population pharmacokinetic analysis of Phase I and Phase II data.

Dapagliflozin

The mean dapagliflozin AUC_{ss} in females was estimated to be about 22% higher than in males.

There were no clinically relevant differences in systemic exposures between White, Black or Asian races.

Dapagliflozin exposure was found to decrease with increased weight. Consequently, low-weight patients may have somewhat increased exposure and patients with high weight somewhat decreased exposure. However, the differences in exposure were not considered clinically meaningful.

5.3 Preclinical safety data

The following data are findings in studies performed with sitagliptin or dapagliflozin individually.

Sitagliptin

Renal and liver toxicity were observed in rodents at systemic exposure values 58 times the human exposure level, while the no-effect level was found at 19 times the human exposure level. Incisor teeth abnormalities were observed in rats at exposure levels 67 times the clinical exposure level; the no-effect level for this finding was 58-fold based on the 14-week rat study. The relevance of these findings for humans is unknown. Transient treatment-related physical signs, some of which suggest neural toxicity, such as open-mouth breathing, salivation, white foamy emesis, ataxia, trembling, decreased activity, and/or hunched posture were observed in dogs at exposure levels approximately 23 times the clinical exposure level. In addition, very slight to slight skeletal muscle degeneration was also observed histologically at doses resulting in systemic exposure levels of approximately 23 times the human exposure

level. A no-effect level for these findings was found at an exposure 6-fold the clinical exposure level.

Sitagliptin has not been demonstrated to be genotoxic in preclinical studies. Sitagliptin was not carcinogenic in mice. In rats, there was an increased incidence of hepatic adenomas and carcinomas at systemic exposure levels 58 times the human exposure level. Since hepatotoxicity has been shown to correlate with induction of hepatic neoplasia in rats, this increased incidence of hepatic tumours in rats was likely secondary to chronic hepatic toxicity at this high dose. Because of the high safety margin (19-fold at this no-effect level), these neoplastic changes are not considered relevant for the situation in humans.

No adverse effects upon fertility were observed in male and female rats given sitagliptin prior to and throughout mating.

In a pre-/postnatal development study performed in rats sitagliptin showed no adverse effects. Reproductive toxicity studies showed a slight treatment-related increased incidence of foetal rib malformations (absent, hypoplastic and wavy ribs) in the offspring of rats at systemic exposure levels more than 29 times the human exposure levels. Maternal toxicity was seen in rabbits at more than 29 times the human exposure levels. Because of the high safety margins, these findings do not suggest a relevant risk for human reproduction. Sitagliptin is secreted in considerable amounts into the milk of lactating rats (milk/plasma ratio: 4:1).

Dapagliflozin

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential and fertility. Dapagliflozin did not induce tumours in either mice or rats at any of the doses evaluated in two-year carcinogenicity studies.

Direct administration of dapagliflozin to weanling juvenile rats and indirect exposure during late pregnancy (time periods corresponding to the second and third trimesters of pregnancy with respect to human renal maturation) and lactation are each associated with increased incidence and/or severity of renal pelvic and tubular dilatations in progeny.

In a juvenile toxicity study, when dapagliflozin was dosed directly to young rats from postnatal day 21 until postnatal day 90, renal pelvic and tubular dilatations were reported at all dose levels; pup exposures at the lowest dose tested were ≥ 15 times the maximum recommended human dose. These findings were associated with dose-related increases in kidney weight and macroscopic kidney enlargement observed at all doses. The renal pelvic and tubular dilatations observed in juvenile animals did not fully reverse within the approximate 1-month recovery period.

In a separate study of pre- and postnatal development, maternal rats were dosed from gestation day 6 through postnatal day 21, and pups were indirectly exposed in utero and throughout lactation. (A satellite study was conducted to assess dapagliflozin exposures in milk and pups.) Increased incidence or severity of renal pelvic dilatation was observed in adult offspring of treated dams, although only at the highest dose tested (associated maternal and pup dapagliflozin exposures were 1,415 times and 137 times, respectively, the human values at the maximum recommended human dose). Additional developmental toxicity was limited to dose-related reductions in pup body weights, and observed only at doses ≥ 15 mg/kg/day (associated with pup exposures that are ≥ 29 times the human values at the maximum recommended human dose). Maternal toxicity was evident only at the highest dose tested, and limited to transient reductions in body weight and food consumption at dose. The no observed adverse effect level (NOAEL) for developmental toxicity, the lowest dose tested, is associated with a maternal systemic exposure multiple that is approximately 19 times the human value at the maximum recommended human dose.

In additional studies of embryo-foetal development in rats and rabbits, dapagliflozin was administered for intervals coinciding with the major periods of organogenesis in each species. Neither maternal nor developmental toxicities were observed in rabbits at any dose tested; the highest dose tested is associated with a systemic exposure multiple of approximately 1,191 times the maximum recommended human dose. In rats, dapagliflozin was neither embryo-lethal nor teratogenic at exposures up to 1,441 times the maximum recommended human dose.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Cellulose, Microcrystalline

Lactose monohydrate

Sodium starch glycolate type A

Magnesium stearate

Film-coating

Polyvinyl alcohol, part. hydrolysed

Iron oxide yellow

Macrogol

Talc

Iron oxide red

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

18 months

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

OPA/Alu/PVC-Alu perforated or non-perforated blisters containing 28, 30, 60 or 98 film-coated tablets. Each blister strip contains 7 or 10 film-coated tablets. The blisters are packed in cartons.

OPA/Alu/PVC-Alu perforated unit-dose blisters containing 28x1 or 98x1 film-coated tablets. Each blister strip contains 7 film-coated tablets. The blisters are packed in cartons.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

<To be completed nationally>

8. MARKETING AUTHORISATION NUMBER(S)

<To be completed nationally>

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE
AUTHORISATION**

<To be completed nationally>

10. DATE OF REVISION OF THE TEXT

19 Nov 2025

<To be completed nationally>

Detailed information on this medicinal product is available on the website of: {name of
Member State/Agency}

LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

OUTER CARTON EU full/**reduced** harmonised labelling text
Baltic cluster (LT/LV/EE)

1. NAME OF THE MEDICINAL PRODUCT

Sitagliptin/Dapagliflozin Teva 100 mg/10 mg film-coated tablets

sitagliptin/dapagliflozin

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each **film-coated** tablet contains 100 mg sitagliptin (as sitagliptin hydrochloride monohydrate) and 10 mg dapagliflozin (as dapagliflozin amorphous base).

3. LIST OF EXCIPIENTS

Contains lactose. **See package leaflet for further information.**

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

Film-coated tablet

[Product name] 100 mg/10 mg

28 **film-coated** tablets

30 **film-coated** tablets

60 **film-coated** tablets

98 **film-coated** tablets

28 ×1 **film-coated** tablets

98 ×1 **film-coated** tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.

Oral use.

<QR code to be included> + URL

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

<[To be completed nationally]>

12. MARKETING AUTHORISATION NUMBER(S)

<[To be completed nationally]>

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

<[To be completed nationally]>

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

[Product name] 100 mg/10 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTER EU full/*reduced* harmonised labelling text
Baltic cluster (LT/LV/EE)

1. NAME OF THE MEDICINAL PRODUCT

Sitagliptin/Dapagliflozin Teva 100 mg/10 mg *film-coated* tablets

sitagliptin/dapagliflozin

2. NAME OF THE MARKETING AUTHORISATION HOLDER

<[To be completed nationally]>

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PACKAGE LEAFLET

Package leaflet: Information for the patient

Sitagliptin/Dapagliflozin Teva 100 mg/10 mg film-coated tablets sitagliptin/dapagliflozin

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [Product name] is and what it is used for
2. What you need to know before you take [Product name]
3. How to take [Product name]
4. Possible side effects
5. How to store [Product name]
6. Contents of the pack and other information

1. What [Product name] is and what it is used for

What [Product name] is:

[Product name] contains the active substances sitagliptin and dapagliflozin.

Sitagliptin is a member of a class of medicines called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors) that lower blood sugar levels in adult patients with type 2 diabetes mellitus. Dapagliflozin belongs to a group of medicines called “sodium glucose co-transporter-2 (SGLT2) inhibitors”. They work by blocking the SGLT2 protein in your kidney. By blocking this protein, blood sugar (glucose), salt (sodium) and water are removed from your body via the urine.

[Product name] is used for type 2 diabetes.

- in adults aged 18 years and older
- if your type 2 diabetes cannot be controlled with diet and exercise
- if metformin and sitagliptin cannot control your diabetes,
- you are already being treated with sitagliptin and dapagliflozin as single tablets. Your doctor may ask you to switch to this medicine

Diet and exercise can help your body use its blood sugar better. If you have diabetes, it is important to stay on any diet and exercise program recommended by your doctor while taking [Product name].

What is type 2 diabetes and how does [Product name] help?

In type 2 diabetes your body does not make enough insulin or is not able to use the insulin it makes properly. This leads to a high level of sugar in your blood. This can lead to serious problems like heart or kidney disease, blindness, and poor circulation in your arms and legs.

This medicine helps to increase the levels of insulin produced after a meal, decreases the amount of sugar made by the body and removes excess sugar from your body.

Your doctor has prescribed this medicine to help lower your blood sugar, which is too high because of your type 2 diabetes.

2. What you need to know before you take [Product name]

Do not take [Product name]

- if you are allergic to sitagliptin, dapagliflozin or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking [Product name] :

- if you have “type 1 diabetes” – the type that usually starts when you are young, and your body does not produce any insulin. [Product name] should not be used to treat this condition.
- if you have or had a disease of the pancreas (such as pancreatitis), or if you have or had gallstones, alcohol dependence or very high levels of triglycerides (a form of fat) in your blood. These medical conditions can increase your chance of getting pancreatitis. Possible signs of pancreatitis are listed in section 4.
- if you have or had a kidney problem
- if you have or had a liver problem
- if you have or had very high levels of sugar in your blood which may make you dehydrated (lose too much body fluid). Possible signs of dehydration are listed in section 4. Tell your doctor before you start taking [Product name] if you have any of these signs.
- if you have or had or develop nausea (feeling sick), vomiting or fever or if you are not able to eat or drink. These conditions can cause dehydration. Your doctor may ask you to stop taking [Product name] until you recover to prevent dehydration.
- if you have or had diabetic ketoacidosis (a complication of diabetes with high blood sugar, rapid weight loss, nausea or vomiting). Possible signs of diabetic ketoacidosis are listed in section 4.
- if you often get infections of the urinary tract

If any of the above applies to you (or you are not sure), talk to your doctor, pharmacist or nurse before taking [Product name].

Diabetes and foot care

If you have diabetes, it is important to check your feet regularly and adhere to any other advice regarding foot care given by your health care professional.

Urine glucose

Because of how [Product name] works, your urine will test positive for sugar while you are on this medicine.

Children and adolescents

Do not give this medicine to children under 18 years of age, because [Product name] has not been studied in children.

Other medicines and [Product name]

Tell your doctor, pharmacist or nurse if you are taking, have recently taken, or might take any other medicines.

Especially, tell your doctor if you are taking any of the following medicines:

- Medicines used to treat irregular heartbeat and other heart problems (digoxin). The level of digoxin in your blood may need to be checked if taking with [Product name].
- Medicines used to remove water from the body (diuretic).
- Other medicines that lower the amount of sugar in your blood such as insulin or a “sulphonylurea” medicine. Your doctor may want to lower the dose of these other medicines, to prevent you from getting low blood sugar levels (hypoglycaemia).
- medicines to lower your blood pressure (anti-hypertensives) and have a history of low blood pressure (hypotension).
- Lithium, used for mental disorders [Product name] can lower the amount of lithium in your blood.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You should not take this medicine during pregnancy. You should stop taking this medicine if you become pregnant.

You should not take this medicine if you are breast-feeding or plan to breast-feed. It is not known if this medicine passes into breast milk.

Driving and using machines

This medicine has no or negligible influence on the ability to drive and use machines. However, dizziness and drowsiness have been reported, which may affect your ability to drive or use machines.

Do not drive or use any tools or machines, if you feel dizzy taking [Product name].

Taking this medicine with other medicines called sulphonylureas or with insulin can cause too low blood sugar levels (hypoglycaemia), which may cause symptoms such as shaking, sweating and change in vision, and may affect your ability to drive and use machines.

[Product name] contains lactose and sodium

[Product name] contains lactose (milk sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

[Product name] contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially “sodium-free”.

3. How to take [Product name]

Always take this medicine exactly as your doctor has told you. Check with your doctor, pharmacist or nurse if you are not sure.

The usual recommended dose is:

- one [Product name] film-coated tablet. Your doctor will prescribe the strength that is right for you.
- once a day
- swallow the tablet whole with a glass of water, to ensure accurate dosing and avoid unpleasant taste
- you can take your tablet with or without food

Your doctor may prescribe [Product name] alone or with certain other medicines that lower blood sugar. Remember to take these other medicine(s) as your doctor has told you. This will

help get the best results for your health.

If you take more [Product name] than you should

If you take more [Product name] tablets than you should, talk to a doctor or go to a hospital immediately.

Take the medicine pack with you.

If you forget to take [Product name]

What to do if you forget to take a tablet.

- If it is 12 hours or more until your next dose, take a dose of [Product name] as soon as you remember.
- Then take your next dose at the usual time.
- If it is less than 12 hours until your next dose, skip the missed dose. Then take your next dose at the usual time.
- Do not take a double dose of [Product name] to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking [Product name] and contact a doctor or the nearest hospital immediately if you notice any of the following serious side effects:

- **Pancreatitis** - severe and persistent pain in the abdomen (stomach area) which might reach through to your back with or without nausea and vomiting, as these could be signs of an inflamed pancreas. (frequency not known)
- **Serious allergic reaction (anaphylactic reaction, angioedema)** (very rare). These are signs:
 - swelling of the face, tongue, lips or throat
 - difficulties swallowing and/or breathing
 - hives and breathing problems
 - rash
 - blisters on the skin/peeling skin
- **diabetic ketoacidosis** - this is rare in patients with type 2 diabetes (may affect up to 1 in 1 000 people). These are the signs:
 - increased levels of “ketone bodies” in your urine or blood
 - feeling sick or being sick
 - stomach pain
 - excessive thirst
 - fast and deep breathing
 - confusion
 - unusual sleepiness or tiredness
 - a sweet smell to your breath, a sweet or metallic taste in your mouth or a different odour to your urine or sweat
 - rapid weight loss.

This may occur regardless of blood sugar level. Your doctor may decide to temporarily or permanently stop your treatment with [Product name].

- **necrotising fasciitis of the perineum** or Fournier’s gangrene, a serious or even life threatening soft tissue infection of the genitals or the area between the genitals and the anus, seen very rarely. These are the signs: a combination of symptoms of pain, tenderness, redness, or swelling of the genitals or the area between the genitals and

the anus with fever or feeling generally unwell. Fournier's gangrene has to be treated immediately (very rare).

- **Stevens-Johnson syndrome**, an exfoliative skin condition that usually starts with flu-like symptoms, followed by a painful red or purple rash that spreads and forms blisters on your skin (frequency not known).
- **Bullous pemphigoid**, a type of skin blistering characterized by large, fluid-filled blisters (frequency not known).

Stop taking [Product name] and see a doctor as soon as possible if you notice any of the following serious side effects:

- **urinary tract infection**, seen commonly (may affect up to 1 in 10 people). These are signs of a severe infection of the urinary tract:
 - fever and/or chills
 - burning sensation when passing water (urinating)
 - pain in your back or side.

Although uncommon, if you see blood in your urine, tell your doctor immediately.

Contact your doctor as soon as possible if you have any of the following side effects:

- **low blood sugar levels** (hypoglycaemia), seen very commonly (may affect more than 1 in 10 people) in patients with diabetes taking this medicine with a sulphonylurea or insulin. These are the signs of low blood sugar:
 - shaking, sweating, feeling very anxious, fast heart beat
 - feeling hungry, headache, change in vision
 - a change in your mood or feeling confused.

Your doctor will tell you how to treat low blood sugar levels and what to do if you get any of the signs above.

Other side effects when taking [Product name]:

Common (may affect up to 1 in 10 people)

- genital infection (thrush) of your penis or vagina (signs may include irritation, itching, unusual discharge or odour)
- back pain
- passing more water (urine) than usual or needing to pass water more often
- changes in the amount of cholesterol or fats in your blood (shown in tests)
- increases in the amount of red blood cells in your blood (shown in tests)
- decreases in creatinine renal clearance (shown in tests) in the beginning of treatment
- dizziness
- rash
- headache

Uncommon (may affect up to 1 in 100 people)

- loss of too much fluid from your body (dehydration, signs may include very dry or sticky mouth, passing little or no urine or fast heartbeat)
- thirst
- constipation
- awakening from sleep at night to pass urine
- dry mouth
- weight decreased
- increases in creatinine (shown in laboratory blood tests) in the beginning of treatment
- increases in urea (shown in laboratory blood tests)
- Fungal infection
- Itching, vulvovaginal itching, genital itching

Rare (may affect up to 1 in 1 000 people)

- Low Blood platelets (Thrombocytopenia)

Very rare (may affect up to 1 in 10 000 people)

- Inflammation of part of the kidneys

Frequency not known

- Scarring of the lungs (Interstitial lung disease)
- Vomiting
- Hives, itchy skin rash (urticaria)
- Inflammation of the small blood vessels in the skin (cutaneous vasculitis)
- Joint pain
- Muscle pain
- Joint disease or joint problems

Kidney problems (sometimes requiring dialysis)

Reporting of side effects

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

[To be completed nationally]

5. How to store [Product name]

Keep this medicine out of sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the blister after “EXP.” The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [Product name] contains

The active substances are sitagliptin and dapagliflozin.

- [Product name] 100 mg/10 mg film-coated tablets: each film-coated tablet contains 100 mg sitagliptin (as sitagliptin hydrochloride monohydrate) and 10 mg dapagliflozin (as dapagliflozin amorphous base).

The other ingredients in the tablet core are cellulose, microcrystalline, lactose monohydrate, sodium starch glycolate type A, magnesium stearate.

The film-coating contains poly (vinyl) alcohol– part. hydrolyzed, iron oxide yellow, macrogol, talc, iron oxide red.

What [Product name] looks like and contents of the pack

[Product name] 100 mg/10 mg film-coated tablet is a light/brown, round film-coated tablet with “101” engraved on one side and with diameter 9.9 mm.

Pack sizes

Cartons of 28, 30, 60, 98 film-coated tablets

Cartons of 28 ×1, 98 ×1 film-coated tablets in unit-dose blisters.

Not all pack sizes may be marketed .

Marketing Authorisation Holder

Marketing Authorisation Holder:

<[To be completed nationally]>

Manufacturer

<[To be completed nationally]>

This medicinal product is authorised in the Member States of the European Economic Area under the following names:

<[To be completed nationally]>

This leaflet was last revised in <{MM/YYYY}> <{month YYYY}>.

19 Nov 2025

<[To be completed nationally]>

<Other sources of information>

<Latest approved information on this medicine is available by scanning the QR code included in the <package leaflet> <outer carton> with a smartphone/device. The same information is also available on the following URL: [URL to be included] <and the <NCA> website>>

In case that only the URL is mentioned in the PI and not linked via mobile technology, please include: <Latest approved information on this product is available on the following URL: [URL to be included] <and the <NCA> website>>