

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Vaftor 150 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 150 mg of ivacaftor.

Excipient with known effect

Each film-coated tablet contains 167.2 mg of mannitol.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet

Light blue, capsule-shaped film-coated tablets, “150” embossed in one side and plain on the other, with dimensions 16.5 x 8.5 mm.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

<Product name> is indicated as monotherapy for the treatment of adults, adolescents, and children aged 6 years and older and weighing 25 kg or more with cystic fibrosis (CF) who have an *R117H* *CFTR* mutation or one of the following gating (class III) mutations in the cystic fibrosis transmembrane conductance regulator (*CFTR*) gene: *G551D*, *G1244E*, *G1349D*, *G178R*, *G551S*, *S1251N*, *S1255P*, *S549N* or *S549R* (see sections 4.4 and 5.1).

4.2 Posology and method of administration

<Product name> should only be prescribed by physicians with experience in the treatment of cystic fibrosis. If the patient's genotype is unknown, an accurate and validated genotyping method should be performed before starting treatment to confirm the presence of an indicated mutation in the *CFTR* gene (see section 4.1). The phase of the poly-T variant identified with the *R117H* mutation should be determined in accordance with local clinical recommendations.

Posology

Adults, adolescents, and children aged 6 years and older should be dosed according to Table 1.

Table 1: Dosing recommendations

	Morning	Evening
Ivacaftor as monotherapy		
6 years and older, ≥ 25 kg	One ivacaftor 150 mg tablet	One ivacaftor 150 mg tablet

The morning and evening dose should be taken approximately 12 hours apart with fat-containing food (see Method of administration).

Missed dose

If 6 hours or less have passed since the missed morning or evening dose, the patient should be advised to take it as soon as possible and then take the next dose at the regularly scheduled time. If more than 6 hours have passed since the time the dose is usually taken, the patient should be advised to wait until the next scheduled dose.

Concomitant use of CYP3A inhibitors

When co-administered with moderate or strong inhibitors of CYP3A, the dose should be reduced (see Table 2 for the recommended dose). Dosing intervals should be modified according to clinical response and tolerability (see sections 4.4 and 4.5).

Table 2: Dosing recommendations for concomitant use with moderate or strong CYP3A inhibitors

	Moderate CYP3A inhibitors	Strong CYP3A inhibitors
Ivacaftor as monotherapy		
6 years and older, ≥ 25 kg	One morning tablet of ivacaftor 150 mg once daily. No evening dose.	One morning tablet of ivacaftor 150 mg twice a week, approximately 3 to 4 days apart. No evening dose.

Special populations

Elderly

Very limited data are available for elderly patients treated with ivacaftor administered as monotherapy. No dose adjustment specific to this patient population is required (see section 5.2).

Renal impairment

No dose adjustment is necessary for patients with mild to moderate renal impairment. Caution is recommended in patients with severe renal impairment (creatinine clearance less than or equal to 30 mL/min) or end-stage renal disease (see sections 4.4 and 5.2).

Hepatic impairment

No dose adjustment is necessary in patients with mild hepatic impairment (Child-Pugh Class A).

In patients with moderate hepatic impairment (Child-Pugh Class B) or severe hepatic impairment (Child-Pugh Class C), the ivacaftor dose should be adjusted as detailed in Table 3 (see sections 4.4, 4.8, and 5.2).

Table 3: Dosing recommendations for patients with moderate or severe hepatic impairment

	Moderate (Child-Pugh Class B)	Severe (Child-Pugh Class C)
Ivacaftor as monotherapy		
6 years and older, ≥ 25 kg	One morning tablet of ivacaftor 150 mg once daily. No evening dose.	Use is not recommended unless the benefits are expected to outweigh the risks. If used: one morning tablet of ivacaftor 150 mg every other day or less frequently. Dosing interval should be modified according to clinical response and tolerability. No evening dose.

Paediatric population

The safety and efficacy of ivacaftor as monotherapy have not been established in children less than 1 month of age or in children less than 6 months of age born prematurely (less than 37 weeks of gestational age). No data are available.

Limited data are available in patients less than 6 years of age with an R117H mutation in the CFTR gene. Available data in patients aged 6 years and older are described in sections 4.8, 5.1 and 5.2.

Method of administration

For oral use.

Patients should be instructed to swallow the tablets whole. The tablets should not be chewed, crushed, or broken before swallowing because there are no clinical data currently available to support other methods of administration.

Ivacaftor tablets should be taken with fat-containing food.

Food or drink containing grapefruit should be avoided during treatment (see section 4.5).

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Only patients with CF who had a *G551D*, *G1244E*, *G1349D*, *G178R*, *G551S*, *S1251N*, *S1255P*, *S549N*, *S549R* gating (class III), *G970R* or *R117H* mutation in at least one allele of the *CFTR* gene were included in studies 770-102, 770-103, 770-111 and 770-110 (see section 5.1).

In study 770-111, four patients with the *G970R* mutation were included. In three of four patients the change in the sweat chloride test was < 5 mmol/L and this group did not demonstrate a clinically relevant improvement in FEV₁ after 8 weeks of treatment. Clinical efficacy in patients with the *G970R* mutation of the *CFTR* gene could not be established (see section 5.1).

Efficacy results from a phase 2 study in patients with CF who are homozygous for the *F508del* mutation in the *CFTR* gene showed no statistically significant difference in FEV₁ over 16 weeks of ivacaftor treatment compared to placebo (see section 5.1). Therefore, use of ivacaftor as monotherapy in these patients is not recommended.

Less evidence of a positive effect of ivacaftor has been shown for patients with an *R117H-7T* mutation associated with less severe disease in study 770-110 (see section 5.1).

Elevated transaminases and hepatic injury

Moderate transaminase (alanine transaminase [ALT] or aspartate transaminase [AST]) elevations are common in subjects with CF. Transaminase elevations have been observed in some patients treated with ivacaftor as monotherapy. Therefore, assessments of transaminases (ALT and AST) are recommended for all patients prior to initiating ivacaftor, every 3 months during the first year of treatment and annually thereafter. For all patients with a history of liver disease or transaminase elevations, more frequent monitoring of liver function tests should be considered. In the event of significant elevations of transaminases (e.g., patients with ALT or AST $> 5 \times$ the upper limit of normal (ULN)), dosing should be interrupted, and laboratory tests closely followed until the abnormalities resolve. Following resolution of transaminase elevations, the benefits and risks of resuming treatment should be considered (see sections 4.2, 4.8 and 5.2).

Hepatic impairment

Use of ivacaftor is not recommended in patients with severe hepatic impairment unless the benefits are expected to outweigh the risks (see Table 3 and sections 4.2, 4.8 and 5.2).

Depression

Depression (including suicidal ideation and suicide attempt) has been reported in patients while receiving ivacaftor, usually occurring within three months of treatment initiation and in patients with a history of psychiatric disorders. In some cases, symptom improvement was reported after dose reduction or treatment discontinuation. Patients (and caregivers) should be alerted about the need to monitor for depressed mood, suicidal thoughts, or unusual changes in behaviour and to seek medical advice immediately if these symptoms present.

Renal impairment

Caution is recommended while using ivacaftor in patients with severe renal impairment or end-stage renal disease (see sections 4.2 and 5.2).

Patients after organ transplantation

Ivacaftor has not been studied in patients with CF who have undergone organ transplantation. Therefore, use in transplanted patients is not recommended. See section 4.5 for interactions with ciclosporin or tacrolimus.

Interactions with medicinal products

CYP3A inducers

Exposure to ivacaftor is significantly decreased by the concomitant use of CYP3A inducers, potentially resulting in the loss of ivacaftor efficacy; therefore, co-administration of ivacaftor with strong CYP3A inducers is not recommended (see section 4.5).

CYP3A inhibitors

Exposure to ivacaftor is increased when co-administered with strong or moderate CYP3A inhibitors. The dose of ivacaftor must be adjusted when used concomitantly with strong or moderate CYP3A inhibitors (see Table 2 and sections 4.2 and 4.5).

Paediatric population

Cases of non-congenital lens opacities/cataracts without impact on vision have been reported in paediatric patients treated with ivacaftor. Although other risk factors were present in some cases (such as corticosteroid use and exposure to radiation), a possible risk attributable to treatment with ivacaftor cannot be excluded. Baseline

and follow-up ophthalmological examinations are recommended in paediatric patients initiating ivacaftor treatment (see section 5.3).

Sodium content

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Ivacaftor is a substrate of CYP3A4 and CYP3A5. It is a weak inhibitor of CYP3A and P-gp and a potential inhibitor of CYP2C9. *In vitro* studies showed that ivacaftor is not a substrate for P-gp.

Medicinal products affecting the pharmacokinetics of ivacaftor

CYP3A inducers

Co-administration of ivacaftor with rifampicin, a strong CYP3A inducer, decreased ivacaftor exposure (AUC) by 89% and decreased hydroxymethyl ivacaftor (M1) to a lesser extent than ivacaftor.

Co-administration of ivacaftor with strong CYP3A inducers, such as rifampicin, rifabutin, phenobarbital, carbamazepine, phenytoin and St. John's wort (*Hypericum perforatum*), is not recommended (see section 4.4).

No dose adjustment is recommended when ivacaftor is used with moderate or weak CYP3A inducers.

CYP3A inhibitors

Ivacaftor is a sensitive CYP3A substrate. Co-administration with ketoconazole, a strong CYP3A inhibitor, increased ivacaftor exposure (measured as area under the curve [AUC]) by 8.5-fold and increased M1 to a lesser extent than ivacaftor. A reduction of the ivacaftor dose is recommended for co-administration with strong CYP3A inhibitors, such as ketoconazole, itraconazole, posaconazole, voriconazole, telithromycin and clarithromycin (see Table 2 and sections 4.2 and 4.4).

Co-administration with fluconazole, a moderate inhibitor of CYP3A, increased ivacaftor exposure by 3-fold and increased M1 to a lesser extent than ivacaftor. A reduction of the ivacaftor dose is recommended for patients taking concomitant moderate CYP3A inhibitors, such as fluconazole, erythromycin, and verapamil (see Table 2 and sections 4.2 and 4.4).

Co-administration of ivacaftor with grapefruit juice, which contains one or more components that moderately inhibit CYP3A, may increase exposure to ivacaftor. Food or drink containing grapefruit should be avoided during treatment with ivacaftor (see section 4.2).

Potential for ivacaftor to interact with transporters

In vitro studies showed that ivacaftor is not a substrate for OATP1B1 or OATP1B3. Ivacaftor and its metabolites are substrates of BCRP *in vitro*. Due to its high intrinsic permeability and low likelihood of being excreted intact, co-administration of BCRP inhibitors is not expected to alter exposure of ivacaftor and M1-IVA, while any potential changes in M6-IVA exposures are not expected to be clinically relevant.

Ciprofloxacin

Co-administration of ciprofloxacin with ivacaftor did not affect the exposure of ivacaftor. No dose adjustment is required when ivacaftor is co-administered with ciprofloxacin.

Medicinal products affected by ivacaftor

Administration of ivacaftor may increase systemic exposure of medicinal products that are sensitive substrates of CYP2C9, and/or P-gp, and/or CYP3A which may increase or prolong their therapeutic effect and adverse reactions.

CYP2C9 substrates

Ivacaftor may inhibit CYP2C9. Therefore, monitoring of the international normalised ratio (INR) is recommended during co-administration of warfarin with ivacaftor. Other medicinal products for which exposure may be increased include glimepiride and glipizide; these medicinal products should be used with caution.

Digoxin and other P-gp substrates

Co-administration with digoxin, a sensitive P-gp substrate, increased digoxin exposure by 1.3-fold, consistent with weak inhibition of P-gp by ivacaftor. Administration of ivacaftor may increase systemic exposure of medicinal products that are sensitive substrates of P-gp, which may increase or prolong their therapeutic effect and adverse reactions. When used concomitantly with digoxin or other substrates of P-gp with a narrow therapeutic index, such as ciclosporin, everolimus, sirolimus or tacrolimus, caution and appropriate monitoring should be used.

CYP3A substrates

Co-administration with (oral) midazolam, a sensitive CYP3A substrate, increased midazolam exposure 1.5-fold, consistent with weak inhibition of CYP3A by ivacaftor. No dose adjustment of CYP3A substrates, such as midazolam, alprazolam, diazepam or triazolam, is required when these are co-administered with ivacaftor.

Hormonal contraceptives

Ivacaftor has been studied with an oestrogen/progesterone oral contraceptive and was found to have no significant effect on the exposures of the oral contraceptive. Therefore, no dose adjustment of oral contraceptives is necessary.

Paediatric population

Interaction studies have only been performed in adults.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no or limited amount of data (less than 300 pregnancy outcomes) from the use of ivacaftor in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). As a precautionary measure, it is preferable to avoid the use of ivacaftor during pregnancy.

Breast-feeding

Limited data show that ivacaftor is excreted into human milk. A risk to the newborns/infants cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from ivacaftor therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility

There are no data available on the effect of ivacaftor on fertility in humans. Ivacaftor had an effect on fertility in rats (see section 5.3).

4.7 Effects on ability to drive and use machines

Ivacaftor has minor influence on the ability to drive and use machines. Ivacaftor may cause dizziness (see section 4.8) and, therefore, patients experiencing dizziness should be advised not to drive or use machines until symptoms abate.

4.8 Undesirable effects

Summary of the safety profile

The most common adverse reactions experienced by patients aged 6 years and older who received ivacaftor are headache (23.9%), oropharyngeal pain (22.0%), upper respiratory tract infection (22.0%), nasal congestion (20.2%), abdominal pain (15.6%), nasopharyngitis (14.7%), diarrhoea (12.8%), dizziness (9.2%), rash (12.8%) and bacteria in sputum (12.8%). Transaminase elevations occurred in 12.8% of ivacaftor-treated patients versus 11.5% of placebo-treated patients.

Serious adverse reactions in patients who received ivacaftor included abdominal pain and transaminase elevations (see section 4.4).

Tabulated list of adverse reactions

Table 4 reflects the adverse reactions observed with ivacaftor monotherapy in clinical trials (placebo- controlled and uncontrolled studies) in which the length of exposure to ivacaftor ranged from 16 weeks to 144 weeks. The frequency of adverse reactions is defined as follows: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 4: Adverse reactions

System organ class	Adverse reactions	Frequency
Infections and infestations	Upper respiratory tract infection	very common
	Nasopharyngitis	very common
	Rhinitis	common
Nervous system disorders	Headache	very common
	Dizziness	very common
Psychiatric disorders	Depression	not known
Ear and labyrinth disorders	Ear pain	common
	Ear discomfort	common
	Tinnitus	common
	Tympanic membrane hyperaemia	common
	Vestibular disorder	common
	Ear congestion	uncommon
Respiratory, thoracic and mediastinal disorders	Oropharyngeal pain	very common
	Nasal congestion	very common
	Sinus congestion	common
	Pharyngeal erythema	common
Gastrointestinal disorders	Abdominal pain	very common
	Diarrhoea	very common
Hepatobiliary disorders	Transaminase elevations	very common
Skin and subcutaneous tissue	Rash	very common
Reproductive system and breast disorders	Breast mass	common
	Breast inflammation	uncommon
	Gynaecomastia	uncommon
	Nipple disorder	uncommon
	Nipple pain	uncommon
Investigations	Bacteria in sputum	very common

Description of selected adverse reactions

Transaminase elevations

During the 48-week placebo-controlled studies 770-102 and 770-103 of ivacaftor as monotherapy in patients aged 6 years and older, the incidence of maximum transaminase (ALT or AST) > 8 , > 5 or $> 3 \times \text{ULN}$ was 3.7%, 3.7% and 8.3% in ivacaftor-treated patients and 1.0%, 1.9% and 8.7% in placebo-treated patients, respectively. Two patients, one on placebo and one on ivacaftor permanently discontinued treatment for elevated transaminases, each $> 8 \times \text{ULN}$. No ivacaftor-treated patients experienced a transaminase elevation $> 3 \times \text{ULN}$ associated with elevated total bilirubin $> 1.5 \times \text{ULN}$. In ivacaftor-treated patients, most transaminase elevations up to $5 \times \text{ULN}$ resolved without treatment interruption. Ivacaftor dosing was interrupted in most patients with transaminase elevations $> 5 \times \text{ULN}$. In all instances where dosing was interrupted for elevated transaminases and subsequently resumed, ivacaftor dosing was able to be resumed successfully (see section 4.4).

Post-marketing cases of treatment discontinuation due to elevated transaminases have been reported (see section 4.4).

Paediatric population

Safety of ivacaftor as monotherapy for 24 weeks was evaluated in 43 patients between 1 month to less than 24 months of age (with 7 of them less than 4 months old), 34 patients between 2 to less than 6 years of age, 61 patients between 6 to less than 12 years of age and 94 patients between 12 to less than 18 years of age.

The safety profile of ivacaftor is generally consistent among paediatric patients and is also consistent with adult patients.

The incidence of transaminase elevations (ALT or AST) observed in studies 770-103, 770-111 and 770-110 (patients aged 6 to less than 12 years) is described in Table 5. In the placebo-controlled studies, the incidence of transaminase elevations was similar between treatment with ivacaftor (15.0%) and placebo (14.6%). Peak LFT elevations were generally higher in paediatric patients than in older patients. Across all populations, peak LFT elevations returned to baseline levels following interruption, and in almost all instances where dosing was interrupted for elevated transaminases and subsequently resumed, ivacaftor dosing was able to be resumed successfully (see section 4.4). Cases suggestive of positive rechallenge were observed.

Table 5: Transaminase elevations in patients aged 1 month to < 12 years treated with ivacaftor as monotherapy

Age group	n	% of Patients $> 3 \times \text{ULN}$	% of Patients $> 5 \times \text{ULN}$	% of Patients $> 8 \times \text{ULN}$
6 to < 12 years	40	15.0% (6)	2.5% (1)	2.5% (1)
2 to < 6 years	34	14.7% (5)	14.7% (5)	14.7% (5)
12 to < 24 months	18	27.8% (5)	11.1% (2)	11.1% (2)
1 to < 12 months	24	8.3% (2)	4.2% (1)	4.2% (1)

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

No specific antidote is available for overdose with ivacaftor. Treatment of overdose consists of general supportive measures including monitoring of vital signs, liver function tests and observation of the clinical status of the patient.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Other respiratory system products, ATC code: R07AX02 Mechanism

of action

Ivacaftor is a potentiator of the CFTR protein, i.e., *in vitro* ivacaftor increases CFTR channel gating to enhance chloride transport in specified gating mutations (as listed in section 4.1) with reduced channel-open probability compared to normal CFTR. Ivacaftor also potentiated the channel-open probability of R117H-CFTR, which has both low channel-open probability (gating) and reduced channel current amplitude (conductance). The *G970R* mutation causes a splicing defect resulting in little-to-no CFTR protein at the cell surface which may explain the results observed in subjects with this mutation in study 770-111 (see Pharmacodynamic effects and Clinical efficacy and safety).

In vitro responses seen in single channel patch clamp experiments using membrane patches from rodent cells expressing mutant CFTR forms do not necessarily correspond to *in vivo* pharmacodynamic response (e.g., sweat chloride) or clinical benefit. The exact mechanism leading ivacaftor to potentiate the gating activity of normal and some mutant CFTR forms in this system has not been completely elucidated.

Pharmacodynamic effects

Ivacaftor as monotherapy

In studies 770-102 and 770-103 in patients with the *G551D* mutation in one allele of the *CFTR* gene, ivacaftor led to rapid (15 days), substantial (the mean change in sweat chloride from baseline through week 24 was -48 mmol/L [95% CI -51, -45] and -54 mmol/L [95% CI -62, -47], respectively) and sustained (through 48 weeks) reductions in sweat chloride concentration.

In study 770-111, part 1 in patients who had a non-*G551D* gating mutation in the *CFTR* gene, treatment with ivacaftor led to a rapid (15 days) and substantial mean change from baseline in sweat chloride of -49 mmol/L (95% CI -57, -41) through 8 weeks of treatment. However, in patients with the *G970R-CFTR* mutation, the mean (SD) absolute change in sweat chloride at week 8 was -6.25 (6.55) mmol/L. Similar results to part 1 were seen in part 2 of the study. At the 4-week follow-up visit (4 weeks after dosing with ivacaftor ended), mean sweat chloride values for each group were trending to pre-treatment levels.

In study 770-110 in patients aged 6 years or older with CF who had an *R117H* mutation in the *CFTR* gene, the treatment difference in mean change in sweat chloride from baseline through 24 weeks of treatment was -24 mmol/L (95% CI -28, -20). In subgroup analyses by age, the treatment difference was -21.87 mmol/L (95% CI: -26.46, -17.28) in patients aged 18 years or older, and -27.63 mmol/L (95% CI: -37.16, -18.10) in patients aged 6-11 years. Two patients 12 to 17 years of age were enrolled in this study.

Clinical efficacy and safety

Ivacaftor as monotherapy

Studies 770-102 and 770-103: studies in patients with CF with *G551D* gating mutations

The efficacy of ivacaftor has been evaluated in two phase 3 randomised, double-blind, placebo-controlled, multi-centre studies of clinically stable patients with CF who had the *G551D* mutation in the *CFTR* gene on at least 1 allele and had FEV₁ ≥ 40% predicted.

Patients in both studies were randomised 1:1 to receive either 150 mg of ivacaftor or placebo every 12 hours with food containing fat for 48 weeks in addition to their prescribed CF therapies (e.g., tobramycin, dornase alfa). The use of inhaled hypertonic sodium chloride was not permitted.

Study 770-102 evaluated 161 patients who were 12 years of age or older; 122 (75.8%) patients had the *F508del* mutation in the second allele. At the start of the study, patients in the placebo group used some medicinal

products at a higher frequency than the ivacaftor group. These medicinal products included dornase alfa (73.1% versus 65.1%), salbutamol (53.8% versus 42.2%), tobramycin (44.9% versus 33.7%) and salmeterol/fluticasone (41.0% versus 27.7%). At baseline, mean predicted FEV₁ was 63.6% (range: 31.6% to 98.2%) and mean age was 26 years (range: 12 to 53 years).

Study 770-103 evaluated 52 patients who were 6 to 11 years of age at screening; mean (SD) body weight was 30.9 (8.63) kg; 42 (80.8%) patients had the *F508del* mutation in the second allele. At baseline, mean predicted FEV₁ was 84.2% (range: 44.0% to 133.8%) and mean age was 9 years (range: 6 to 12 years); 8 (30.8%) patients in the placebo group and 4 (15.4%) patients in the ivacaftor group had an FEV₁ less than 70% predicted at baseline.

The primary efficacy endpoint in both studies was the mean absolute change from baseline in percent predicted FEV₁ through 24 weeks of treatment.

The treatment difference between ivacaftor and placebo for the mean absolute change (95% CI) in percent predicted FEV₁ from baseline through week 24 was 10.6 percentage points (8.6, 12.6) in study 770-102 and 12.5 percentage points (6.6, 18.3) in study 770-103. The treatment difference between ivacaftor and placebo for the mean relative change (95% CI) in percent predicted FEV₁ from baseline through week 24 was 17.1% (13.9, 20.2) in study 770-102 and 15.8% (8.4, 23.2) in study 770-103. The mean change from baseline through week 24 in FEV₁ (L) was 0.37 L in the ivacaftor group and 0.01 L in the placebo group in study 770-102 and 0.30 L in the ivacaftor group and 0.07 L in the placebo group in study 770-103. In both studies, improvements in FEV₁ were rapid in onset (day 15) and durable through 48 weeks.

The treatment difference between ivacaftor and placebo for the mean absolute change (95% CI) in percent predicted FEV₁ from baseline through week 24 in patients 12 to 17 years of age in study 770-102 was 11.9 percentage points (5.9, 17.9). The treatment difference between ivacaftor and placebo for the mean absolute change (95% CI) in percent predicted FEV₁ from baseline through week 24 in patients with baseline predicted FEV₁ greater than 90% in study 770-103 was 6.9 percentage points (-3.8, 17.6).

The results for clinically relevant secondary endpoints are shown in Table 6.

Table 6: Effect of ivacaftor on other efficacy endpoints in studies 770-102 and 770-103

Endpoint	Study 770-102		Study 770-103	
	Treatment difference ^a (95% CI)	P value	Treatment difference ^a (95% CI)	P value
Mean absolute change from baseline in CFQ-R^b respiratory domain score (points)^c				
Through week 24	8.1 (4.7, 11.4)	< 0.0001	6.1 (-1.4, 13.5)	0.1092
Through week 48	8.6 (5.3, 11.9)	< 0.0001	5.1 (-1.6, 11.8)	0.1354
Relative risk of pulmonary exacerbation				
Through week 24	0.40 ^d	0.0016	NA	NA
Through week 48	0.46 ^d	0.0012	NA	NA
Mean absolute change from baseline in body weight (kg)				
At week 24	2.8 (1.8, 3.7)	< 0.0001	1.9 (0.9, 2.9)	0.0004
At week 48	2.7 (1.3, 4.1)	0.0001	2.8 (1.3, 4.2)	0.0002
Mean absolute change from baseline in BMI (kg/m²)				
At week 24	0.94 (0.62, 1.26)	< 0.0001	0.81 (0.34, 1.28)	0.0008
At week 48	0.93 (0.48, 1.38)	< 0.0001	1.09 (0.51, 1.67)	0.0003
Mean change from baseline in z-scores				
Weight-for-age z-score at week 48 ^e	0.33 (0.04, 0.62)	0.0260	0.39 (0.24, 0.53)	< 0.0001
BMI-for-age z-score at week 48 ^e	0.33 (0.002, 0.65)	0.0490	0.45 (0.26, 0.65)	< 0.0001

CI: confidence interval; NA: not analysed due to low incidence of events

^a Treatment difference = effect of ivacaftor – effect of placebo

^b CFQ-R: Cystic Fibrosis Questionnaire-Revised is a disease-specific, health-related quality-of-life measure for CF.

^c Study 770-102 data were pooled from CFQ-R for adults/adolescents and CFQ-R for children 12 to 13 years of age; Study 770-103 data were obtained from CFQ-R for children 6 to 11 years of age.

^d Hazard ratio for time to first pulmonary exacerbation

^e In subjects under 20 years of age (CDC growth charts)

Study 770-111: study in patients with CF with non-G551D gating mutations

Study 770-111 was a phase 3, two-part, randomised, double-blind, placebo-controlled, crossover study (part 1) followed by a 16-week open-label extension period (part 2) to evaluate the efficacy and safety of ivacaftor in patients with CF aged 6 years and older who have a *G970R* or non-*G551D* gating mutation in the *CFTR* gene (*G178R*, *S549N*, *S549R*, *G551S*, *G1244E*, *S1251N*, *S1255P* or *G1349D*).

In part 1, patients were randomised 1:1 to receive either 150 mg of ivacaftor or placebo every 12 hours with fat-containing food for 8 weeks in addition to their prescribed CF therapies and crossed over to the other treatment for the second 8 weeks after a 4- to 8-week washout period. The use of inhaled hypertonic saline was not permitted. In part 2, all patients received ivacaftor as indicated in part 1 for 16 additional weeks. The duration of continuous ivacaftor treatment was 24 weeks for patients randomised to part 1 placebo/ivacaftor treatment sequence and 16 weeks for patients randomised to part 1 ivacaftor/placebo treatment sequence.

Thirty-nine patients (mean age 23 years) with baseline $FEV_1 \geq 40\%$ predicted (mean FEV_1 78% predicted [range: 43% to 119%]) were enrolled. Sixty-two percent (24/39) of them carried the *F508del-CFTR* mutation in the second allele. A total of 36 patients continued into part 2 (18 per treatment sequence).

In part 1 of study 770-111, the mean FEV_1 percent predicted at baseline in placebo-treated patients was 79.3% while in ivacaftor-treated patients this value was 76.4%. The mean overall post-baseline value was 76.0% and 83.7%, respectively. The mean absolute change from baseline through week 8 in percent predicted FEV_1 (primary efficacy endpoint) was 7.5% in the ivacaftor period and -3.2% in the placebo period. The observed treatment difference (95% CI) between ivacaftor and placebo was 10.7% (7.3, 14.1) ($P < 0.0001$).

The effect of ivacaftor in the overall population of study 770-111 (including the secondary endpoints absolute change in BMI at 8 weeks of treatment and absolute change in the respiratory domain score of the CFQ-R through 8 weeks of treatment) and by individual mutation (absolute change in sweat chloride and in percent predicted FEV_1 at week 8) is shown in Table 7. Based on clinical (percent predicted FEV_1) and pharmacodynamic (sweat chloride) responses to ivacaftor, efficacy in patients with the *G970R* mutation could not be established.

Table 7: Effect of ivacaftor for efficacy variables in the overall population and for specific *CFTR* mutations

Absolute change in percent predicted FEV_1	BMI (kg/m ²)	CFQ-R respiratory domain score (points)
Through week 8	At week 8	Through week 8
All patients (N = 39)		
Results shown as mean (95% CI) change from baseline ivacaftor vs placebo-treated patients:		
10.7 (7.3, 14.1)	0.66 (0.34, 0.99)	9.6 (4.5, 14.7)

G551S (2)	-68 [†]	3 [†]
G970R [#] (4)	-6 (-16, -2)	3 (-1, 5)
S1251N (8)	-54 (-84, -7)	9 (-20, 21)
S1255P (2)	-78 (-82, -74)	3 (-1, 8)
S549N (6)	-74 (-93, -53)	11 (-2, 20)
S549R (4)	-61 ^{††} (-71, -54)	5 (-3, 13)

* Statistical testing was not performed due to small numbers for individual mutations.

[†] Reflects results from the one patient with the G551S mutation with data at the 8-week time point.

^{††} n = 3 for the analysis of absolute change in sweat chloride.

[#] Causes a splicing defect resulting in little-to-no CFTR protein at the cell surface.

In part 2 of study 770-111, the mean (SD) absolute change in percent predicted FEV₁ following 16 weeks (patients randomised to the ivacaftor/placebo treatment sequence in part 1) of continuous ivacaftor treatment was 10.4% (13.2%). At the follow-up visit, 4 weeks after ivacaftor dosing had ended, the mean (SD) absolute change in percent predicted FEV₁ from part 2 week 16 was -5.9% (9.4%). For patients randomised to the placebo/ivacaftor treatment sequence in part 1 there was a further mean (SD) change of 3.3% (9.3%) in percent predicted FEV₁ after the additional 16 weeks of treatment with ivacaftor. At the follow up visit, 4 weeks after ivacaftor dosing had ended, the mean (SD) absolute change in percent predicted FEV₁ from part 2 week 16 was -7.4% (5.5%).

Study 770-104: study in patients with CF with the F508del mutation in the CFTR gene

Study 770-104 (part A) was a 16-week, 4:1 randomised, double-blind, placebo-controlled, parallel-group phase 2 study of ivacaftor (150 mg every 12 hours) in 140 patients with CF aged 12 years and older who were homozygous for the F508del mutation in the CFTR gene and who had FEV₁ ≥ 40% predicted.

The mean absolute change from baseline through week 16 in percent predicted FEV₁ (primary efficacy endpoint) was 1.5 percentage points in the ivacaftor group and -0.2 percentage points in the placebo group. The estimated treatment difference for ivacaftor versus placebo was 1.7 percentage points (95% CI -0.6, 4.1); this difference was not statistically significant (P = 0.15).

Study 770-105: open-label extension study

In study 770-105 patients who completed treatment in studies 770-102 and 770-103 with placebo were switched to ivacaftor while patients on ivacaftor continued to receive it for a minimum of 96 weeks, i.e., the length of treatment with ivacaftor was at least 96 weeks for patients in the placebo/ivacaftor group and at least 144 weeks for patients in the ivacaftor/ivacaftor group.

One hundred and forty-four (144) patients from study 770-102 were rolled over in study 770-105, 67 in the placebo/ivacaftor group and 77 in the ivacaftor/ivacaftor group. Forty-eight (48) patients from study 770-103 were rolled over in study 770-105, 22 in the placebo/ivacaftor group and 26 in the ivacaftor/ivacaftor group.

Table 8 shows the results of the mean (SD) absolute change in percent predicted FEV₁ for both groups of patients. For patients in the placebo/ivacaftor group baseline percent predicted FEV₁ is that of study 770-105 while for patients in the ivacaftor/ivacaftor group the baseline value is that of studies 770-102 and 770-103.

Table 8: Effect of ivacaftor on percent predicted FEV₁ in study 770-105

Original study and treatment group	Duration of ivacaftor treatment (weeks)	Absolute change from baseline in percent predicted FEV ₁ (percentage points)	
		N	Mean (SD)
Study 770-102			
Ivacaftor	48*	77	9.4 (8.3)
	144	72	9.4 (10.8)
Placebo	0*	67	-1.2 (7.8) [†]
	96	55	9.5 (11.2)
Study 770-103			
Ivacaftor	48*	26	10.2 (15.7)
	144	25	10.3 (12.4)
Placebo	0*	22	-0.6 (10.1) [†]
	96	21	10.5 (11.5)

* Treatment occurred during blinded, controlled, 48-week phase 3 study.

[†] Change from prior study baseline after 48 weeks of placebo treatment.

When the mean (SD) absolute change in percent predicted FEV₁ is compared from study 770-105 baseline for patients in the ivacaftor/ivacaftor group (n = 72) who rolled over from study 770-102, the mean (SD) absolute change in percent predicted FEV₁ was 0.0% (9.05), while for patients in the ivacaftor/ivacaftor group (n = 25) who rolled over from study 770-103 this figure was 0.6% (9.1). This shows that patients in the ivacaftor/ivacaftor group maintained the improvement seen at week 48 of the initial study (day 0 through week 48) in percent predicted FEV₁ through week 144. There were no additional improvements in study 770-105 (week 48 through week 144).

For patients in the placebo/ivacaftor group from study 770-102, the annualised rate of pulmonary exacerbations was higher in the initial study when patients were on placebo (1.34 events/year) than during the subsequent study 770-105 when patients rolled over to ivacaftor (0.48 events/year across day 1 to week 48, and 0.67 events/year across weeks 48 to 96). For patients in the ivacaftor/ivacaftor group from study 770-102, the annualised rate of pulmonary exacerbations was 0.57 events/year across day 1 to week 48 when patients were on ivacaftor. When they rolled over into study 770-105, the rate of annualised pulmonary exacerbations was 0.91 events/year across day 1 to week 48 and 0.77 events/year across weeks 48 to 96.

For patients who rolled over from study 770-103 the number of events was, overall, low.

Study 770-110: study in patients with CF with an R117H mutation in the CFTR gene

Study 770-110 evaluated 69 patients who were 6 years of age or older; 53 (76.8%) patients had the *F508del* mutation in the second allele. The confirmed *R117H* poly-T variant was *5T* in 38 patients and *7T* in 16 patients. At baseline, mean predicted FEV₁ was 73% (range: 32.5% to 105.5%) and mean age was 31 years (range: 6 to 68 years). The mean absolute change from baseline through week 24 in percent predicted FEV₁ (primary efficacy endpoint) was 2.57 percentage points in the ivacaftor group and 0.46 percentage points in the placebo group. The estimated treatment difference for ivacaftor versus placebo was 2.1 percentage points (95% CI -1.1, 5.4).

A pre-planned subgroup analysis was conducted in patients aged 18 years and older (26 patients on placebo and 24 on ivacaftor). Treatment with ivacaftor resulted in a mean absolute change in percent predicted FEV₁ through week 24 of 4.5 percentage points in the ivacaftor group versus -0.46 percentage points in the placebo group. The estimated treatment difference for ivacaftor versus placebo was 5.0 percentage points (95% CI 1.1, 8.8).

In a subgroup analysis in patients with a confirmed *R117H-5T* genetic variant, the difference in the mean absolute change from baseline through week 24 in percent predicted FEV₁ between ivacaftor and placebo was 5.3% (95% CI 1.3, 9.3). In patients with a confirmed *R117H-7T* genetic variant, the treatment difference between ivacaftor and placebo was 0.2% (95% CI -8.1, 8.5).

For secondary efficacy variables, no treatment differences were observed for ivacaftor versus placebo in absolute change from baseline in BMI at week 24 or time to first pulmonary exacerbation.

Treatment differences were observed in absolute change in CFQ-R respiratory domain score through week 24 (treatment difference of ivacaftor versus placebo was 8.4 [95% CI 2.2, 14.6] points) and for the mean change from baseline in sweat chloride (see Pharmacodynamic effects).

5.2 Pharmacokinetic properties

The pharmacokinetics of ivacaftor are similar between healthy adult volunteers and patients with CF.

After oral administration of a single 150 mg dose to healthy volunteers in a fed state, the mean ($\pm$ SD) for AUC and C_{max} were 10600 (5260) ng*hr/mL and 768 (233) ng/mL, respectively. After every 12-hour dosing, steady-state plasma concentrations of ivacaftor were reached by days 3 to 5, with an accumulation ratio ranging from 2.2 to 2.9.

Absorption

Following multiple oral dose administrations of ivacaftor, the exposure of ivacaftor generally increased with dose from 25 mg every 12 hours to 450 mg every 12 hours. When given with fat-containing food, the exposure of

ivacaftor increased approximately 2.5- to 4-fold. Therefore, ivacaftor should be administered with fat-containing food. The median (range) t_{max} is approximately 4.0 (3.0; 6.0) hours in the fed state.

Distribution

Ivacaftor is approximately 99% bound to plasma proteins, primarily to alpha 1-acid glycoprotein and albumin. Ivacaftor does not bind to human red blood cells. After oral administration of ivacaftor 150 mg every 12 hours for 7 days in healthy volunteers in a fed state, the mean ($\pm$ SD) apparent volume of distribution was 353 L (122).

Biotransformation

Ivacaftor is extensively metabolised in humans. *In vitro* and *in vivo* data indicate that ivacaftor is primarily metabolised by CYP3A. M1 and M6 are the two major metabolites of ivacaftor in humans. M1 has approximately one-sixth the potency of ivacaftor and is considered pharmacologically active. M6 has less than one-fiftieth the potency of ivacaftor and is not considered pharmacologically active.

Elimination

Following oral administration in healthy volunteers, the majority of ivacaftor (87.8%) was eliminated in the faeces after metabolic conversion. The major metabolites M1 and M6 accounted for approximately 65% of the total dose eliminated with 22% as M1 and 43% as M6. There was negligible urinary excretion of ivacaftor as unchanged parent. The apparent terminal half-life was approximately 12 hours following a single dose in the fed state. The apparent clearance (CL/F) of ivacaftor was similar for healthy subjects and patients with CF. The mean ($\pm$ SD) CL/F for a single 150 mg dose was 17.3 (8.4) L/hr in healthy subjects.

Linearity/non-linearity

The pharmacokinetics of ivacaftor are generally linear with respect to time or dose ranging from 25 mg to 250 mg.

Special populations

Hepatic impairment

Following a single dose of 150 mg of ivacaftor, adult subjects with moderately impaired hepatic function (Child-Pugh Class B, score 7 to 9) had similar ivacaftor C_{max} (mean [$\pm$ SD] of 735 [331] ng/mL) but an approximately two-fold increase in ivacaftor $AUC_{0-\infty}$ (mean [$\pm$ SD] of 16800 [6140] ng*hr/mL) compared with healthy subjects matched for demographics. Simulations for predicting the steady-state exposure of ivacaftor showed that by reducing the dosage from 150 mg q12h to 150 mg once daily, adults with moderate hepatic impairment would have comparable steady-state C_{min} values as those obtained with a dose of 150 mg q12h in adults without hepatic impairment.

The impact of severe hepatic impairment (Child Pugh Class C, score 10 to 15) on the pharmacokinetics of ivacaftor has not been studied. The magnitude of increase in exposure in these patients is unknown but is expected to be higher than that observed in patients with moderate hepatic impairment.

For guidance on appropriate use and dose modification see Table 3 in section 4.2.

Renal impairment

Pharmacokinetic studies have not been performed with ivacaftor in patients with renal impairment. In a human pharmacokinetic study with ivacaftor monotherapy, there was minimal elimination of ivacaftor and its metabolites in urine (only 6.6% of total radioactivity was recovered in the urine). There was negligible urinary excretion of ivacaftor as unchanged parent (less than 0.01% following a single oral dose of 500 mg).

No dose adjustments are recommended for mild and moderate renal impairment. Caution is recommended when administering ivacaftor to patients with severe renal impairment (creatinine clearance less than or equal to 30

mL/min) or end-stage renal disease (see sections 4.2 and 4.4).

Race

Race had no clinically meaningful effect on the PK of ivacaftor in white (n = 379) and non-white (n = 29) patients based on a population PK analysis.

Gender

The pharmacokinetic parameters of ivacaftor are similar in males and females.

Elderly

Clinical studies of ivacaftor did not include sufficient numbers of patients aged 65 years and older to determine whether pharmacokinetic parameters are similar or not to those in younger adults.

Paediatric population

Predicted ivacaftor exposure based on observed ivacaftor concentrations in phase 2 and 3 studies as determined using compartmental analysis is presented by age group in Table 9.

Table 9: Mean (SD) ivacaftor exposure by age group

Age group	Dose	C _{min, ss} (ng/mL)	AUC _{τ, ss} (ng*h/mL)
1 month to less than 2 months (≥ 3 kg)*	13.4 mg q24h	300 (221) [†]	5840 (2980) [†]
2 months to less than 4 months (≥ 3 kg)*	13.4 mg q24h	406 (266) [†]	6450 (3430) [†]
4 months to less than 6 months (≥ 5 kg)*	25 mg q12h	371 (183)	6480 (2520)
6 months to less than 12 months (≥ 5 kg to < 7 kg) [†]	25 mg q12h	336	5410
6 months to less than 12 months (7 kg to < 14 kg)	50 mg q12h	508 (252)	9140 (4200)
12 months to less than 24 months (7 kg to < 14 kg)	50 mg q12h	440 (212)	9050 (3050)
12 months to less than 24 months (≥ 14 kg to < 25 kg)	75 mg q12h	451 (125)	9600 (1800)
2- to 5-year-olds (< 14 kg)	50 mg q12h	577 (317)	10500 (4260)
2- to 5-year-olds (≥ 14 kg to < 25 kg)	75 mg q12h	629 (296)	11300 (3820)
6- to 11-year-olds [§] (≥ 14 kg to < 25 kg)	75 mg q12h	641 (329)	10760 (4470)
6- to 11-year-olds [§] (≥ 25 kg)	150 mg q12h	958 (546)	15300 (7340)
12- to 17-year-olds	150 mg q12h	564 (242)	9240 (3420)
Adults (≥ 18 years old)	150 mg q12h	701 (317)	10700 (4100)

* Patients 1 month to less than 6 months of age were ≥ 37 weeks gestational age.

[†] Exposures for 1 month to less than 4 months of age are predictions based on simulations from the physiologically based PK model incorporating data from the given age group.

[‡] Values based on data from a single patient; standard deviation not reported.

[§] Exposures in 6- to 11-year-olds are predictions based on simulations from the population PK model using data obtained for this age group.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, and carcinogenic potential.

Pregnancy and fertility

Ivacaftor was associated with slight decreases of the seminal vesicle weights, a decrease of overall fertility index and number of pregnancies in females mated with treated males and significant reductions in number of corpora lutea and implantation sites with subsequent reductions in the average litter size and average number of viable embryos per litter in treated females. The

No-Observed-Adverse-Effect-Level (NOAEL) for fertility findings provides an exposure level of approximately 4 times the systemic exposure of ivacaftor and its metabolites when administered as ivacaftor monotherapy in adult humans at the maximum recommended human dose (MRHD).

Placental transfer of ivacaftor was observed in pregnant rats and rabbits.

Peri- and post-natal development

Ivacaftor decreased survival and lactation indices and caused a reduction in pup body weights. The NOAEL for viability and growth in the offspring provides an exposure level of approximately 3 times the systemic exposure of ivacaftor and its metabolites when administered as ivacaftor monotherapy in adult humans at the MRHD.

Juvenile animals studies

Findings of cataracts were observed in juvenile rats dosed from postnatal day 7 through 35 at ivacaftor exposure levels of 0.22 times the MRHD based on systemic exposure of ivacaftor and its metabolites when administered as ivacaftor monotherapy. This finding has not been observed in fetuses derived from rat dams treated with ivacaftor on gestation days 7 to 17, in rat pups exposed to ivacaftor through milk ingestion up to postnatal day 20, in 7-week old rats, nor in 3.5 to 5-month old dogs treated with ivacaftor. The potential relevance of these findings in humans is unknown.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Core

Hypromellose acetate succinate

Sodium laurilsulfate (E487)

Silicified microcrystalline cellulose

(comprises of microcrystalline cellulose and silica, colloidal anhydrous)

Mannitol (E421)

Croscarmellose sodium

Magnesium stearate

Coating

Polyvinyl alcohol part. hydrolyzed (E1203)

Calcium carbonate (E170)

Macrogol (E1521)

Talc (E553b)

Indigo carmine aluminum lake (E132)

Carnauba wax powder

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

OPA/Aluminum/PVC//Aluminum blisters or transparent PVC/PCTFE(Aclar)//Aluminum blisters

The following pack sizes are available:

Blister packs containing 28 & 56 film-coated tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

[To be completed nationally]

8. MARKETING AUTHORISATION NUMBER(S)

[To be completed nationally]

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

[To be completed nationally]

10. DATE OF REVISION OF THE TEXT

08 May 2025