

Package leaflet: Information for the user

Venlafaxin 1A Farma 37.5 mg prolonged-release tablets

Venlafaxin 1A Farma 75 mg prolonged-release tablets

Venlafaxin 1A Farma 150 mg prolonged-release tablets

Venlafaxin 1A Farma 225 mg prolonged-release tablets

venlafaxine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Venlafaxin 1A Farma is and what it is used for
2. What you need to know before you take Venlafaxin 1A Farma
3. How to take Venlafaxin 1A Farma
4. Possible side effects
5. How to store Venlafaxin 1A Farma
6. Contents of the pack and other information

1. What Venlafaxin 1A Farma is and what it is used for

Venlafaxin 1A Farma contains the active substance venlafaxine.

Venlafaxin 1A Farma is an antidepressant that belongs to a group of medicines called serotonin and norepinephrine reuptake inhibitors (SNRIs). This group of medicines is used to treat depression and other conditions, such as anxiety disorders. It is not fully understood how antidepressants work, but they may help by increasing the levels of serotonin and noradrenaline in the brain.

Venlafaxin 1A Farma is a treatment for adults with depression. Venlafaxin 1A Farma is also a treatment for adults with the following anxiety disorders: generalised anxiety disorder, social anxiety disorder (fear or avoidance of social situations) and panic disorder (panic attacks). Treating depression or anxiety disorders properly is important to help you get better. If it is not treated, your condition may not go away and may become more serious and more difficult to treat.

2. What you need to know before you take Venlafaxin 1A Farma

Do not take Venlafaxin 1A Farma

- If you are allergic to venlafaxine or any of the other ingredients of this medicine (listed in section 6).
- If you are also taking or have taken within the last 14 days any medicines known as irreversible monoamine oxidase inhibitors (MAOIs), used to treat depression or Parkinson's disease. Taking an irreversible MAOI together with Venlafaxin 1A Farma, can cause serious or even life-threatening side effects. Also, you must wait at least 7 days after you stop taking Venlafaxin 1A Farma before you take any MAOI (see also the section entitled "Other medicines and Venlafaxin 1A Farma" and the information in that section about "Serotonin Syndrome").

Warnings and precautions

Talk to your doctor or pharmacist **before** taking Venlafaxin 1A Farma:

- If you use other medicines that taken together with Venlafaxin 1A Farma could increase the risk of developing serotonin syndrome (see the section “Other medicines and Venlafaxin 1A Farma ”).
- If you have a swallowing, stomach or intestinal disorder that reduces your ability to swallow or pass foods by normal bowel movements
- If you have eye problems, such as certain kinds of glaucoma (increased pressure in the eye).
- If you have a history of high blood pressure.
- If you have a history of heart problems.
- If you have been told you have an abnormal heart rhythm.
- If you have a history of fits (seizures).
- If you have a history of low sodium levels in your blood (hyponatraemia).
- If you have a history of bleeding disorders (tendency to develop bruises or a tendency to bleed easily) or if you are taking other medicines that may increase the risk of bleeding e.g. warfarin (used to prevent blood clots) or if you are pregnant (see ‘Pregnancy and breast-feeding’)
- If you have a history of, or if someone in your family has had, mania or bipolar disorder (feeling over-excited or euphoric).
- If you have a history of aggressive behaviour.

Venlafaxin 1A Farma may cause a sensation of restlessness or an inability to sit or stand still during the first few weeks of treatment. You should tell your doctor if this happens to you.

Do not drink alcohol while being treated with Venlafaxin 1A Farma as it can lead to extreme tiredness and unconsciousness. Concomitant use with alcohol and/or certain medicines can make your symptoms of depression and other conditions, such as anxiety disorders worse.

Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders, you can sometimes have thoughts of harming or killing yourself. These may be increased when you first start taking antidepressants, since these medicines all take time to work, usually about two weeks, but sometimes longer. These thoughts may also occur when your dose is decreased or during discontinuation of treatment with Venlafaxin 1A Farma.

You may be more likely to think like this:

- If you have previously had thoughts about killing yourself or harming yourself.
- If you are a young adult. Information from clinical trials has shown an increased risk of suicidal behaviour in young adults (less than 25 years old) with psychiatric conditions who were treated with an antidepressant.

If you have thoughts of harming or killing yourself at any time, contact your doctor or go to a hospital straight away.

You may find it helpful to tell a relative or close friend that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

Dry mouth

Dry mouth is reported in 10% of patients treated with venlafaxine. This may increase the risk of tooth decay (caries). Therefore, you should take special care in your dental hygiene.

Diabetes

Your blood glucose levels may be altered due to Venlafaxin 1A Farma. Therefore, the dosages of our diabetes medicines may need to be adjusted.

Sexual problems

Medicines like Venlafaxin 1A Farma (so called SSRIs/SNRIs) may cause symptoms of sexual dysfunction (see section 4). In some cases, these symptoms have continued after stopping treatment.

Children and adolescents

Venlafaxin 1A Farma should normally not be used for children and adolescents under 18 years. Also, you should know that patients under 18 have an increased risk of side effects, such as suicide attempt, suicidal thoughts and hostility (predominantly aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe this medicine for patients under 18 because he/she decides that this is in their best interests. If your doctor has prescribed this medicine for a patient under 18, and you want to discuss this, please go back to your doctor. You should inform your doctor if any of the symptoms listed above develop or worsen when patients under 18 are taking Venlafaxin 1A Farma. Also, the long-term safety effects concerning growth, maturation and cognitive and behavioural development of this medicine in this age group has not yet been demonstrated.

Other medicines and Venlafaxin 1A Farma

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Your doctor should decide whether you can take Venlafaxin 1A Farma with other medicines.

Do not start or stop taking any medicines, including those bought without a prescription, natural and herbal remedies, before checking with your doctor or pharmacist.

- Monoamine oxidase inhibitors which are used to treat depression or Parkinson's disease **must not be taken with Venlafaxin 1A Farma**. Tell your doctor if you have taken these medicines within the last 14 days. (MAOIs: see the section "What you need to know before you take Venlafaxin 1A Farma").

- **Serotonin syndrome:**

A potentially life-threatening condition or Neuroleptic Malignant Syndrome (NMS)-like reactions (see the section "Possible Side Effects"), may occur with venlafaxine treatment, particularly when taken with other medicines.

Examples of these medicines include:

- Triptans (used for migraine)
- Other medicines to treat depression, for instance SNRI, SSRIs, tricyclics, or medicines containing lithium
- Medicines containing amphetamines (used to treat attention deficit hyperactivity disorder (ADHD), narcolepsy and obesity)
- Medicines containing linezolid, an antibiotic (used to treat infections)
- Medicines containing moclobemide, a MAOI (used to treat depression)
- Medicines containing sibutramine (used for weight loss)
- Medicines containing opioids (e.g., buprenorphine, tramadol, fentanyl, tapentadol, pethidine, or pentazocine) (used to treat severe pain)
- Medicines containing dextromethorphan (used to treat coughing)
- Medicines containing methadone (used to treat opioid drug addiction or severe pain)
- Medicines containing methylene blue (used to treat high levels of methaemoglobin in the blood)
- Products containing St. John's Wort (also called *Hypericum perforatum*, a natural or herbal remedy used to treat mild depression)
- Products containing tryptophan (used for problems such as sleep and depression)
- Antipsychotics (used to treat a disease with symptoms such as hearing, seeing or sensing things which are not there, mistaken beliefs, unusual suspiciousness, unclear reasoning and becoming withdrawn)

Signs and symptoms of serotonin syndrome may include a combination of the following: restlessness, hallucinations, loss of coordination, fast heartbeat, increased body temperature, fast changes in blood pressure, overactive reflexes, diarrhoea, coma, nausea, vomiting.

In its most severe form, serotonin syndrome can resemble Neuroleptic Malignant Syndrome (NMS). Signs and symptoms of NMS may include a combination of fever, fast heartbeat, sweating, severe muscle stiffness, confusion, increased muscle enzymes (determined by a blood test).

Tell your doctor immediately or go to the casualty department at your nearest hospital if you think serotonin syndrome is happening to you.

You must tell your doctor if you are taking medicines that can affect your heart rhythm.

Examples of these medicines include:

- Antiarrhythmics such as quinidine, amiodarone, sotalol or dofetilide (used to treat abnormal heart rhythm)
- Antipsychotics such as thioridazine (See also Serotonin syndrome above)
- Antibiotics such as erythromycin or moxifloxacin (used to treat bacterial infections)
- Antihistamines (used to treat allergy)

The following medicines may also interact with Venlafaxin 1A Farma and should be used with caution. It is especially important to mention to your doctor or pharmacist if you are taking medicines containing:

- Ketoconazole (an antifungal medicine)
- Haloperidol or risperidone (to treat psychiatric conditions)
- Metoprolol (a beta blocker to treat high blood pressure and heart problems)

Venlafaxin 1A Farma with food, drink and alcohol

Venlafaxin 1A Farma should be taken with food (see section 3 “How to take Venlafaxin 1A Farma”).

Do not drink alcohol while being treated with Venlafaxin 1A Farma. Concomitant use with alcohol can lead to extreme tiredness and unconsciousness and can make your symptoms of depression and other conditions, such as anxiety disorders worse.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. You should use Venlafaxin 1A Farma only after discussing the potential benefits and the potential risks to your unborn child with your doctor.

Make sure your midwife and/or doctor knows you are on Venlafaxin 1A Farma. When taken during pregnancy, similar drugs (SSRIs) may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear bluish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

If you take Venlafaxin 1A Farma near the end of your pregnancy there may be an increased risk of heavy vaginal bleeding shortly after birth, especially if you have a history of bleeding disorders. Your doctor or midwife should be aware that you are taking Venlafaxin 1A Farma so they can advise you.

If you are taking this medicine during pregnancy, in addition to having trouble breathing, another symptom your baby might have when it is born is not feeding properly. If your baby has these symptoms when it is born and you are concerned, contact your doctor and/or midwife who will be able to advise you.

Venlafaxin 1A Farma passes into breast milk. There is a risk of an effect on the baby. Therefore, you should discuss the matter with your doctor, and he/she will decide whether you should stop breast-feeding or stop the therapy with this medicine.

Driving and using machines

Do not drive or use any tools or machines until you know how this medicine affects you.

Venlafaxin 1A Farma contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Venlafaxin 1A Farma

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual recommended starting dose for treatment of depression, generalised anxiety disorder and social anxiety disorder is 75 mg per day. The dose can be raised by your doctor gradually, and if needed, even up to a maximum dose of 375 mg daily for depression. If you are being treated for panic disorder, your doctor will start with a lower dose (37.5 mg) and then increase the dose gradually. The maximum dose for generalised anxiety disorder, social anxiety disorder and panic disorder is 225 mg/day.

Take Venlafaxin 1A Farma at approximately the same time each day, either in the morning or in the evening. Tablets must be swallowed whole with fluid and not opened, crushed, chewed or dissolved.

Venlafaxin 1A Farma should be taken with food.

If you have liver or kidney problems, talk to your doctor, since your dose of this medicine may need to be different.

Do not stop taking this medicine without talking to your doctor (see the section “If you stop taking Venlafaxin 1A Farma”).

If you take more Venlafaxin 1A Farma than you should

Call your doctor or pharmacist immediately if you take more of this medicine than the amount prescribed by your doctor.

Overdose can be life-threatening, especially with concomitant use of alcohol and/or certain medicines (see “Other medicines and Venlafaxin 1A Farma”).

The symptoms of a possible overdose may include a rapid heartbeat, changes in level of alertness (ranging from sleepiness to coma), blurred vision, seizures or fits, and vomiting.

If you forget to take Venlafaxin 1A Farma

If you miss a dose, take it as soon as you remember. However, if it is time for your next dose, skip the missed dose and take only a single dose as usual. Do not take a double dose to make up for a forgotten dose. Do not take more than the daily amount of Venlafaxin 1A Farma that has been prescribed for you in one day.

If you stop taking Venlafaxin 1A Farma

Do not stop taking your treatment or reduce the dose without the advice of your doctor even if you feel better. If your doctor thinks that you no longer need Venlafaxin 1A Farma, he/she may ask you to reduce your dose slowly, before stopping treatment altogether. Side effects are known to occur when people stop using this medicine, especially when it is stopped suddenly or the dose is reduced too quickly. Some patients may experience symptoms such as suicidal thoughts, aggressiveness, tiredness, dizziness, light-headedness, headache, sleeplessness, nightmares, dry mouth, loss of appetite, nausea, diarrhoea, nervousness, agitation, confusion, ringing in the ears, tingling or rarely electric shock sensations, weakness, sweating, seizures, or flu-like symptoms problems with eyesight and increase in blood pressure (which can cause headache, dizziness, ringing in the ears, sweating, etc.)

Your doctor will advise you on how you should gradually discontinue Venlafaxin 1A Farma treatment. This can take a period of several weeks or months. In some patients, discontinuation may need to occur very gradually over periods of months or longer. If you experience any of these or other symptoms that are troublesome, ask your doctor for further advice.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If any of the following happen, do not take more Venlafaxin 1A Farma. **Tell your doctor immediately, or go to the casualty department at your nearest hospital:**

Uncommon (may affect up to 1 in 100 people)

- Swelling of the face, mouth, tongue, throat, hands, or feet, and/or a raised itchy rash (hives), trouble swallowing or breathing

Rare (may affect up to 1 in 1,000 people)

- Chest tightness, wheezing, trouble swallowing or breathing
- Severe skin rash, itching, or hives (elevated patches of red or pale skin that often itch)
- Signs and symptoms of serotonin syndrome which may include restlessness, hallucinations, loss of coordination, fast heartbeat, increased body temperature, fast changes in blood pressure, overactive reflexes, diarrhoea, coma, nausea, vomiting.
In its most severe form, serotonin syndrome can resemble Neuroleptic Malignant Syndrome (NMS). Signs and symptoms of NMS may include a combination of fever, fast heartbeat, sweating, severe muscle stiffness, confusion, increased muscle enzymes (determined by a blood test).
- Signs of infection, such as high temperature, chills, shivering, headaches, sweating, flu-like symptoms. This may be the result of a blood disorder which leads to an increased risk of infection.
- Severe rash, which may lead to severe blistering and peeling of the skin.
- Unexplained muscle pain, tenderness or weakness. This may be a sign of rhabdomyolysis.

Frequency not known (cannot be estimated from the available data)

- Signs and symptoms of a condition called “stress cardiomyopathy” which may include chest pain, shortness of breath, dizziness, fainting, irregular heartbeat.

Other side effects that you should **tell your doctor about include** (The frequency of these side effects are included in the list “Other side effects that may occur” below):

- Coughing, wheezing and, shortness of breath which may be accompanied by a high temperature
- Black (tarry) stools or blood in stools
- Itchiness, yellow skin or eyes, or dark urine, which may be symptoms of inflammation of the liver (hepatitis)
- Heart problems, such as fast or irregular heart rate, increased blood pressure
- Eye problems, such as blurred vision, dilated pupils
- Nerve problems, such as dizziness, pins and needles, movement disorder (muscle spasms or stiffness), seizures or fits
- Psychiatric problems, such as hyperactivity and feeling unusually overexcited
- Withdrawal effects (see the section “How to take Venlafaxin 1A Farma, if you stop taking Venlafaxin 1A Farma”).

- Prolonged bleeding - if you cut or injure yourself, it may take slightly longer than usual for bleeding to stop.

Do not be concerned if you see a tablet in your stools after taking this medicine. As the tablet travels the length of your gastrointestinal tract, venlafaxine is slowly released. The shape of the tablet remains undissolved and is eliminated in your stools. Therefore, even though you may see a tablet in your stools, your dose of venlafaxine has been absorbed.

Other side effects that may occur

Very common (may affect more than 1 in 10 people)

- Dizziness; headache; drowsiness
- Insomnia
- Nausea; dry mouth; constipation
- Sweating (including night sweats)

Common (may affect up to 1 in 10 people)

- Appetite decreased
- Confusion; feeling separated (or detached) from yourself; lack of orgasm; decreased libido; agitation; nervousness; abnormal dreams
- Tremor, a sensation of restlessness or an inability to sit or stand still; pins and needles; altered taste sensation; increased muscle tone
- Visual disturbance including blurred vision; dilated pupils; inability of the eye to automatically change focus from distant to near objects
- Ringing in the ears (tinnitus)
- Fast heartbeat; palpitations
- Increase in blood pressure; flushing
- Shortness of breath; yawning
- Vomiting; diarrhoea
- Mild rash; itching
- Increased frequency in urination; inability to pass urine; difficulties passing urine
- Menstrual irregularities such as increased bleeding or increased irregular bleeding; abnormal ejaculation/orgasm (males); erectile dysfunction (impotence)
- Weakness (asthenia); fatigue; chills
- Weight gain; weight loss
- Increased cholesterol

Uncommon (may affect up to 1 in 100 people)

- Over activity, racing thoughts and decreased need for sleep (mania)
- Hallucinations; feeling separated (or detached) from reality; abnormal orgasm; lack of feeling or emotion; feeling over-excited; grinding of the teeth
- Fainting; involuntary movements of the muscles; impaired coordination and balance
- Feeling dizzy (particularly when standing up too quickly); decrease in blood pressure
- Vomiting blood, black tarry stools (faeces) or blood in stools; which can be a sign of internal bleeding
- Sensitivity to sunlight; bruising; abnormal hair loss
- Inability to control urination
- Stiffness, spasms and involuntary movements of the muscles
- Slight changes in blood levels of liver enzymes

Rare (may affect up to 1 in 1,000 people)

- Seizures or fits
- Coughing, wheezing and shortness of breath which may be accompanied by a high temperature
- Disorientation and confusion often accompanied by hallucination (delirium)

- Excessive water intake (known as SIADH)
- Decrease in blood sodium levels
- Severe eye pain and decreased or blurred vision
- Abnormal, rapid or irregular heartbeat, which could lead to fainting
- Severe abdominal or back pains (which could indicate a serious problem in the gut, liver or pancreas)
- Itchiness, yellow skin or eyes, dark urine, or flu-like symptoms, which are symptoms of inflammation of the liver (hepatitis)

Very rare (may affect up to 1 in 10,000 people)

- Prolonged bleeding, which may be a sign of reduced number of platelets in your blood, leading to an increased risk of bruising or bleeding
- Abnormal breast milk production
- Unexpected bleeding, e.g. bleeding gums, blood in the urine or in vomit, or the appearance of unexpected bruises or broken blood vessels (broken veins)

Frequency not known (frequency cannot be estimated from the available data)

- Suicidal ideation and suicidal behaviours; cases of suicidal ideation and suicidal behaviours have been reported during venlafaxine therapy or early after treatment discontinuation (see section 2, what you need to know before you take Venlafaxin 1A Farma)
- Aggression
- Vertigo
- Heavy vaginal bleeding shortly after birth (postpartum haemorrhage), see 'Pregnancy and breast-feeding' in section 2 for more information.

Venlafaxin 1A Farma sometimes causes unwanted effects that you may not be aware of, such as increases in blood pressure or abnormal heartbeat; slight changes in blood levels or liver enzymes, sodium or cholesterol. More rarely, Venlafaxin 1A Farma may reduce the function of platelets in your blood, leading to an increased risk of bruising or bleeding. Therefore, your doctor may wish to do blood tests occasionally, particularly if you have been taking Venlafaxin 1A Farma for a long time.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Venlafaxin 1A Farma

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the packaging. The expiry date refers to the last day of that month.

This medicinal product does not require any special temperature storage conditions.

Blisters: Store in the original package in order to protect from moisture.

Plastic bottle: Keep the bottle tightly closed in order to protect from moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What **Venlafaxin 1A Farma** contains
The active substance is venlafaxine.

Each prolonged-release tablet contains 37.5, 75, 150 or 225 mg venlafaxine (as hydrochloride).

The other ingredients are

Core: Mannitol (E421), Povidone K-90, Macrogol 400, Cellulose microcrystalline (E 460(i)), Colloidal anhydrous silica, Magnesium stearate (E 470b).

Coat: Cellulose acetate, Macrogol 400, hypromellose, lactose monohydrate, titanium dioxide (E171), triacetin.

What Venlafaxin 1A Farma looks like and contents of the pack

37.5 mg prolonged-release tablets: 7 mm round, biconvex, white to off white coloured tablets with a (little) pierce in one side.

75 mg prolonged-release tablets: 7.5 mm round, biconvex, white to off white coloured tablets with a (little) pierce in one side.

150 mg prolonged-release tablets: 9.5 mm round, biconvex, white to off white coloured tablets with a (little) pierce in one side.

225 mg prolonged-release tablets: 11 mm round, biconvex, white to off white coloured tablets with a (little) pierce in one side.

Venlafaxin 1A Farma is available in blisters of 10, 14, 20, 28, 30, 50, 56, 60, 100 and 500 tablets; and in plastic bottles of 10, 14, 20, 28, 30, 50, 56, 60, 100, 106 and 500 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

<To be completed nationally>

This medicinal product is authorised in the Member States of the European Economic Area and in the United Kingdom (Northern Ireland) under the following names:

Name of the Member States	Name of the medicinal product
Sweden	Venlafaxin 1A Farma 37,5/75/150/225 mg depottabletter
Estonia	Venlafaxine Wörwag Pharma 37,5/75/150/225 mg, toimeainet prolongeeritult vabastavad tabletid
Greece	SEROSMINE 37.5/75/150/225 mg Δισκίο παρατεταμένης αποδέσμευσης
France	VENLAFAXINE BIPHAR 150/225 mg comprimé a liberation prolongée
Italy	Zarelix 37.5/75/150/225 mg compressa a rilascio prolungato
Netherlands	Venlafaxine Aurobindo 225 mg Tablet met verlengde afgifte
Poland	Axyven, 37,5/75/150 mg, tabletki o przedłużonym uwalnianiu
Portugal	Zarelix 37,5/75/150/225 mg comprimidos de libertação prolongada
United kingdom	Venlafaxine XL 37.5/75/150/225 mg prolonged-release tablets

This leaflet was last approved in 16 October 2025

Other sources of information

Detailed information on this medicine is available on the website of {name of Member State Agency (link)}.