

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Zyfoqam 5 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 5 mg of linagliptin.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Red, round, biconvex film-coated tablets with a nominal diameter of 8.1 mm, debossed with “5” on one side and plain on the other side.

4. CLINICAL PARTICULARS

4.1. Therapeutic indications

[Product Name] is indicated in adults with type 2 diabetes mellitus as an adjunct to diet and exercise to improve glycaemic control as:

monotherapy

- when metformin is inappropriate due to intolerance, or contraindicated due to renal impairment.

combination therapy

- in combination with other medicinal products for the treatment of diabetes, including insulin, when these do not provide adequate glycaemic control (see sections 4.4, 4.5 and 5.1 for available data on different combinations).

4.2. Posology and method of administration

Posology

The dose of linagliptin is 5 mg once daily. When linagliptin is added to metformin, the dose of metformin should be maintained, and linagliptin administered concomitantly.

When linagliptin is used in combination with a sulphonylurea or with insulin, a lower dose of the sulphonylurea or insulin, may be considered to reduce the risk of hypoglycaemia (see section 4.4)

Special populations

Renal impairment

For patients with renal impairment, no dose adjustment for linagliptin is required.

Hepatic impairment

Pharmacokinetic studies suggest that no dose adjustment is required for patients with hepatic impairment but clinical experience in such patients is lacking.

Elderly

No dose adjustment is necessary based on age.

Paediatric population

A clinical trial did not establish efficacy in paediatric patients 10 to 17 years of age (see section 4.8, 5.1 and 5.2). Therefore, treatment of children and adolescents with linagliptin is not recommended. Linagliptin has not been studied in paediatric patients under 10 years of age.

Method of administration

The tablets can be taken with or without a meal at any time of the day. If a dose is missed, it should be taken as soon as the patient remembers. A double dose should not be taken on the same day.

4.3. Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4. Special warnings and precautions for use

General

Linagliptin should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis.

Hypoglycaemia

Linagliptin alone showed a comparable incidence of hypoglycaemia to placebo.

In clinical trials of linagliptin as part of combination therapy with medicinal products not known to cause hypoglycaemia (metformin), rates of hypoglycaemia reported with linagliptin were similar to rates in patients taking placebo.

When linagliptin was added to a sulphonylurea (on a background of metformin), the incidence of hypoglycaemia was increased over that of placebo (see section 4.8).

Sulphonylureas and insulin are known to cause hypoglycaemia. Therefore, caution is advised when linagliptin is used in combination with a sulphonylurea and/or insulin. A dose reduction of the sulphonylurea or insulin may be considered (see section 4.2).

Acute pancreatitis

Use of DPP-4 inhibitors has been associated with a risk of developing acute pancreatitis. Acute pancreatitis has been observed in patients taking linagliptin. In a cardiovascular and renal safety study (CARMELINA) with median observation period of 2.2 years, adjudicated acute pancreatitis was reported in 0.3% of patients treated with linagliptin and in 0.1% of patients treated with placebo.

Patients should be informed of the characteristic symptoms of acute pancreatitis. If pancreatitis is suspected, [Product Name] should be discontinued; if acute pancreatitis is confirmed, [Product Name] should not be restarted. Caution should be exercised in patients with a history of pancreatitis.

Bullous pemphigoid

Bullous pemphigoid has been observed in patients taking linagliptin. In the CARMELINA study, bullous pemphigoid was reported in 0.2% of patients on treatment with linagliptin and in no patient on placebo. If bullous pemphigoid is suspected, [Product Name] should be discontinued.

4.5. Interaction with other medicinal products and other forms of interaction

In vitro assessment of interactions

Linagliptin is a weak competitive and a weak to moderate mechanism-based inhibitor of CYP isozyme CYP3A4, but does not inhibit other CYP isozymes. It is not an inducer of CYP isozymes.

Linagliptin is a P-glycoprotein substrate, and inhibits P-glycoprotein mediated transport of digoxin with low potency. Based on these results and *in vivo* interaction studies, linagliptin is considered unlikely to cause interactions with other P-gp substrates.

In vivo assessment of interactions

Effects of other medicinal products on linagliptin

Clinical data described below suggest that the risk for clinically meaningful interactions by co-administered medicinal products is low.

Rifampicin: multiple co-administration of 5 mg linagliptin with rifampicin, a potent inducer of P-glycoprotein and CYP3A4, resulted in a 39.6% and 43.8% decreased linagliptin steady-state AUC and C_{max} , respectively, and about 30% decreased DPP-4 inhibition at trough. Thus, full efficacy of linagliptin in combination with strong P-gp inducers might not be achieved, particularly if these are administered long-term. Co-administration with other potent inducers of P-glycoprotein and CYP3A4, such as carbamazepine, phenobarbital and phenytoin has not been studied.

Ritonavir: co-administration of a single 5 mg oral dose of linagliptin and multiple 200 mg oral doses of ritonavir, a potent inhibitor of P-glycoprotein and CYP3A4, increased the AUC and C_{max} of linagliptin approximately twofold and threefold, respectively. The unbound concentrations, which are usually less than 1% at the therapeutic dose of linagliptin, were increased 4-5-fold after co-administration with ritonavir. Simulations of steady-state plasma concentrations of linagliptin with and without ritonavir indicated that the increase in exposure will be not associated with an increased accumulation. These changes in linagliptin pharmacokinetics were not considered to be clinically relevant. Therefore, clinically relevant interactions would not be expected with other P-glycoprotein/CYP3A4 inhibitors.

Metformin: co-administration of multiple three times daily doses of 850 mg metformin with 10 mg linagliptin once daily did not clinical meaningfully alter the pharmacokinetics of linagliptin in healthy volunteers.

Sulphonylureas: the steady-state pharmacokinetics of 5 mg linagliptin was not changed by concomitant administration of a single 1.75 mg dose glibenclamide (glyburide).

Effects of linagliptin on other medicinal products

In clinical studies, as described below, linagliptin had no clinically relevant effect on the pharmacokinetics of metformin, glyburide, simvastatin, warfarin, digoxin or oral contraceptives providing *in vivo* evidence of a low propensity for causing medicinal product interactions with substrates of CYP3A4, CYP2C9, CYP2C8, P-glycoprotein, and organic cationic transporter (OCT).

Metformin: co-administration of multiple daily doses of 10 mg linagliptin with 850 mg metformin, an OCT substrate, had no relevant effect on the pharmacokinetics of metformin in healthy volunteers. Therefore, linagliptin is not an inhibitor of OCT-mediated transport.

Sulphonylureas: co-administration of multiple oral doses of 5 mg linagliptin and a single oral dose of 1.75 mg glibenclamide (glyburide) resulted in clinically not relevant reduction of 14% of both AUC and C_{max} of glibenclamide. Because glibenclamide is primarily metabolised by CYP2C9, these data also support the conclusion that linagliptin is not a CYP2C9 inhibitor. Clinically meaningful interactions would not be expected with other sulphonylureas (e.g., glipizide, tolbutamide, and glimepiride) which, like glibenclamide, are primarily eliminated by CYP2C9.

Digoxin: co-administration of multiple daily doses of 5 mg linagliptin with multiple doses of 0.25 mg digoxin had no effect on the pharmacokinetics of digoxin in healthy volunteers. Therefore, linagliptin is not an inhibitor of P-glycoprotein-mediated transport *in vivo*.

Warfarin: multiple daily doses of 5 mg linagliptin did not alter the pharmacokinetics of S(-) or R(+) warfarin, a CYP2C9 substrate, administered in a single dose.

Simvastatin: multiple daily doses of linagliptin had a minimal effect on the steady-state pharmacokinetics of simvastatin, a sensitive CYP3A4 substrate, in healthy volunteers. Following administration of a supratherapeutic dose of 10 mg linagliptin concomitantly with 40 mg of simvastatin daily for 6 days, the plasma AUC of simvastatin was increased by 34%, and the plasma C_{max} by 10%.

Oral contraceptives: co-administration with 5 mg linagliptin did not alter the steady-state pharmacokinetics of levonorgestrel or ethinylestradiol.

4.6. Fertility, pregnancy and lactation

Pregnancy

The use of linagliptin has not been studied in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). As a precautionary measure, it is preferable to avoid the use of linagliptin during pregnancy.

Breastfeeding

Available pharmacokinetic data in animals have shown excretion of linagliptin/metabolites in milk. A risk to the breastfed child cannot be excluded. A decision must be made whether to discontinue breastfeeding or to discontinue/abstain from linagliptin therapy taking into account the benefit of breastfeeding for the child and the benefit of therapy for the woman.

Fertility

No studies on the effect on human fertility have been conducted for linagliptin. Animal studies do not indicate direct or indirect harmful effects with respect to fertility (see section 5.3).

4.7. Effects on ability to drive and use machines

Linagliptin has no or negligible influence on the ability to drive and use machines. However patients should be alerted to the risk of hypoglycaemia especially when combined with sulphonylurea and/or insulin.

4.8. Undesirable effects

Summary of the safety profile

In the pooled analysis of the placebo-controlled trials, the overall incidence of adverse events in patients treated with placebo was similar to linagliptin 5 mg (63.4% versus 59.1%).

Discontinuation of therapy due to adverse events was higher in patients who received placebo as compared to linagliptin 5 mg (4.3% versus 3.4%).

The most frequently reported adverse reaction was “hypoglycaemia” observed under the triple combination, linagliptin plus metformin plus sulphonylurea 14.8% versus 7.6% in placebo.

In the placebo-controlled studies 4.9% of patients experienced “hypoglycaemia” as an adverse reaction under linagliptin. Of these, 4.0% were mild and 0.9% were moderate and 0.1% were classified as severe in intensity. Pancreatitis was reported more often in patients randomized to linagliptin (7 events in 6,580 patients receiving linagliptin versus 2 events in 4,383 patients receiving placebo).

Tabulated list of adverse reactions

Due to the impact of the background therapy on adverse reactions (e.g. on hypoglycaemias), adverse reactions were analysed based on the respective treatment regimens (monotherapy, add-on to metformin, add-on to metformin plus sulphonylurea, and add-on to insulin).

The placebo-controlled studies included studies where linagliptin was given as

- monotherapy with short-term duration of up to 4 weeks
- monotherapy with ≥ 12 week duration
- add-on to metformin
- add-on to metformin + sulphonylurea
- add on to metformin and empagliflozin
- add-on to insulin with or without metformin

Adverse reactions classified by system organ class and MedDRA preferred terms reported in patients who received 5 mg linagliptin in double-blind studies as monotherapy or as add-on therapy are presented in the table below (see Table 1).

The adverse reactions are listed by absolute frequency. Frequencies are defined as very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1\,000$ to $< 1/100$), rare ($\geq 1/10\,000$ to $< 1/1\,000$), very rare ($< 1/10\,000$) or not known (cannot be estimated from the available data).

Table 1. Adverse reactions reported in patients who received linagliptin 5 mg daily as monotherapy or as add-on therapies in clinical trial and from post-marketing experience.

System organ class Adverse reaction	Frequency of adverse reaction
Infections and infestations	
Nasopharyngitis	uncommon
Immune system disorders	
Hypersensitivity (e.g. bronchial hyperreactivity)	uncommon
Metabolism and nutrition disorders	
Hypoglycaemia ¹	very common
Respiratory, thoracic and mediastinal disorders	
Cough	uncommon
Gastrointestinal disorders	
Pancreatitis	rare#
Constipation ²	uncommon
Skin and subcutaneous tissue disorders	
Angioedema*	rare
Urticaria*	rare
Rash*	uncommon
Bullous pemphigoid	rare#
Investigations	
Amylase increased	uncommon
Lipase increased**	common

* Based on post-marketing experience

** Based on lipase elevations $> 3 \times \text{ULN}$ observed in clinical trials

Based on Linagliptin cardiovascular and renal safety study (CARMELINA), see also below

¹ Adverse reaction observed in combination with metformin plus sulphonylurea

² Adverse reaction observed in combination with insulin

Linagliptin cardiovascular and renal safety study (CARMELINA)

The CARMELINA study evaluated the cardiovascular and renal safety of linagliptin versus placebo in patients with type 2 diabetes and with increased CV risk evidenced by a history of established macrovascular or renal disease (see section 5.1). The study included 3494 patients treated with linagliptin (5 mg) and 3485 patients treated with placebo. Both treatments were added to standard of care targeting regional standards for HbA1c and CV risk factors. The overall incidence of adverse events

and serious adverse events in patients receiving linagliptin was similar to that in patients receiving placebo. Safety data from this study was in line with previous known safety profile of linagliptin.

In the treated population, severe hypoglycaemic events (requiring assistance) were reported in 3.0% of patients on linagliptin and in 3.1% on placebo. Among patients who were using sulfonylurea at baseline, the incidence of severe hypoglycaemia was 2.0% in linagliptin-treated patients and 1.7% in placebo treated patients. Among patients who were using insulin at baseline, the incidence of severe hypoglycaemia was 4.4% in linagliptin-treated patients and 4.9% in placebo treated patients.

In the overall study observation period adjudicated acute pancreatitis was reported in 0.3% of patients treated with linagliptin and in 0.1% of patients treated with placebo.

In the CARMELINA study, bullous pemphigoid was reported in 0.2% of patients treated with linagliptin and in no patient treated with placebo.

Paediatric population

Overall, in clinical trials in paediatric patients with type 2 diabetes mellitus aged 10 to 17 years, the safety profile of linagliptin was similar to that observed in the adult population.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

4.9. Overdose

Symptoms

During controlled clinical trials in healthy subjects, single doses of up to 600 mg linagliptin (equivalent to 120 times the recommended dose) were generally well tolerated. There is no experience with doses above 600 mg in humans.

Therapy

In the event of an overdose, it is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring and institute clinical measures if required.

5. PHARMACOLOGICAL PROPERTIES

5.1. Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes, dipeptidyl peptidase 4 (DPP-4) inhibitors, ATC code: A10BH05

Mechanism of action

Linagliptin is an inhibitor of the enzyme DPP-4 (dipeptidyl peptidase 4, EC 3.4.14.5) an enzyme which is involved in the inactivation of the incretin hormones GLP-1 and GIP (glucagon-like peptide1, glucose-dependent insulintropic polypeptide). These hormones are rapidly degraded by the enzyme DPP-4. Both incretin hormones are involved in the physiological regulation of glucose homeostasis. Incretins are secreted at a low basal level throughout the day and levels rise immediately after meal intake. GLP-1 and GIP increase insulin biosynthesis and secretion from pancreatic beta cells in the presence of normal and elevated blood glucose levels. Furthermore GLP-1 also reduces glucagon secretion from pancreatic alpha cells, resulting in a reduction in hepatic glucose output. Linagliptin binds very effectively to DPP-4 in a reversible manner and thus leads to a sustained increase and a prolongation of active incretin levels. Linagliptin glucose-dependently increases insulin secretion and lowers

glucagon secretion thus resulting in an overall improvement in the glucose homeostasis. Linagliptin binds selectively to DPP-4 and exhibits a > 10,000 fold selectivity versus DPP-8 or DPP-9 activity in vitro.

Clinical efficacy and safety

8 phase III randomised controlled trials involving 5,239 patients with type 2 diabetes, of which 3,319 were treated with linagliptin were conducted to evaluate efficacy and safety. These studies had 929 patients of 65 years and over who were on linagliptin. There were also 1,238 patients with mild renal impairment, and 143 patients with moderate renal impairment on linagliptin. Linagliptin once daily produced clinically significant improvements in glycaemic control, with no clinically relevant change in body weight. The reductions in glycosylated haemoglobin A1c (HbA1c) were similar across different subgroups including gender, age, renal impairment and body mass index (BMI). Higher baseline HbA1c was associated with a greater reduction in HbA1c. There was a significant difference in reduction in HbA1c between Asian patients (0.8%) and White patients (0.5%) in the pooled studies.

Linagliptin as monotherapy in patients ineligible for metformin

The efficacy and safety of linagliptin monotherapy was evaluated in a double-blind placebo-controlled study of 24 weeks duration. Treatment with once daily linagliptin at 5 mg provided a significant improvement in HbA1c (-0.69% change compared to placebo), in patients with baseline HbA1c of approximately 8%. Linagliptin also showed significant improvements in fasting plasma glucose (FPG), and 2-hour post-prandial glucose (PPG) compared to placebo. The observed incidence of hypoglycaemia in patients treated with linagliptin was similar to placebo.

The efficacy and safety of linagliptin monotherapy was also evaluated in patients for whom metformin therapy is inappropriate, due to intolerance or contraindicated due to renal impairment, in a double-blind placebo-controlled study of 18 weeks duration. Linagliptin provided significant improvements in HbA1c, (-0.57% change compared to placebo), from a mean baseline HbA1c of 8.09%. Linagliptin also showed significant improvements in fasting plasma glucose (FPG) compared to placebo. The observed incidence of hypoglycaemia in patients treated with linagliptin was similar to placebo.

Linagliptin as add-on to metformin therapy

The efficacy and safety of linagliptin in combination with metformin was evaluated in a double-blind placebo-controlled study of 24 weeks duration. Linagliptin provided significant improvements in HbA1c, (-0.64% change compared to placebo), from a mean baseline HbA1c of 8%. Linagliptin also showed significant improvements in fasting plasma glucose (FPG), and 2-hour post-prandial glucose (PPG) compared to placebo. The observed incidence of hypoglycaemia in patients treated with linagliptin was similar to placebo.

Linagliptin as add-on to a combination of metformin and sulphonylurea therapy

A placebo-controlled study of 24 weeks in duration was conducted to evaluate the efficacy and safety of linagliptin 5 mg to placebo, in patients not sufficiently treated with a combination with metformin and a sulphonylurea. Linagliptin provided significant improvements in HbA1c (-0.62% change compared to placebo), from a mean baseline HbA1c of 8.14%. Linagliptin also showed significant improvements in patients fasting plasma glucose (FPG), and 2-hour post-prandial glucose (PPG), compared to placebo.

Linagliptin as add-on to a combination of metformin and empagliflozin therapy

In patients inadequately controlled with metformin and empagliflozin (10 mg (n=247) or 25 mg (n=217)), 24-weeks treatment with add-on therapy of linagliptin 5 mg provided adjusted mean HbA1c reductions from baseline by -0.53% (significant difference to add-on placebo -0.32% (95% CI -0.52, -0.13) and -0.58% (significant difference to add-on placebo -0.47% (95% CI -0.66; -0.28), respectively. A statistically significant greater proportion of patients with a baseline HbA1c $\geq 7.0\%$ and treated with linagliptin 5 mg achieved a target HbA1c of $<7\%$ compared to placebo.

Linagliptin as add-on to insulin therapy

The efficacy and safety of the addition of linagliptin 5 mg to insulin alone or in combination with metformin and/or pioglitazone has been evaluated in a double-blind placebo-controlled study of 24 weeks duration. Linagliptin provided significant improvements in HbA1c (-0.65% compared to placebo) from a mean baseline HbA1c of 8.3%. Linagliptin also provided significant improvements in fasting plasma glucose (FPG), and a greater proportion of patients achieved a target HbA1c of < 7.0%, compared to placebo. This was achieved with a stable insulin dose (40.1 IU). Body weight did not differ significantly between the groups. Effects on plasma lipids were negligible. The observed incidence of hypoglycaemia in patients treated with linagliptin was similar to placebo (22.2% linagliptin; 21.2% placebo).

Linagliptin 24 month data, as add-on to metformin in comparison with glimepiride

In a study comparing the efficacy and safety of the addition of linagliptin 5 mg or glimepiride (mean dose 3 mg) in patients with inadequate glycaemic control on metformin monotherapy, mean reductions in HbA1c were -0.16% with linagliptin (mean baseline HbA1c 7.69%) and -0.36% with glimepiride (mean baseline HbA1c 7.69%) with a mean treatment difference of 0.20% (97.5% CI: 0.09, 0.299). The incidence of hypoglycaemia in the linagliptin group (7.5%) was significantly lower than that in the glimepiride group (36.1%). Patients treated with linagliptin exhibited a significant mean decrease from baseline in body weight compared to a significant weight gain in patients administered glimepiride (-1.39 vs +1.29 kg).

Linagliptin as add-on therapy in patients with severe renal impairment, 12 week placebo-controlled data (stable background) and 40 week placebo-controlled extension (adjustable background)

The efficacy and safety of linagliptin was also evaluated in type 2 diabetes patients with severe renal impairment in a double-blind study versus placebo for 12 weeks duration, during which background glycaemic therapies were kept stable. Most patients (80.5%) received insulin as background therapy, alone or in combination with other oral anti-diabetics such as sulphonylurea, glinide and pioglitazone. There was a further follow up 40 week treatment period during which dose adjustments in antidiabetes background therapies were allowed.

Linagliptin provided significant improvements in HbA1c (-0.59 % change compared to placebo after 12 weeks), from a mean baseline HbA1c of 8.2%. The observed difference in HbA1c over placebo was -0.72% after 52 weeks.

Body weight did not differ significantly between the groups. The observed incidence of hypoglycaemia in patients treated with linagliptin was higher than placebo, due to an increase in asymptomatic hypoglycaemic events. There was no difference between groups in severe hypoglycaemic events.

Linagliptin as add-on therapy in elderly (age ≥ 70 years) with type 2 diabetes

The efficacy and safety of linagliptin in elderly (age ≥ 70years) with type 2 diabetes was evaluated in a double-blind study of 24 weeks duration. Patients received metformin and/or sulphonylurea and/or insulin as background therapy. Doses of background antidiabetic medicinal products were kept stable during the first 12 weeks, after which adjustments were permitted. Linagliptin provided significant improvements in HbA1c (-0.64 % change compared to placebo after 24 weeks), from a mean baseline HbA1c of 7.8%. Linagliptin also showed significant improvements in fasting plasma glucose (FPG) compared to placebo. Body weight did not differ significantly between the groups.

Linagliptin cardiovascular and renal safety study (CARMELINA)

CARMELINA was a randomized study in 6979 patients with type 2 diabetes with increased CV risk evidenced by a history of established macrovascular or renal disease who were treated with linagliptin 5 mg (3494) or placebo (3485) added to standard of care targeting regional standards for HbA1c, CV risk factors and renal disease. The study population included 1211 (17.4%) patients ≥ 75 years of age and 4348 (62.3%) patients with renal impairment. Approximately 19% of the population had eGFR ≥45 to <60 mL/min/1.73 m², 28% of the population had eGFR ≥30 to <45 mL/min/1.73 m² and 15% had eGFR < 30 mL/min/1.73 m². The mean HbA1c at baseline was 8.0%. The study was designed to demonstrate non-inferiority for the primary cardiovascular endpoint which was a composite of the first

occurrence of cardiovascular death or a non-fatal myocardial infarction (MI) or a non-fatal stroke (3P-MACE). The renal composite endpoint was defined as renal death or sustained end stage renal disease or sustained decrease of 40% or more in eGFR.

After a median follow up of 2.2 years, linagliptin, when added to usual care, did not increase the risk of major adverse cardiovascular events or renal outcome events. There was no increased risk in hospitalization for heart failure which was an additional adjudicated endpoint observed compared to usual care without linagliptin in patients with type 2 diabetes (see Table 2).

Table 2. Cardiovascular and renal outcomes by treatment group in the CARMELINA study

	Linagliptin 5 mg		Placebo		Hazard Ratio
	Number of Subjects (%)	Incidence Rate per 1000 PY*	Number of Subjects (%)	Incidence Rate per 1000 PY*	(95% CI)
Number of patients	3494		3485		
Primary CV composite (Cardiovascular death, non-fatal MI, non-fatal stroke)	434 (12.4)	57.7	420 (12.1)	56.3	1.02 (0.89, 1.17)**
Secondary renal composite (renal death, ESRD, 40% sustained decrease in eGFR)	327 (9.4)	48.9	306 (8.8)	46.6	1.04 (0.89, 1.22)
All-cause mortality	367 (10.5)	46.9	373 (10.7)	48.0	0.98 (0.84, 1.13)
CV death	255 (7.3)	32.6	264 (7.6)	34	0.96 (0.81, 1.14)
Hospitalization for heart failure	209 (6.0)	27.7	226 (6.5)	30.4	0.90 (0.74, 1.08)

* PY=patient years

** Test on non-inferiority to demonstrate that the upper bound of the 95% CI for the hazard ratio is less than 1.3

In analyses for albuminuria progression (change from normoalbuminuria to micro- or macroalbuminuria, or from microalbuminuria to macroalbuminuria) the estimated hazard ratio was 0.86 (95% CI 0.78, 0.95) for linagliptin versus placebo.

Linagliptin cardiovascular safety study (CAROLINA)

CAROLINA was a randomized study in 6033 patients with early type 2 diabetes and increased CV risk or established complications who were treated with linagliptin 5 mg (3023) or glimepiride 1-4mg (3010) added to standard of care (including background therapy with metformin in 83% of patients) targeting regional standards for HbA1c and CV risk factors. The mean age for study population was 64 years and included 2030 (34%) patients $\geq$ 70 years of age. The study population included 2089 (35%) patients with cardiovascular disease and 1130 (19%) patients with renal impairment with an eGFR < 60ml/min/1.73m² at baseline. The mean HbA1c at baseline was 7.15%.

The study was designed to demonstrate non-inferiority for the primary cardiovascular endpoint which was a composite of the first occurrence of cardiovascular death or a non-fatal myocardial infarction (MI) or a non-fatal stroke (3P-MACE).

After a median follow up of 6.25 years, linagliptin did not increase the risk of major adverse cardiovascular events (see Table 3) as compared to glimepiride. Results were consistent for patients treated with or without metformin.

Table 3. Major adverse cardiovascular events (MACE) and mortality by treatment group in the CAROLINA study

	Linagliptin 5 mg		Glimepiride (1-4mg)		Hazard Ratio
	Number of Subjects (%)	Incidence Rate per 1000 PY*	Number of Subjects (%)	Incidence Rate per 1000 PY*	(95% CI)
Number of patients	3023		3010		
Primary CV composite (Cardiovascular death, non-fatal MI, non-fatal stroke)	356 (11.8)	20.7	362 (12.0)	21.2	0.98 (0.84, 1.14)**
All-cause mortality	308 (10.2)	16.8	336 (11.2)	18.4	0.91 (0.78, 1.06)
CV death	169 (5.6)	9.2	168 (5.6)	9.2	1.00 (0.81, 1.24)
Hospitalization for heart failure (HHF)	112 (3.7)	6.4	92 (3.1)	5.3	1.21 (0.92, 1.59)

* PY=patient years

** Test on non-inferiority to demonstrate that the upper bound of the 95% CI for the hazard ratio is less than 1.3

For the entire treatment period (median time on treatment 5.9 years) the rate of patients with moderate or severe hypoglycaemia was 6.5% on linagliptin versus 30.9% on glimepiride, severe hypoglycaemia occurred in 0.3% of patients on linagliptin versus 2.2% on glimepiride.

Paediatric population

The clinical efficacy and safety of empagliflozin 10 mg with potential dose-increase to 25 mg or linagliptin 5 mg once daily has been studied in children and adolescents from 10 to 17 years of age with T2DM in a double-blind, randomised, placebo-controlled, parallel group study (DINAMO) over 26 weeks, with a double-blind active treatment safety extension period up to 52 weeks.

At baseline, the mean HbA1c was 8.03%. Treatment with linagliptin 5 mg did not provide significant improvement in HbA1c. The treatment difference of adjusted mean change in HbA1c after 26 weeks between linagliptin and placebo was -0.34% (95% CI -0.99, 0.30; p=0.2935). The adjusted mean change in HbA1c from baseline was 0.33% in patients treated with linagliptin and 0.68% in patients treated with placebo (see section 4.2).

5.2. Pharmacokinetic properties

The pharmacokinetics of linagliptin has been extensively characterised in healthy subjects and patients with type 2 diabetes. After oral administration of a 5 mg dose to healthy volunteers or patients, linagliptin was rapidly absorbed, with peak plasma concentrations (median T_{max}) occurring 1.5 hours post-dose.

Plasma concentrations of linagliptin decline in a triphasic manner with a long terminal half-life (terminal half-life for linagliptin more than 100 hours), that is mostly related to the saturable, tight binding of linagliptin to DPP-4 and does not contribute to the accumulation of the medicinal product. The effective half-life for accumulation of linagliptin, as determined from oral administration of multiple doses of 5 mg linagliptin, is approximately 12 hours. After once daily dosing of 5 mg linagliptin, steady-state plasma concentrations are reached by the third dose. Plasma AUC of linagliptin increased approximately 33% following 5 mg doses at steady-state compared to the first dose. The intra-subject and inter-subject coefficients of variation for linagliptin AUC were small (12.6% and 28.5%, respectively). Due to the concentration dependent binding of linagliptin to DPP-4, the pharmacokinetics of linagliptin based on total exposure is not linear; indeed total plasma AUC of linagliptin increased in a less than dose-proportional manner while unbound AUC increases in a roughly dose-proportional manner. The pharmacokinetics of linagliptin was generally similar in healthy subjects and in patients with type 2 diabetes.

Absorption

The absolute bioavailability of linagliptin is approximately 30%. Co-administration of a high-fat meal with linagliptin prolonged the time to reach C_{max} by 2 hours and lowered C_{max} by 15% but no influence on AUC_{0-72h} was observed. No clinically relevant effect of C_{max} and T_{max} changes is expected; therefore linagliptin may be administered with or without food.

Distribution

As a result of tissue binding, the mean apparent volume of distribution at steady-state following a single 5 mg intravenous dose of linagliptin to healthy subjects is approximately 1,110 litres, indicating that linagliptin extensively distributes to the tissues. Plasma protein binding of linagliptin is concentration-dependent, decreasing from about 99% at 1 nmol/l to 75-89% at ≥ 30 nmol/l, reflecting saturation of binding to DPP-4 with increasing concentration of linagliptin. At high concentrations, where DPP-4 is fully saturated, 70-80% of linagliptin was bound to other plasma proteins than DPP-4, hence 30-20% were unbound in plasma.

Biotransformation

Following a [^{14}C] linagliptin oral 10 mg dose, approximately 5% of the radioactivity was excreted in urine. Metabolism plays a subordinate role in the elimination of linagliptin. One main metabolite with a relative exposure of 13.3% of linagliptin at steady-state was detected which was found to be pharmacologically inactive and thus does not contribute to the plasma DPP-4 inhibitory activity of linagliptin.

Elimination

Following administration of an oral [^{14}C] linagliptin dose to healthy subjects, approximately 85% of the administered radioactivity was eliminated in faeces (80%) or urine (5%) within 4 days of dosing. Renal clearance at steady-state was approximately 70 ml/min.

Special populations

Renal impairment

A multiple-dose, open-label study was conducted to evaluate the pharmacokinetics of linagliptin (5 mg dose) in patients with varying degrees of chronic renal insufficiency compared to normal healthy control subjects. The study included patients with renal insufficiency classified on the basis of creatinine clearance as mild (50 to <80 ml/min), moderate (30 to <50 ml/min), and severe (<30 ml/min), as well as patients with ESRD on hemodialysis. In addition patients with T2DM and severe renal impairment (<30 ml/min) were compared to T2DM patients with normal renal function. Creatinine clearance was measured by 24-hour urinary creatinine clearance measurements or estimated from serum creatinine based on the Cockcroft-Gault formula. $CrCl = (140 - \text{age}) \times \text{weight}/72 \times \text{serum creatinine}$ [$\times 0.85$ for females], where age is in years, weight in kg, and serum creatinine is in mg/dl.

Under steady-state conditions, linagliptin exposure in patients with mild renal impairment was comparable to healthy subjects. In moderate renal impairment, a moderate increase in exposure of about 1.7 fold was observed compared with control. Exposure in T2DM patients with severe RI was increased by about 1.4 fold compared to T2DM patients with normal renal function. Steady-state predictions for AUC of linagliptin in patients with ESRD indicated comparable exposure to that of patients with moderate or severe renal impairment. In addition, linagliptin is not expected to be eliminated to a therapeutically significant degree by hemodialysis or peritoneal dialysis. Therefore, no dosage adjustment of linagliptin is necessary in patients with any degree of renal insufficiency.

Hepatic impairment

In non-diabetic patients with mild moderate and severe hepatic insufficiency (according to the Child-Pugh classification), mean AUC and $C_{\max}$ of linagliptin were similar to healthy matched controls following administration of multiple 5 mg doses of linagliptin. No dosage adjustment for linagliptin is proposed for diabetic patients with mild, moderate or severe hepatic impairment.

Body Mass Index (BMI)

No dosage adjustment is necessary based on BMI. BMI had no clinically relevant effect on the pharmacokinetics of linagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data. The clinical trials before marketing authorisation have been performed up to a BMI equal to 40 kg/m².

Gender

No dosage adjustment is necessary based on gender. Gender had no clinically relevant effect on the pharmacokinetics of linagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data.

Elderly

No dosage adjustment is required based on age up to 80 years, as age did not have a clinically relevant impact on the pharmacokinetics of linagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data. Older subjects (65 to 80, oldest patient was 78 years) had comparable plasma concentrations of linagliptin compared to younger subjects.

Paediatric population

A paediatric Phase 2 study examined the pharmacokinetics and pharmacodynamics of 1 mg and 5 mg linagliptin in children and adolescents ≥ 10 to <18 years of age with type 2 diabetes mellitus. The observed pharmacokinetic and pharmacodynamic responses were consistent with those found in adult subjects. Linagliptin 5 mg showed superiority over 1 mg with regard to trough DPP-4 inhibition (72% vs 32%, $p=0.0050$) and a numerically larger reduction with regard to adjusted mean change from baseline in HbA1c (-0.63% vs -0.48%, n.s.). Due to the limited nature of the data set the results should be interpreted cautiously.

A paediatric Phase 3 study examined pharmacokinetics and pharmacodynamics (HbA1c change from baseline) of 5 mg linagliptin in children and adolescents 10 to 17 years of age with type 2 diabetes mellitus. The observed exposure-response relationship was generally comparable between paediatric and adult patients, however, with a smaller drug effect estimated in children. Oral administration of linagliptin resulted in exposure within the range observed in adult patients. The observed geometric mean trough concentrations and geometric mean concentrations at 1.5 hours post-administration (representing a concentration around $t_{\max}$) at steady state were 4.30 nmol/L and 12.6 nmol/L, respectively. Corresponding plasma concentrations in adult patients were 6.04 nmol/L and 15.1 nmol/L.

Race

No dosage adjustment is necessary based on race. Race had no obvious effect on the plasma concentrations of linagliptin based on a composite analysis of available pharmacokinetic data, including patients of Caucasian, Hispanic, African, and Asian origin. In addition the pharmacokinetic

characteristics of linagliptin were found to be similar in dedicated phase I studies in Japanese, Chinese and Caucasian healthy volunteers.

5.3. Preclinical safety data

Liver, kidneys and gastrointestinal tract are the principal target organs of toxicity in mice and rats at repeat doses of linagliptin of more than 300 times the human exposure.

In rats effects on reproductive organs, thyroid and the lymphoid organs were seen at more than 1,500 times human exposure. Strong pseudo-allergic reactions were observed in dogs at medium doses, secondarily causing cardiovascular changes, which were considered dog-specific. Liver, kidneys, stomach, reproductive organs, thymus, spleen, and lymph nodes were target organs of toxicity in Cynomolgus monkeys at more than 450 times human exposure. At more than 100 times human exposure, irritation of the stomach was the major finding in these monkeys.

Linagliptin and its main metabolite did not show a genotoxic potential.

Oral 2 year carcinogenicity studies in rats and mice revealed no evidence of carcinogenicity in rats or male mice. A significantly higher incidence of malignant lymphomas only in female mice at the highest dose (> 200 times human exposure) is not considered relevant for humans (explanation: non-treatment related but due to highly variable background incidence). Based on these studies there is no concern for carcinogenicity in humans.

The NOAEL for fertility, early embryonic development and teratogenicity in rats was set at >900 times the human exposure. The NOAEL for maternal-, embryo-fetal-, and offspring toxicity in rats was 49 times human exposure. No teratogenic effects were observed in rabbits at >1,000 times human exposure. A NOAEL of 78 times human exposure was derived for embryo-fetal toxicity in rabbits, and for maternal toxicity the NOAEL was 2.1 times human exposure. Therefore, it is considered unlikely that linagliptin affects reproduction at therapeutic exposures in humans.

6. PHARMACEUTICAL PARTICULARS

6.1. List of excipients

Tablet core

Mannitol (E421)

(Maize) starch, pregelatinised

Maize starch

Povidone (E1201)

Magnesium stearate (E470b)

Film coating

Hypromellose (E464)

Titanium dioxide (E171)

Macrogol (E1521)

Iron oxide red (E172)

6.2. Incompatibilities

Not applicable.

6.3. Shelf life

3 years

6.4. Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5. Nature and contents of container

oPA/aluminium/PVC//aluminium blisters in cartons containing 10, 28, 30, 90 film-coated tablets.
Perforated oPA/aluminium/PVC//aluminium unit-dose blisters in cartons containing 10 x 1, 28 x 1, 30 x 1, 90 x 1 film-coated tablets.

Not all pack sizes may be marketed.

6.6. Special precautions for disposal

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

<To be completed nationally>

8. MARKETING AUTHORISATION NUMBER(S)

<To be completed nationally>

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: <To be completed nationally>

Date of latest renewal: <To be completed nationally>

10. DATE OF REVISION OF THE TEXT

2025-12-18